arXiv:1006.1485v4 [math.AP] 25 Aug 2010

Blowup and Scattering problems for the Nonlinear Schrödinger equations

Takafumi Akahori and Hayato Nawa

Abstract

We consider L^2-supercritical and H^1-subcritical focusing nonlinear Schrödinger equations. We introduce a subset PW of $H^1(\mathbb{R}^d)$ for $d \geq 1$, and investigate behavior of the solutions with initial data in this set. For this end, we divide PW into two disjoint components PW_+ and PW_-. Then, it turns out that any solution starting from a datum in PW_+ behaves asymptotically free, and solution starting from a datum in PW_- blows up or grows up, from which we find that the ground state has two unstable directions. We also investigate some properties of generic global and blowup solutions.

1 Introduction

In this paper, we consider the Cauchy problem for the nonlinear Schrödinger equation

$$2i\frac{\partial\psi}{\partial t}(x,t) + \Delta\psi(x,t) + |\psi(x,t)|^{p-1}\psi(x,t) = 0, \quad (x,t) \in \mathbb{R}^d \times \mathbb{R}, \tag{1.1}$$

where $i := \sqrt{-1}$, ψ is a complex-valued function on $\mathbb{R}^d \times \mathbb{R}$, Δ is the Laplace operator on $\mathbb{R}^d$ and p satisfies the so-called L^2-supercritical and H^1-subcritical condition

$$2 + \frac{4}{d} < p+1 < 2^* := \begin{cases} \infty & \text{if} \quad d = 1,2, \\ \dfrac{2d}{d-2} & \text{if} \quad d \geq 3. \end{cases} \tag{1.2}$$

We associate this equation with the initial datum from the usual Sobolev space $H^1(\mathbb{R}^d)$:

$$\psi(\cdot,0) = \psi_0 \in H^1(\mathbb{R}^d). \tag{1.3}$$

We summarize the basic properties of this Cauchy problem (1.1) and (1.3) (see, e.g., [12, 21, 26, 27, 28, 53]). The unique local existence of solutions is well known: for any $\psi_0 \in H^1(\mathbb{R}^d)$, there exists a unique solution ψ in $C(I_{\max}; H^1(\mathbb{R}^d))$ for some interval $I_{\max} = (-T_{\max}^-, T_{\max}^+) \subset \mathbb{R}$: maximal existence interval including 0; $T_{\max}^+$ $(-T_{\max}^-)$ is the maximal existence time for the future (the past). If $I_{\max} \subsetneqq \mathbb{R}$, then we have

$$\lim_{t \to *T_{\max}^*} \|\nabla\psi(t)\|_{L^2} = \infty \quad \text{(blowup)}, \tag{1.4}$$

provided that $T_{\max}^* < \infty$, where $*$ stands for $+$ or $-$. Besides, the solution ψ satisfies the following conservation laws of the mass $\mathcal{M}$, the Hamiltonian $\mathcal{H}$ and the momentum $\mathcal{P}$ in this order: for all $t \in I_{\max}$,

$$\mathcal{M}(\psi(t)) := \|\psi(t)\|_{L^2}^2 = \mathcal{M}(\psi_0), \tag{1.5}$$

$$\mathcal{H}(\psi(t)) := \|\nabla\psi(t)\|_{L^2}^2 - \frac{2}{p+1}\|\psi(t)\|_{L^{p+1}}^{p+1} = \mathcal{H}(\psi_0), \tag{1.6}$$

$$\mathcal{P}(\psi(t)) := \Im\int_{\mathbb{R}^d} \nabla\psi(x,t)\overline{\psi(x,t)}\,dx = \mathcal{P}(\psi_0). \tag{1.7}$$

If, in addition, $\psi_0 \in L^2(\mathbb{R}^d, |x|^2dx)$, then the corresponding solution ψ also belongs to $C(I_{\max}; L^2(\mathbb{R}^d, |x|^2dx))$ and satisfies the so-called virial identity (see [21]):

$$\begin{split}\int_{\mathbb{R}^d} |x|^2\,|\psi(x,t)|^2\,dx = \int_{\mathbb{R}^d} |x|^2\,|\psi_0(x)|^2\,dx + 2t\Im\int_{\mathbb{R}^d} x\cdot\nabla\psi_0(x)\overline{\psi_0(x)}\,dx \\ + 2\int_0^t\int_0^{t'} \mathcal{K}(\psi(t''))\,dt''dt' \qquad \text{for all } t \in I_{\max},\end{split} \tag{1.8}$$

where $\mathcal{K}$ is a functional defined by

$$\mathcal{K}(f) := \|\nabla f\|_{L^2}^2 - \frac{d(p-1)}{2(p+1)}\|f\|_{L^{p+1}}^{p+1}, \quad f \in H^1(\mathbb{R}^d). \tag{1.9}$$

It is worth while noting that

$$\mathcal{K}(f) = \mathcal{H}(f) - \frac{d}{2(p+1)}\left\{p - \left(1+\frac{4}{d}\right)\right\}\|f\|_{L^{p+1}}^{p+1}, \quad f \in H^1(\mathbb{R}^d), \tag{1.10}$$

so that: for any $f \in H^1(\mathbb{R}^d)\setminus\{0\}$, we have

$$\mathcal{K}(f) > \mathcal{H}(f) \quad \text{if} \quad p < 1+\frac{4}{d}, \tag{1.11}$$

$$\mathcal{K}(f) = \mathcal{H}(f) \quad \text{if} \quad p = 1+\frac{4}{d}, \tag{1.12}$$

$$\mathcal{K}(f) < \mathcal{H}(f) \quad \text{if} \quad p > 1+\frac{4}{d}. \tag{1.13}$$

The virial identity (1.8) tells us the behavior of the "variance" of solution, from which we expect to obtain a kind of propagation or concentration estimates. However, we can not use (1.8) as it is, since we do not require the weight condition $\psi_0 \in L^2(\mathbb{R}^d, |x|^2dx)$. We will work in the pure energy space $H^1(\mathbb{R}^d)$, introducing a generalized version of the virial identity (see (A.20) in the appended section A) to discuss propagation or concentration of a "solution" in Section 4.1 and Section 5.

In some physical literature, nonlinear Schrödinger equations, abbreviated to NLSs, (or Gross-Pitaevskii equation) arise as NLSs with linear potentials:

$$2i\frac{\partial\psi}{\partial t}(x,t) + \Delta\psi(x,t) + |\psi(x,t)|^{p-1}\psi(x,t) = V(x)\psi(x,t), \quad (x,t) \in \mathbb{R}^d\times\mathbb{R}, \tag{1.14}$$

where V is a real potential on $\mathbb{R}^d$. The case $p = d = 3$ corresponds to the conventional ϕ^4-model for Bose gasses with negative scattering length (see, e.g., [33]). Typical examples of the potential V are: $V(x) = c_h|x|^2$ (harmonic potentials with $c_h > 0$); and

$V(x) = g \cdot x$ (the Stark potentials or uniform gravitational fields along $g \in \mathbb{R}^d \setminus \{0\}$). However, it is known that these potentials can be "removed" by appropriate space-time transformations: For $V = g \cdot x$, the Avrov-Herbst formula works well (see [11, 15]); For $V = c_h|x|^2$, the space-time transformations in [10, 47] do. Therefore, the study of NLSs without potentials may reveal the essential structure of the solutions to NLSs with potentials (1.14).

Our equation (1.1) has several kinds of solutions: standing waves, blowup solutions (see (1.4) above), and global-in-time solutions which asymptotically behave like free solutions in the distant future/ distant past. Here, the standing waves are nontrivial solutions to our equation (1.1) of the form

$$\psi(x,t) = e^{\frac{i}{2}\omega t} Q(x), \qquad \omega > 0, \quad Q \in H^1(\mathbb{R}^d) \setminus \{0\}. \tag{1.15}$$

Thus, Q solves the following semilinear elliptic equation (nonlinear scalar field equation):

$$\Delta Q - \omega Q + |Q|^{p-1} Q = 0, \qquad \omega > 0, \quad Q \in H^1(\mathbb{R}^d) \setminus \{0\}. \tag{1.16}$$

Here, we remark that every solution Q to the equation (1.16) satisfies $\mathcal{K}(Q) = 0$. Indeed; since any solution Q to (1.16) belongs to the space $H^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d, |x|^2 dx)$, the standing wave $\psi = e^{\frac{i}{2}\omega t} Q$ enjoys the virial identity (1.8), which immediately leads us to $\mathcal{K}(Q) = 0$.

The standing waves are one of the interesting objects in the study of NLSs for both mathematics and physics: standing waves are considered to be the states of Bose-Einstein condensations. In this paper, we are interested in precise instability mechanism of what we call the ground state (the least action solution to (1.16), see also (1.18) below). For this end, we employ the classical "potential well" theory traced back to Sattinger [50]. To define our potential well PW, we need to know some variational properties of the ground state. We shall give the precise definition of PW in (1.35) below. Anyway, our PW is divided into PW_+ and PW_- according to the sign of the functional $\mathcal{K}$, i.e., $PW_+ = PW \cap [\mathcal{K} > 0]$, $PW_- = PW \cap [\mathcal{K} < 0]$ (see (1.37), (1.38)), and the ground state belongs to $\overline{PW_+} \cap \overline{PW_-}$. We shall show that any solution starting from PW_+ exists globally in time and asymptotically behaves like a free solution in the distant future and past (see Theorem 1.1); in contrast, PW_- gives rise to "singular" solutions (see Theorem 1.2). Thus, the ground state shows at least two type of instability, since it belongs to $\overline{PW_+} \cap \overline{PW_-}$.

In order to define our potential well PW, we shall investigate some properties of the ground states here. There are many literature concerning the elliptic equation (1.16) (see, e.g., [4, 19, 32, 51]). We know that if $d \geq 2$, there are infinitely many solutions (bound states) Q_ω^n ($n = 1, 2, \dots$) such that

$$\mathcal{S}_\omega(Q_\omega^n) := \frac{1}{2} \|\nabla Q_\omega^n\|_{L^2}^2 + \frac{\omega}{2} \|Q_\omega^n\|_{L^2}^2 - \frac{1}{p+1} \|Q_\omega^n\|_{L^{p+1}}^{p+1} \to \infty \quad (n \to \infty), \tag{1.17}$$

where the functional $\mathcal{S}_\omega$ is called the action for (1.16) (see, e.g., [4, 51]). The ground state Q_ω is the least action solution to (1.16), more strongly, it satisfies that

$$\mathcal{S}_\omega(Q_\omega) = \inf \left\{ \mathcal{S}_\omega(Q) \mid Q \in H^1(\mathbb{R}^d) \setminus \{0\}, \ \mathcal{K}(Q) = 0 \right\}. \tag{1.18}$$

In the L^2-supercritical case ($p > 1 + \frac{4}{d}$), it turns out that the ground state solves the following variational problems (see Proposition 1.2 below):

$$N_1 := \inf\Big\{\|f\|_{\widetilde{H}^1}^2 \;\big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\},\ \mathcal{K}(f)\le 0\Big\}, \tag{1.19}$$

$$N_2 := \inf\Big\{\mathcal{N}_2(f) \;\big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\},\ \mathcal{K}(f)\le 0\Big\}, \tag{1.20}$$

$$N_3 := \inf\Big\{\mathcal{I}(f) \;\big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\}\Big\}, \tag{1.21}$$

where

$$\|f\|_{\widetilde{H}^1}^2 := \frac{d(p-1)-4}{d(p-1)}\|\nabla f\|_{L^2}^2 + \|f\|_{L^2}^2, \tag{1.22}$$

$$\mathcal{N}_2(f) := \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\,\|\nabla f\|_{L^2}^{\frac{d}{2}(p-1)-2}, \tag{1.23}$$

$$\mathcal{I}(f) := \frac{\|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla f\|_{L^2}^{\frac{d}{2}(p-1)}}{\|f\|_{L^{p+1}}^{p+1}}. \tag{1.24}$$

We remark that N_3 is positive and gives the best constant of the Gagliardo-Nirenberg inequality, i.e.,

$$\|f\|_{L^{p+1}}^{p+1} \le \frac{1}{N_3}\|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla f\|_{L^2}^{\frac{d}{2}(p-1)} \quad \text{for all } f \in H^1(\mathbb{R}^d) \tag{1.25}$$

(for the details, see [55] and the exposition by Tao [54]). A salient point of our approach to these variational problems (1.19)–(1.21) is that we consider them at the same time, which enables us to obtain some identities automatically (see (1.32) and (1.33) below). The first variational problem (1.19) is introduced to make our variational problems easy to solve. Indeed, all minimizing sequences for (1.19) are bounded in $H^1(\mathbb{R}^d)$; In contrast, minimizing sequences in the other problems (1.20) and (1.21) are not necessarily bounded in $H^1(\mathbb{R}^d)$, this fact comes from the invariance of the functionals $\mathcal{K}$, $\mathcal{N}_2$ and $\mathcal{I}$ under the scaling

$$Q \to \lambda^{\frac{2}{p-1}}Q(\lambda\,\cdot), \quad \lambda > 0. \tag{1.26}$$

The variational values N_1, N_2 and N_3 are closely related each other, as stated in the following:

Proposition 1.1. *Assume that $d \ge 1$ and $2+\frac{4}{d} < p+1 < 2^*$. Then, we have the following relations:*

$$N_1^{\frac{p-1}{2}} = \left(\frac{2}{d}\right)^{\frac{p-1}{2}}\left\{\frac{d(p-1)}{(d+2)-(d-2)p}\right\}^{\frac{1}{4}\{(d+2)-(d-2)p\}} N_2 \tag{1.27}$$

and

$$N_3 = \frac{d(p-1)}{2(p+1)}N_2. \tag{1.28}$$

The next proposition tells us that the variational problems (1.19), (1.20) and (1.21) have the same minimizer.

Proposition 1.2. *Assume that* $d \geq 1$ *and* $2 + \frac{4}{d} < p + 1 < 2^*$. *Then, there exists a positive function* Q *in the Schwartz space* $\mathcal{S}(\mathbb{R}^d)$, *which is unique modulo translation and phase shift, such that*

$$N_1 = \|Q\|_{\widetilde{H}^1}^2, \tag{1.29}$$

$$N_2 = \mathcal{N}_2(Q), \tag{1.30}$$

$$N_3 = \mathcal{I}(Q), \tag{1.31}$$

$$\mathcal{K}(Q) = 0, \tag{1.32}$$

$$\|Q\|_{L^2}^2 = \frac{d+2-(d-2)p}{d(p-1)} \|\nabla Q\|_{L^2}^2 = \frac{d+2-(d-2)p}{2(p+1)} \|Q\|_{L^{p+1}}^{p+1}, \tag{1.33}$$

and Q *solves the equation (1.16) with* $\omega = 1$.

Remark 1.1. *It is known that the function* Q *found in Proposition 1.2 has the following properties:*
(i) *There exists* $y \in \mathbb{R}^d$ *such that* $Q(x-y)$ *is radially symmetric (see [19]).*
(ii) Q *is the ground state of the equation (1.16) with* $\omega = 1$. *Moreover, for all* $\omega > 0$, *the rescaled function* $\omega^{-\frac{1}{p-1}} Q(\omega^{-\frac{1}{2}} \cdot)$ *becomes the ground state of (1.16), and denoting this function by* Q_ω, *we easily verify that*

$$N_2 = \mathcal{N}_2(Q_\omega), \quad N_3 = \mathcal{I}(Q_\omega), \quad \mathcal{K}(Q_\omega) = 0 \qquad \textit{for all } \omega > 0. \tag{1.34}$$

Now, using the variational value N_2, we define a "potential well" PW by

$$PW = \left\{ f \in H^1 \setminus \{0\} \;\middle|\; \mathcal{H}(f) < \mathcal{B}(f) \right\}, \tag{1.35}$$

where

$$\mathcal{B}(f) = \frac{d(p-1)-4}{d(p-1)} \left(\frac{N_2}{\|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}} \right)^{\frac{4}{d(p-1)-4}}. \tag{1.36}$$

We divide PW into two components according to the sign of $\mathcal{K}$:

$$PW_+ = \left\{ f \in PW \;\middle|\; \mathcal{K}(f) > 0 \right\}, \tag{1.37}$$

$$PW_- = \left\{ f \in PW \;\middle|\; \mathcal{K}(f) < 0 \right\}. \tag{1.38}$$

It is worth while noting the following facts:

1. PW_+ and PW_- are unbounded open sets in $H^1(\mathbb{R}^d)$: Indeed, one can easily verify this fact by considering the scaled functions $f_\lambda(x) := \lambda^{\frac{2}{p-1}} f(\lambda x)$ for $f \in H^1(\mathbb{R}^d)$ and $\lambda > 0$.

2. $PW = PW_+ \cup PW_-$ and $PW_+ \cap PW_- = \emptyset$ (see Lemma 2.1)

3. PW_+ and PW_- are invariant under the flow defined by the equation (1.1) (see Proposition 2.5 and Proposition 2.6).

4. The ground state Q_ω belongs to $\overline{PW_+} \cap \overline{PW_-}$ and $Q_\omega \notin PW_+ \cup PW_-$ for all $\omega > 0$, where $\overline{PW_+}$ and $\overline{PW_-}$ are the closures of PW_+ and PW_- in the H^1-topology, respectively (see Theorem 1.3). Moreover, the orbit under the action $((0,\infty) \ltimes \mathbb{R}^d) \times S^1$

$$\left\{ \lambda^{\frac{2}{p-1}} e^{i\theta} Q_\omega(\lambda(\cdot - a)) \;\middle|\; \lambda > 0,\ a \in \mathbb{R}^d,\ \theta \in S^1 \right\} \tag{1.39}$$

is contained in $\overline{PW_+} \cap \overline{PW_-}$.

Here, the last fact above is the key to show the instability of the ground state. We will prove these facts in Section 2, among other properties of these sets PW, PW_+ and PW_-.

For the later convenience, we prepare another expression of PW_+: Besides the functional $\mathcal{N}_2$ and the variational value N_2, we define a functional $\widetilde{\mathcal{N}_2}$ and a number $\widetilde{N}_2$ by

$$\widetilde{\mathcal{N}_2}(f) := \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}(f)}^{\frac{d}{2}(p-1)-2}, \quad f \in H^1(\mathbb{R}^d) \text{ with } \mathcal{H}(f) \geq 0, \tag{1.40}$$

and

$$\widetilde{N}_2 := \sqrt{\frac{d(p-1)-4}{d(p-1)}}^{\frac{d}{2}(p-1)-2} N_2. \tag{1.41}$$

Then, it follows from (1.30) and (1.33) in Proposition 1.2 that

$$\widetilde{N}_2 = \widetilde{\mathcal{N}_2}(Q). \tag{1.42}$$

We can easily verify that: if $f \in H^1(\mathbb{R}^d)$ with $\mathcal{H}(f) \geq 0$, then

$$f \in PW \quad \text{if and only if} \quad \widetilde{\mathcal{N}_2}(f) < \widetilde{N}_2. \tag{1.43}$$

Therefore, we have another expression of PW_+:

$$PW_+ = \left\{ f \in H^1(\mathbb{R}^d) \mid \mathcal{K}(f) > 0,\ \ \widetilde{\mathcal{N}_2}(f) < \widetilde{N}_2 \right\}. \tag{1.44}$$

Here, in order to consider the wave operators, we introduce a set Ω which is a subset of PW_+ (see Remark 1.2, (ii) below):

$$\Omega := \left\{ f \in H^1(\mathbb{R}^d) \setminus \{0\} \mid \mathcal{N}_2(f) < \widetilde{N}_2 \right\}. \tag{1.45}$$

Remark 1.2. (i) *When $p = 1 + \frac{4}{d}$, by (1.40) and (1.41), the condition $\widetilde{\mathcal{N}_2}(f) < \widetilde{N}_2 = \widetilde{\mathcal{N}_2}(Q)$ can be reduced to*

$$\|f\|_{L^2} < \|Q\|_{L^2},$$

since

$$\lim_{p \downarrow 1+\frac{4}{d}} \sqrt{\frac{d(p-1)-4}{d(p-1)}}^{\frac{d}{2}(p-1)-2} = 1.$$

Hence, PW_+ formally becomes in this L^2-critical case

$$\begin{aligned} PW_+ &= \left\{ f \in H^1(\mathbb{R}^d) \;\middle|\; \mathcal{H}(f) > 0,\ \|f\|_{L^2} < \|Q\|_{L^2} \right\} \\ &= \left\{ f \in H^1(\mathbb{R}^d) \setminus \{0\} \;\middle|\; \|f\|_{L^2} < \|Q\|_{L^2} \right\}, \end{aligned} \tag{1.46}$$

where we have used the facts that $\mathcal{K} = \mathcal{H}$ *(see (1.12)) and that* $\|f\|_{L^2} < \|Q\|_{L^2}$ *implies that* $\mathcal{H}(f) > 0$ *(see [55, 45]). On the other hand,* PW_- *formally becomes*

$$PW_- = \left\{ f \in H^1(\mathbb{R}^d) \mid \mathcal{H}(f) < 0 \right\}, \tag{1.47}$$

since $\mathcal{K} = \mathcal{H}$ *(1.12) again. It is well known that the solutions of (1.1) with* $p = 1 + \frac{4}{d}$ *with initial data from the set (1.46) exist globally in time (see [55]), and the ones with data from (1.47) blow up or grow up (see [46]). Thus, we may say that our potential wells* PW_+ *and* PW_- *in (1.37) and (1.38) are natural extensions of those in (1.46) and (1.47) to the case of* $2 + \frac{4}{d} < p + 1 < 2^*$.
(ii) *Since* $\widetilde{N}_2 < N_2$, *we find by the definition of* N_2 *(see (1.20)) that* $\Omega \subset PW_+$.

Now, we are in a position to state our main results. When symbols with ± appear in the following theorems and propositions, we always take both upper signs or both lower signs in the double signs.

The first theorem below is concerned with the behavior of the solutions with initial data from PW_+.

Theorem 1.1 (Global existence and scattering)**.** *Assume that* $d \geq 1$, $2 + \frac{4}{d} < p + 1 < 2^*$ *and* $\psi_0 \in PW_+$. *Then, the corresponding solution* ψ *to the equation (1.1) exists globally in time and has the following properties:*

(i) ψ *stays in* PW_+ *for all time, and satisfies that*

$$\inf_{t \in \mathbb{R}} \mathcal{K}(\psi(t)) \geq \left(1 - \frac{\widetilde{N}_2(\psi_0)}{\widetilde{N}_2} \right) \mathcal{H}(\psi_0) > 0. \tag{1.48}$$

(ii) ψ *belongs to* $L^\infty(\mathbb{R}; H^1(\mathbb{R}))$. *In particular,*

$$\sup_{t \in \mathbb{R}} \|\nabla \psi(t)\|_{L^2}^2 \leq \frac{d(p-1)}{d(p-1) - 4} \mathcal{H}(\psi_0). \tag{1.49}$$

Furthermore,
(iii) *There exist unique* $\phi_+ \in \Omega$ *and* $\phi_- \in \Omega$ *such that*

$$\lim_{t \to \pm\infty} \left\| \psi(t) - e^{\frac{i}{2} t \Delta} \phi_\pm \right\|_{H^1} = \lim_{t \to \pm\infty} \left\| e^{-\frac{i}{2} t \Delta} \psi(t) - \phi_\pm \right\|_{H^1} = 0. \tag{1.50}$$

This formula defines the operators $W_\pm^* \colon \psi_0 \mapsto \phi_\pm = \lim_{t \to \pm\infty} e^{-\frac{i}{2} t \Delta} \psi(t)$. *These operators become homeomorphisms from* PW_+ *to* Ω, *so that we can define the scattering operator* $S := W_+^* W_-$ *from* Ω *into itself, where* $W_- := (W_-^*)^{-1}$*:*

$$S = W_+^* W_- : \begin{array}{ccc} \Omega & \to & \Omega \\ \rotatebox{90}{$\in$} & & \rotatebox{90}{$\in$} \\ \phi_- & \mapsto & \phi_+ \end{array} \quad . \tag{1.51}$$

Remark 1.3. (i) *Theorem 1.1 is an extension of the result by Duyckaerts, Holmer and Roudenko [16]. See* **Notes and Comments** *below for the details.*

(ii) ϕ_+ *and* ϕ_- *found in Theorem 1.1 are called the asymptotic states at* $+\infty$ *and* $-\infty$, *respectively.*

(iii) *To prove the surjectivity of $W_{\pm}^{*}$, we need to construct so-called wave operators $W_{\pm}$ (see Section 3.4). In fact, the operator W_{+}^{*} (W_{-}^{*}) is the inverse of the wave operator W_{+} (W_{-}). According to the terminology of spectrum scattering theory for linear Schrödinger equation, we might say that Ω is a set of scattering states.*

In contrast to the case of PW_{+}, the solutions with initial data from PW_{-} become singular:

Theorem 1.2 (Blowup or growup)**.** *Assume that $d \geq 1$, $2 + \frac{4}{d} < p+1 < 2^{*}$ and $\psi_0 \in PW_{-}$. Then, the corresponding solution ψ to the equation (1.1) satisfies the followings:*

(i) *ψ stays PW_{-} as long as it exists and satisfies that*

$$\mathcal{K}(\psi(t)) < -\left(\mathcal{B}(\psi_0) - \mathcal{H}(\psi_0)\right) < 0 \quad \text{for all } t \in I_{\max}. \tag{1.52}$$

(ii) *ψ blows up in a finite time or grows up, that is,*

$$\sup_{t\in[0,T_{\max}^{+})} \|\nabla\psi(t)\|_{L^2} = \sup_{t\in(-T_{\max}^{-},0]} \|\nabla\psi(t)\|_{L^2} = \infty. \tag{1.53}$$

In particular, if $T_{\max}^{\pm} = \infty$, then we have

$$\limsup_{t\to\pm\infty} \int_{|x|>R} |\nabla\psi(x,t)|^2\,dx = \infty \quad \text{for all } R > 0. \tag{1.54}$$

Remark 1.4. (i) *We do not know whether a solution growing up at infinity exists.*

(ii) *We know (see [20]) that if $\psi_0 \in H^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d, |x|^2dx)$, then $T_{\max}^{\pm} < \infty$ and the corresponding solution ψ satisfies that*

$$\lim_{t\to\pm T_{\max}^{\pm}} \|\nabla\psi(t)\|_{L^2} = \infty.$$

For the case $\psi_0 \notin L^2(\mathbb{R}^d, |x|^2dx)$, see Theorem 1.6 below (see also [48]).

Combining Theorems 1.1 and 1.2, we can show the instability of the ground states: Precisely;

Theorem 1.3 (Instability of ground state)**.** *Let Q_ω be the ground state of the equation (1.16) for $\omega > 0$. Then, Q_ω has two unstable directions in the sense that $Q_\omega \in \overline{PW_{+}} \cap \overline{PW_{-}}$. In particular, for any $\varepsilon > 0$, there exist $f_{+} \in PW_{+}$ and $f_{-} \in PW_{-}$ such that*

$$\|Q_\omega - f_{\pm}\|_{H^1} \leq \varepsilon.$$

Remark 1.5. (i) *An example of $f_{\pm}$ is $\left(1 \mp \frac{\varepsilon}{\|Q_\omega\|_{H^1}}\right) Q_\omega$, where both upper or both lower signs should be chosen in the double signs.*
(ii) *The ground state Q_ω also has an (orbitally) stable "direction". Indeed, if we start with $e^{ivx}Q_\omega$ for "small" $v \in \mathbb{R}^d$, then $e^{i(vx-\frac{1}{2}v^2t)}e^{\frac{i}{2}\omega t}Q_\omega(x-vt)$ solves the equation (1.1) and stays in a neighborhood of the orbit of Q_ω under the action of $\mathbb{R}^d \times S^1$*

$$\left\{ e^{i\theta}Q_\omega(\cdot - a) \;\middle|\; a \in \mathbb{R}^d,\ \theta \in S^1 \right\}.$$

We state further properties of solutions.

Theorem 1.4. *Assume that* $d \geq 1$ *and* $2 + \frac{4}{d} < p + 1 < 2^*$. *Let* ψ *be a global solution to the equation (1.1) with an initial datum in* $H^1(\mathbb{R}^d)$. *Then, the following five conditions are equivalent:*

(i) *Decay of* L^{p+1}*-norm:*

$$\lim_{t \to \infty} \|\psi(t)\|_{L^{p+1}} = 0. \tag{1.55}$$

(ii) *Decay of* L^q*-norm:*

$$\lim_{t \to \infty} \|\psi(t)\|_{L^q} = 0 \quad \text{for all } q \in (2, 2^*). \tag{1.56}$$

(iii) *Boundedness of the Strichartz norms:*

$$\left\|(1 - \Delta)^{\frac{1}{2}} \psi\right\|_{L^r([0,\infty);L^q)} < \infty \quad \text{for all admissible pair } (q, r). \tag{1.57}$$

(iv) *Boundedness of* X*-norm (see (3.9) for the definition of the space* X*):*

$$\|\psi\|_{X([0,\infty))} < \infty, \quad \|\psi\|_{L^\infty([0,\infty);H^1)} < \infty. \tag{1.58}$$

(v) *Existence of an asymptotic state at* $+\infty$*: There exists* $\phi_+ \in H^1(\mathbb{R}^d)$ *such that*

$$\lim_{t \to \infty} \left\|\psi(t) - e^{\frac{i}{2} t\Delta} \phi_+\right\|_{H^1} = 0.$$

A similar result holds for the negative time case.

As a corollary of Theorem 1.1, with the help of Theorem 1.4, we obtain the following result:

Corollary 1.5. $PW_+ \cup \{0\}$ *is connected in the* $L^q(\mathbb{R}^d)$*-topology for all* $q \in (2, 2^*)$.

If, for example, $PW_+ \cup \{0\}$ contains a neighborhood of 0 in the L^q-topology, then we can say that PW_+ is connected in the L^q-topology, without adding $\{0\}$. However, for any $\varepsilon > 0$, there is a function $f_\varepsilon \in H^1(\mathbb{R}^d)$ such that $f_\varepsilon \notin PW_+$ and $\|f_\varepsilon\|_{L^{p+1}} = \varepsilon$, so that we need to add $\{0\}$ in Corollary 1.5 above. On the other hand, we can verify that $PW_+ \cup \{0\}$ contains a sufficiently small ball in $H^1(\mathbb{R}^d)$ (clearly, PW_+ does not contain 0). However, we do not know that PW_+ is connected in the H^1-topology.

Next, we consider singular solutions. The following theorem tells us that solutions with radially symmetric data from PW_- blow up in a finite time:

Theorem 1.6 (Existence of blowup solution)**.** *Assume that* $d \geq 2$, $2 + \frac{4}{d} < p + 1 < 2^*$, *and* $p \leq 5$ *if* $d = 2$. *Let* ψ_0 *be a radially symmetric function in* PW_- *and let* ψ *be the corresponding solution to the equation (1.1). Then, we have*

$$T_{\max}^{\pm} < \infty \quad \text{and} \quad \lim_{t \to \pm T_{\max}^{\pm}} \|\nabla \psi(t)\|_{L^2} = \infty. \tag{1.59}$$

Furthermore, we have the followings:
(i) *For all* $m > 0$, *there exists a constant* $R_m > 0$ *such that*

$$\int_{|x|>R} |\psi(x,t)|^2 \, dx < m \quad \text{for all } R \geq R_m \text{ and } t \in I_{\max}. \tag{1.60}$$

(ii) *For all sufficiently large $R>0$, we have*

$$\int_0^{T_{\max}^+}(T_{\max}^+-t)\left(\int_{|x|>R}|\nabla\psi(x,t)|^2dx\right)dt<\infty, \tag{1.61}$$

$$\int_0^{T_{\max}^+}(T_{\max}^+-t)\left(\int_{|x|>R}|\psi(x,t)|^{p+1}dx\right)dt<\infty. \tag{1.62}$$

(iii) *For all sufficiently large $R>0$, we have*

$$\int_0^{T_{\max}^+}(T_{\max}^+-t)\|\psi(t)\|_{L^\infty(|x|>R)}^4\,dt<\infty, \tag{1.63}$$

$$\int_0^{T_{\max}^+}(T_{\max}^+-t)\|\psi(t)\|_{L^{p+1}(|x|>R)}^{\frac{4(p+1)}{p-1}}\,dt<\infty, \tag{1.64}$$

$$\liminf_{t\to T_{\max}^+}(T_{\max}^+-t)\|\psi(t)\|_{L^\infty(|x|>R)}^2=0, \tag{1.65}$$

$$\liminf_{t\to T_{\max}^+}(T_{\max}^+-t)^{\frac{p-1}{2}}\|\psi(t)\|_{L^{p+1}(|x|>R)}^{p+1}=0. \tag{1.66}$$

For the negative time case, the corresponding results to (ii) *and* (iii) *hold valid.*

In Theorem 1.6, we require the condition $p\le 5$ if $d=2$, as well as Ogawa and Tsutsumi [48]. A difficulty in the case $p>5$ with $d=2$ might come from a kind of quantum forces such that the stronger the nonlinear effect becomes, the stronger the dispersion effect does.

We do not know a lot of things about the asymptotic behavior of such singular solutions as found in Theorem 1.2. What we can say is the following (for simplicity, we state the forward time case only):

Proposition 1.3 (Asymptotic profiles of singular solutions)**.** *Assume that $d\ge 1$ and $2+\frac{4}{d}<p+1<2^*$. Let ψ be a solution to the equation (1.1) such that*

$$\limsup_{t\to T_{\max}^+}\|\nabla\psi(t)\|_{L^2}=\limsup_{t\to T_{\max}^+}\|\psi(t)\|_{L^{p+1}}=\infty, \tag{1.67}$$

and let $\{t_n\}_{n\in\mathbb{N}}$ be a sequence in $[0,T_{\max}^+)$ such that

$$\lim_{n\to\infty}t_n=T_{\max}^+,\qquad \|\psi(t_n)\|_{L^{p+1}}=\sup_{t\in[0,t_n)}\|\psi(t)\|_{L^{p+1}}. \tag{1.68}$$

For this sequence $\{t_n\}$, we put

$$\lambda_n=\|\psi(t_n)\|_{L^{p+1}}^{-\frac{(p-1)(p+1)}{d+2-(d-2)p}}, \tag{1.69}$$

and consider the scaled functions

$$\psi_n(x,t) := \lambda_n^{\frac{2}{p-1}} \overline{\psi(\lambda_n x, t_n - \lambda_n^2 t)}, \quad t \in \Big(-\frac{T_{\max}^{+} - t_n}{\lambda_n^2}, \frac{t_n}{\lambda_n^2} \Big], \quad n \in \mathbb{N}. \tag{1.70}$$

Suppose that

$$\|\psi\|_{L^\infty([0,T_{\max}^{+});L^{\frac{d}{2}(p-1)})} < \infty. \tag{1.71}$$

Then, there exists a subsequence of $\{\psi_n\}$ *(still denoted by the same symbol) with the following properties: There exist*
(i) *a nontrivial function* $\psi_\infty \in L^\infty([0,\infty); \dot{H}^1(\mathbb{R}^d) \cap L^{p+1}(\mathbb{R}^d))$ *solving the equation (1.1) in the* $\mathcal{D}'([0,\infty); \dot{H}^{-1}(\mathbb{R}^d) + L^{\frac{p+1}{p}}(\mathbb{R}^d))$*-sense and*
(ii) *a sequence* $\{\gamma_n\}$ *in* $\mathbb{R}^d$
such that, putting $\widetilde{\psi}_n(x,t) := \psi_n(x+\gamma_n, t)$*, we have, for any* $T > 0$*,*

$$\lim_{n\to\infty} \widetilde{\psi}_n = \psi_\infty \quad \textit{strongly in } L^\infty([0,T]; L^q_{loc}(\mathbb{R}^d)) \quad \textit{for all } q \in [1, 2^*),$$

$$\lim_{n\to\infty} \nabla\widetilde{\psi}_n = \nabla\psi_\infty \quad \textit{weakly* in } L^\infty([0,T]; L^2_{loc}(\mathbb{R}^d)).$$

Furthermore, for any $\varepsilon > 0$*, there exists* $R > 0$ *such that, for* $y_n = \lambda_n \gamma_n$*,*

$$\lim_{n\to\infty} \int_{|x-y_n|\le\lambda_n R} |\psi(x,t_n)|^{\frac{d}{2}(p-1)} dx \ge (1-\varepsilon) \|\psi_\infty(0)\|_{L^{\frac{d}{2}(p-1)}}^{\frac{d}{2}(p-1)}. \tag{1.72}$$

Remark 1.6. *The* $L^{\frac{d(p-1)}{2}}(\mathbb{R}^d)$*-norm is invariant under the scaling leaving the equation (1.1) invariant: Precisely, when* ψ *is a solution to (1.1), putting*

$$\psi_\lambda(x,t) := \lambda^{\frac{d(p-1)}{2}} \psi(\lambda x, \lambda^2 t), \quad \lambda > 0, \tag{1.73}$$

we see that ψ_λ *solves (1.1) and satisfies that*

$$\|\psi_\lambda(t)\|_{L^{\frac{d}{2}(p-1)}} = \|\psi(\lambda t)\|_{L^{\frac{d}{2}(p-1)}} \quad \textit{for all } t \in (-\tfrac{T_{\max}^{-}}{\lambda}, \tfrac{T_{\max}^{+}}{\lambda}). \tag{1.74}$$

Proposition 1.3 tells us that the $L^{\frac{d(p-1)}{2}}(\mathbb{R}^d)$-norm of a singular solution concentrates at some point under the assumption (1.71). In general, it is difficult to check whether the assumption (1.71) holds (cf. Merle and Raphaël [39]).

Without the assumption (1.71), we have the following:

Proposition 1.4. *Under the same assumptions except (1.71), definitions and notation in Proposition 1.3, we define the "renormalized" functions* Φ_n^{RN} *by*

$$\Phi_n^{RN}(x,t) = \psi_n(x,t) - e^{\frac{i}{2}t\Delta}\psi_n(x,0), \quad n \in \mathbb{N}. \tag{1.75}$$

Then, for any $T > 0$*,* $\{\Phi_n^{RN}\}_{n\in\mathbb{N}}$ *is a uniformly bounded sequence in* $C([0,T]; H^1(\mathbb{R}^d))$*, and satisfies the following alternatives* (i) *and* (ii)*:*
(i) *If*

$$\lim_{n\to\infty} \sup_{t\in[0,T]} \|\Phi_n^{RN}(t)\|_{L^{\frac{d}{2}(p-1)}} = 0, \tag{1.76}$$

then

$$\lim_{n\to\infty} \sup_{t\in[t_n-\lambda_n^2 T,, t_n]} \Big\| \psi(t) - e^{\frac{i}{2}(t-t_n)\Delta}\psi(t_n) \Big\|_{L^{\frac{d}{2}(p-1)}} = 0. \tag{1.77}$$

(ii) *If*

$$\limsup_{n\to\infty} \sup_{t\in[0,T]} \left\|\Phi_n^{RN}(t)\right\|_{L^{\frac{d}{2}(p-1)}} > 0, \tag{1.78}$$

then there exists a subsequence of $\{\Phi_n^{RN}\}$ *(still denoted by the same symbol) with the following properties: There exist a nontrivial function* $\Phi \in L^\infty([0,\infty); H^1(\mathbb{R}^d))$ *and a sequence* $\{y_n\}$ *in* $\mathbb{R}^d$ *such that, putting* $\widetilde{\Phi}_n^{RN}(x,t) = \Phi_n^{RN}(x+y_n,t)$, *we have*

$$\lim_{n\to\infty} \widetilde{\Phi}_n^{RN} = \Phi \quad \textit{weakly* in } L^\infty([0,T]; H^1(\mathbb{R}^d)). \tag{1.79}$$

Here, Φ *solves the following equation*

$$2i\frac{\partial\Phi}{\partial t} + \Delta\Phi = -F, \tag{1.80}$$

where F *is the nontrivial function in* $L^\infty([0,\infty); L^{\frac{p+1}{p}}(\mathbb{R}^d))$ *given by*

$$\lim_{n\to\infty} |\psi_n|^{p-1}\psi_n = F \quad \textit{weakly* in} \;\; L^\infty([0,T]; L^{\frac{p+1}{p}}(\mathbb{R}^d)). \tag{1.81}$$

Furthermore, for any $\varepsilon > 0$, *there exists* $R > 0$ *such that*

$$\lim_{n\to\infty} \int_{|x-y_n|\le\lambda_n R} \left|\psi(x, t_n - \lambda_n^2 T) - e^{-\frac{i}{2}\lambda_n^2 T\Delta}\psi(x,t_n)\right|^{\frac{d}{2}(p-1)} dx \ge (1-\varepsilon)\left\|\Phi(T)\right\|_{L^{\frac{d}{2}(p-1)}}^{\frac{d}{2}(p-1)}. \tag{1.82}$$

If the case (i) of Proposition 1.4 occurs, then we may say that the dynamics of the solution is composed of the free evolution and the dilation (1.73). On the other hand, in (ii), concentration of $L^{\frac{d(p-1)}{2}}$-mass occurs further. We remark that the left-hand side of (1.82) is finite.

Here, we discuss some relations between the previous works and our results:
Notes and Comments.

1. Our analysis in PW_+ is inspired by the previous work by Duyckaerts, Holmer and Roudenko [16, 23] (also Kenig and Merle [30]). They considered a typical nonlinear Schrödinger equation, the equation (1.1) with $d = p = 3$, and proved, in [16], that: if $\psi_0 \in H^1(\mathbb{R}^3)$ satisfies that
$$\mathcal{M}(\psi_0)\mathcal{H}(\psi_0) < \mathcal{M}(Q)\mathcal{H}(Q), \quad \|\psi_0\|_{L^2}\|\nabla\psi_0\|_{L^2} < \|Q\|_{L^2}\|\nabla Q\|_{L^2}, \tag{1.83}$$
then the corresponding solution exists globally in time and has asymptotic states at ±∞, where Q denotes the ground state of the equation (1.16) with $\omega = 1$. In our terminology, we see that the condition (1.83) is equivalent to that $\psi_0 \in PW_+ \cup \{0\}$ via the variational problem for N_2 (see (1.20)). In this paper, we intensively study the scattering problem on PW_+, so that we have Theorem 1.1, which is an extension of the result by Duycaerts et al [16, 23] to all spatial dimensions $d \ge 1$ and L^2-supercritical and H^1-subcritical powers $2 + \frac{4}{d} < p+1 < 2^*$. Furthermore, we establish the so-called asymptotic completeness: the wave operators $W_\pm$ exists on Ω and they are homeomorphisms from Ω to PW_+.
For the nonlinear Klein-Gordon equation, the corresponding result to Theorem 1.1 was obtained by Ibrahim, Masmoudi and Nakanishi [25].

2. In order to prove PW_+ being a set of scattering states, we basically employ the argument of Kenig and Merle [30]. In their argument, the Bahouri-Gerard type compactness [1] plays an important role: Duyckaerts et al [16] also used such a compactness (the profile decomposition due to Keraani [31]). However, we employ the classical compactness device due to Brezis and Lieb [8] instead of that due to Bahouri and Gerard; The Brezis-Lieb type compactness device is also used to prove the existence of the ground states (see Section F), and to investigate the blowup solutions (see the proof of Proposition 1.4 in Section 5). As long as we consider the H^1-solutions, this classical compactness devise seems to be enough for our analysis.

3. The decomposition scheme in Sections 4.2 (also the profile decomposition due to Bahouri-Gerard [1], Keraani [31]) seems to be a kind of perturbation methods employed in quantum physics like a "Born type approximation scheme".

4. In the course of the proof of Theorem 1.1, we encounter a "fake soliton" (critical element in the terminology of Kenig-Merle [30]). Then, we make a slightly different approach from Duyckaerts et al [16] to trace its motion (for details, see Section 4.1). Moreover, our choice of function spaces is different from theirs [16, 23] (see Section 3.1). We are choosing our function spaces so that the generalized inhomogeneous Strichartz estimates due to Foshci [17] work well there.

5. In [23] (also in [24]), Holmer and Roudenko also considered the equation (1.1) with $d = p = 3$ and proved that if $\psi_0 \in H^1(\mathbb{R}^3)$ satisfies that

$$\psi_0 \text{ is radially symmetric,} \tag{1.84}$$

$$\mathcal{M}(\psi_0)\mathcal{H}(\psi_0) < \mathcal{M}(Q)\mathcal{H}(Q), \quad \|\psi_0\|_{L^2} \|\nabla\psi_0\|_{L^2} > \|Q\|_{L^2} \|\nabla Q\|_{L^2}, \tag{1.85}$$

   then the corresponding solution blows up in a finite time. In our terminology, we see that the condition (1.85) is equivalent to $\psi_0 \in PW_-$ via the variational problem for N_2. Hence, Theorem 1.2, together with Theorem 1.6, is an extension of their result, in particular, to all spatial dimensions $d \geq 1$ and powers $2 + \frac{4}{d} < p + 1 < 2^*$.

6. Our PW_+ and PW_- are naturally introduced by the potential well PW by appealing to the variational structure of the ground states. We note again that the functional $\mathcal{K}$ divides the PW into PW_+ and PW_-. Our potential well PW seems new. One may find a similarity between PW and the set of initial data given in Theorem 4.1 in Begout [2]. However, the relevance is not clear.

7. Stubbe [52] already introduced the condition (1.83) and proved the global existence of the solutions with initial data satisfying it. He also conjectured that the condition is sharp in the sense that there exists an initial datum such that it does not satisfy the condition (1.83) and leads to a solution blowing up in a finite time. Our result concerning PW_- gives an affirmative answer to his conjecture.

This paper is organized as follows. In Section 2, we discuss properties of the potential well PW. In Section 3, we introduce function spaces in which Strichartz type estimates work well. We also give a small date theory and a long time perturbation theory. Theorem 1.4 is proved here (see Section 3.2). In Section 4, we give the proofs of Theorem 1.1 and Corollary 1.5. Section 5 is devoted to the proofs of Theorem 1.2, Theorem 1.6, and Propositions 1.3 and 1.4. In Appendices A, B, C, D and E are devoted to preliminaries and auxiliary results. Finally, in Section F, we give the proofs of Propositions 1.1 and 1.2.

Notation. We summarize the notation used in this paper.

We keep the letters d and p to denote the spatial dimension and the power of nonlinearity of the equation (1.1), respectively.

$\mathbb{N}$ denotes the set of natural numbers, i.e., $\mathbb{N} = \{1, 2, 3, \ldots\}$.

$I_{\max}$ denotes the maximal existence interval of the considering solution, which has the form

$$I_{\max} = (-T_{\max}^{-}, T_{\max}^{+}),$$

where $T_{\max}^{+} > 0$ is the maximal existence time for the future, and $T_{\max}^{-} > 0$ is the one for the past.

Functionals concerned with conservation laws for the equation (1.1) are: the mass

$$\mathcal{M}(f) := \|f\|_{L^2}^2 \qquad \text{(see (1.5))},$$

the Hamiltonian

$$\mathcal{H}(f) := \|\nabla f\|_{L^2}^2 - \frac{2}{p+1}\|f\|_{L^{p+1}}^{p+1} \qquad \text{(see (1.6))},$$

and the momentum

$$\mathcal{P}(f) =: \Im \int_{\mathbb{R}^d} \nabla f(x)\overline{f(x)}\, dx \qquad \text{(see (1.7))}.$$

We also use the functional

$$\mathcal{K}(f) := \|\nabla f\|_{L^2}^2 - \frac{d(p-1)}{2(p+1)}\|f\|_{L^{p+1}}^{p+1} \qquad \text{(see (1.9))}.$$

The symbol $\mathcal{K}$ might stand for “Kamiltonian” (?).

Functionals concerned with variational problems are the followings:

$$\|f\|_{\widetilde{H}^1}^2 := \frac{d(p-1)-4}{d(p-1)}\|\nabla f\|_{L^2}^2 + \|f\|_{L^2}^2 \qquad \text{(see (1.22))},$$

$$\mathcal{N}_2(f) := \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla f\|_{L^2}^{\frac{d}{2}(p-1)-2} \qquad \text{(see (1.23))},$$

$$\mathcal{I}(f) := \frac{\|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla f\|_{L^2}^{\frac{d}{2}(p-1)}}{\|f\|_{L^{p+1}}^{p+1}} \qquad \text{(see (1.24)))}.$$

Variational values concerned with these functionals are

$$N_1 := \inf\Big\{\|f\|_{\widetilde{H}^1}^2 \;\Big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\},\ \mathcal{K}(f) \le 0\Big\} \qquad \text{(see (1.19))},$$

$$N_2 := \inf\Big\{\mathcal{N}_2(f) \;\Big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\},\ \mathcal{K}(f) \le 0\Big\} \qquad \text{(see (1.20))},$$

$$N_3 := \inf\Big\{\mathcal{I}(f) \;\Big|\; f \in H^1(\mathbb{R}^d)\setminus\{0\}\Big\} \qquad \text{(see (1.21))}.$$

We define our “potential well” by

$$PW := \big\{f \in H^1\setminus\{0\} \;\big|\; \mathcal{H}(f) < \mathcal{B}(f)\big\} \qquad \text{(see (1.35))},$$

where

$$\mathcal{B}(f) := \frac{d(p-1)-4}{d(p-1)} \left(\frac{N_2}{\|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)}} \right)^{\frac{4}{d(p-1)-4}} \qquad \text{(see (1.36))}.$$

We put

$$\varepsilon_0 := \mathcal{B}(\psi_0) - \mathcal{H}(\psi_0).$$

We divide the set PW into two disjoint components:

$$PW_+ := \{ f \in PW \mid \mathcal{K}(f) > 0 \} \qquad \text{(see (1.37))},$$

$$PW_- := \{ f \in PW \mid \mathcal{K}(f) < 0 \} \qquad \text{(see (1.38))}.$$

We can rewrite the set PW_+ in the form

$$PW_+ = \left\{ f \in H^1(\mathbb{R}^d) \;\middle|\; \mathcal{K}(f) > 0,\ \widetilde{\mathcal{N}}_2(f) < \widetilde{N}_2 \right\},$$

where

$$\widetilde{\mathcal{N}}_2(f) := \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}(f)}^{\frac{d}{2}(p-1)-2} \qquad \text{(see (1.40))}$$

and

$$\widetilde{N}_2 := \sqrt{\frac{d(p-1)-4}{d(p-1)}}^{\frac{d}{2}(p-1)-2} N_2 \qquad \text{(see (1.41))}.$$

We need a subset Ω of PW_+ below to consider the wave operators:

$$\Omega := \left\{ f \in H^1(\mathbb{R}^d) \setminus \{0\} \;\middle|\; \mathcal{N}_2(f) < \widetilde{N}_2 \right\} \qquad \text{(see (1.45))}.$$

s_p stands for the critical regularity of our equation (1.1), i.e.,

$$s_p := \frac{d}{2} - \frac{2}{p-1} \qquad \text{(see (3.2))}.$$

We will fix a number $q_1 \in (p+1, 2^*)$ in Sections 3 and 4. The number r_0 is chosen for the pair (q_1, r_0) being admissible, i.e.,

$$\frac{1}{r_0} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1} \right).$$

Furthermore, r_1 and $\widetilde{r}_1$ are defined respectively by

$$\frac{1}{r_1} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1} - \frac{s_p}{d} \right),$$

$$\frac{1}{\widetilde{r}_1} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1} + \frac{s_p}{d} \right).$$

A pair (q_2, r_2) is defined by

$$\frac{1}{q_2} := \frac{1}{p-1}\left(1 - \frac{2}{q_1} \right), \quad \frac{1}{r_2} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_2} - \frac{s_p}{d} \right).$$

For an interval I, we define the Strichartz type space $X(I)$ by

$$X(I) := L^{r_1}(I; L^{q_1}(\mathbb{R}^d)) \cap L^{r_2}(I; L^{q_2}(\mathbb{R}^d)) \qquad \text{(see (3.9))}.$$

Besides, we define the usual Strichartz space $S(I)$ by

$$S(I) := L^{\infty}(I; L^2(\mathbb{R}^d)) \cap L^{r_0}(I; L^{q_1}(\mathbb{R}^d)) \qquad \text{(see (3.10))}.$$

In order to show that the solutions starting from PW_+ have asymptotic states, we will introduce a set

$$PW_+(\delta) := \left\{ f \in PW_+ \,\middle|\, \widetilde{\mathcal{N}}_2(f) < \delta \right\}, \quad \delta > 0 \qquad \text{(see (4.1))}.$$

We will also consider a variational value

$$\begin{aligned} \widetilde{N}_c &:= \sup \left\{ \delta > 0 \,\middle|\, \forall \psi_0 \in PW_+(\delta),\ \|\psi\|_{X(\mathbb{R})} < \infty \right\} \\ &= \inf \left\{ \delta > 0 \,\middle|\, \exists \psi_0 \in PW_+(\delta),\ \|\psi\|_{X(\mathbb{R})} = \infty \right\} \qquad \text{(see (4.2))}, \end{aligned}$$

where ψ denotes the solution to (1.1) with $\psi(0) = \psi_0$.

If $(A_0, \|\cdot\|_{A_0})$ and $(A_1, \|\cdot\|_{A_1})$ are "compatible" normed vector spaces (see [6], p.24.), then $\|\cdot\|_{A_0 \cap A_1}$ denotes the norm of their intersection $X \cap Y$, i.e.,

$$\|f\|_{A_0 \cap A_1} := \max\{\|f\|_{A_0}, \|f\|_{A_1}\} \quad \text{for all } f \in A_0 \cap A_1.$$

The symbol $(\cdot, \cdot)$ denotes the inner product of $L^2(\mathbb{R}^d) = L^2(\mathbb{R}^d; \mathbb{C})$, i.e.,

$$(f, g) := \int_{\mathbb{R}^d} f(x)\overline{g(x)}\, dx, \quad f, g \in L^2(\mathbb{R}^d).$$

$C_c^{\infty}(\mathbb{R}^d)$ denotes the set of infinitely differentiable functions from $\mathbb{R}^d \to \mathbb{C}$ with compact supports.

Using the Fourier transformation $\mathcal{F}$, we define differential operators $|\nabla|^s$, $(-\Delta)^{\frac{s}{2}}$ and $(1 - \Delta)^{\frac{s}{2}}$, for $s \in \mathbb{R}$, by

$$|\nabla|^s f = (-\Delta)^{\frac{s}{2}} f := \mathcal{F}^{-1}[|\xi|^s \mathcal{F}[f]], \qquad (1 - \Delta)^{\frac{s}{2}} f := \mathcal{F}^{-1}[(1 + |\xi|^2)^{\frac{s}{2}} \mathcal{F}[f]].$$

2 Potential well *PW*

In this section, we shall discuss fundamental properties of the sets PW, PW_- and PW_+. In particular, we will prove that these sets are invariant under the flow defined by the equation (1.1) (see Propositions 2.4, 2.5 and 2.6). Moreover, we prove Theorem 1.3 here.

We begin with the following fact:

Lemma 2.1. *The set PW does not contain any function f with $\mathcal{K}(f) = 0$, i.e.,*

$$\left\{ f \in H^1(\mathbb{R}^d) \,\middle|\, \mathcal{K}(f) = 0 \right\} \cap PW = \emptyset, \tag{2.1}$$

so that

$$PW = PW_+ \cup PW_-, \qquad PW_+ \cap PW_- = \emptyset. \tag{2.2}$$

Proof of Lemma 2.1. Let f be a function in $H^1(\mathbb{R}^d)$ with $\mathcal{K}(f)=0$. Then, the condition $\mathcal{K}(f)=0$ leads to that

$$\mathcal{H}(f)=\frac{d(p-1)-4}{d(p-1)}\|\nabla f\|_{L^2}^2. \tag{2.3}$$

Using (2.3) and the definitions of N_2 (see (1.20)) and $\widetilde{N}_2$ (see (1.41)), we obtain that

$$\begin{aligned}\widetilde{\mathcal{N}}_2(f)&=\sqrt{\frac{d(p-1)-4}{d(p-1)}}^{\frac{d}{2}(p-1)-2}\mathcal{N}_2(f)\\ &\ge\sqrt{\frac{d(p-1)-4}{d(p-1)}}^{\frac{d}{2}(p-1)-2}N_2=\widetilde{N}_2.\end{aligned} \tag{2.4}$$

Hence, it follows from (1.43) that $f\notin PW$.

We immediately obtain (2.2) from (2.1) and the definitions of PW_+ and PW_- (see (1.37) and (1.38)). ☐

In the next lemma, we consider a path constructed from the ground state:

Lemma 2.2. *Let Q_ω be the ground state of the equation (1.16) for $w>0$. We consider a path $\Gamma_\omega\colon[0,\infty)\to H^1(\mathbb{R}^d)$ given by $\Gamma_\omega(s)=sQ_\omega$ for $s\ge 0$. Then, Γ_ω is continuous and satisfies that*

$$\Gamma_\omega(s)\in PW_+\quad\textit{for all } s\in(0,1), \tag{2.5}$$

$$\Gamma_\omega(1)=Q_\omega\notin PW=PW_+\cup PW_-, \tag{2.6}$$

$$\Gamma_\omega(s)\in\Big\{f\in H^1(\mathbb{R}^d)\ \Big|\ \mathcal{H}(f)>0\Big\}\cap PW_-\quad\textit{for all } s\in\left(1,\ \left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}}\right), \tag{2.7}$$

$$\Gamma_\omega(s)\in\Big\{f\in H^1(\mathbb{R}^d)\setminus\{0\}\ \Big|\ \mathcal{H}(f)\le 0\Big\}\subset PW_-\quad\textit{for all } s\in\left[\left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}},\ \infty\right). \tag{2.8}$$

In particular, $PW_-\neq\emptyset$ and $PW_+\neq\emptyset$.

Proof of Lemma 2.2. The continuity of $\Gamma_\omega\colon(0,\infty)\to H^1(\mathbb{R}^d)$ is obvious from its definition. We shall prove the properties (2.5)–(2.8). As stated in (1.34) (see also (1.18)), the ground state Q_ω satisfies that $\mathcal{K}(Q_\omega)=0$, which immediately yields that

$$\|\nabla Q_\omega\|_{L^2}^2=\frac{d(p-1)}{2(p+1)}\|Q_\omega\|_{L^{p+1}}^{p+1}. \tag{2.9}$$

Using (2.9), we obtain that

$$\mathcal{H}(\Gamma_\omega(s))=s^2\left(1-\frac{4}{d(p-1)}s^{p-1}\right)\|\nabla Q_\omega\|_{L^2}^2. \tag{2.10}$$

This formula gives us that

$$\mathcal{H}(\Gamma_\omega(s))>0\quad\text{for all } s\in\left(0,\ \left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}}\right), \tag{2.11}$$

$$\mathcal{H}(\Gamma_\omega(s))\le 0\quad\text{for all } s\in\left[\left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}},\ \infty\right). \tag{2.12}$$

Similarly, we can verify that

$$\mathcal{K}(\Gamma_\omega(s)) > 0 \quad \text{for all } s \in (0,1), \tag{2.13}$$

$$\mathcal{K}(\Gamma_\omega(1)) = \mathcal{K}(Q_\omega) = 0, \tag{2.14}$$

$$\mathcal{K}(\Gamma_\omega(s)) < 0 \quad \text{for all } s \in (1,\infty). \tag{2.15}$$

Now, it follows from (2.10) and (1.34) that

$$\widetilde{\mathcal{N}}_2(\Gamma_\omega(s)) = s^{p-1}\left(1 - \frac{4}{d(p-1)}s^{p-1}\right)^{\frac{d}{4}(p-1)-1} N_2 \qquad \text{for all } s \in \left(0,\ \left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}}\right). \tag{2.16}$$

This relation (2.16) shows that

$$\widetilde{\mathcal{N}}_2(\Gamma_\omega(s)) < \widetilde{\mathcal{N}}_2(\Gamma_\omega(1)) = \widetilde{N}_2 \qquad \text{for all } s \in \left(0,\ \left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}}\right) \setminus \{1\}. \tag{2.17}$$

Hence, (2.11) and (2.17), with the help of (1.43), lead us to the conclusion that

$$\Gamma_\omega(s) \in PW \qquad \text{for all } s \in \left(0,\ \left\{\tfrac{d(p-1)}{4}\right\}^{\frac{1}{p-1}}\right) \setminus \{1\}. \tag{2.18}$$

Then, (2.5) follows from (2.13). Moreover, (2.7) follows from (2.11) and (2.15). The second claim (2.6) is a direct consequence of (2.14) and Lemma 2.1.

It remains to prove (2.8). Since $\mathcal{B}(f) \geq 0$ and $\mathcal{K}(f) < \mathcal{H}(f)$ for all $f \in H^1(\mathbb{R}^d) \setminus \{0\}$, we have a relation

$$\left\{ f \in H^1(\mathbb{R}^d) \setminus \{0\} \;\middle|\; \mathcal{H}(f) \leq 0 \right\} \subset PW_-. \tag{2.19}$$

Then, (2.8) immediately follows from (2.12) and this relation (2.19). ☐

Now, we are in a position to prove Theorem 1.3.

Proof of Theorem 1.3. We consider the path Γ_ω given in Lemma 2.2. Then, (2.5) and the continuity of Γ_ω yield that

$$\Gamma_\omega(s) \in PW_+ \quad \text{for all } s \in (0,1), \qquad \lim_{s\uparrow 1}\Gamma_\omega(s) = \Gamma_\omega(1) = Q_\omega \quad \text{strongly in } H^1(\mathbb{R}^d). \tag{2.20}$$

Hence, we have $Q_\omega \in \overline{PW_+}$. Similarly, (2.7) and the continuity of Γ_ω show that $Q_\omega \in \overline{PW_-}$. ☐

We find from the following lemma that PW_+ has a “foliate structure”.

Lemma 2.3. *For all $\eta \in (0, \widetilde{N}_2)$ and $\alpha > 0$, there exists $f \in PW_+$ such that $\widetilde{\mathcal{N}}_2(f) = \eta$ and $\|f\|_{L^2} = \alpha$.*

Proof of Lemma 2.3. We construct a desired function from the continuous path $\Gamma_\omega \colon [0,\infty) \to H^1(\mathbb{R}^d)$ ($\omega > 0$) given in Lemma 2.2. Let us remind you that

$$\Gamma_\omega(s) \in PW_+ \quad \text{for all } s \in (0,1), \tag{2.21}$$

$$\widetilde{\mathcal{N}}_2(\Gamma_\omega(s)) = s^{p-1}\left(1 - \frac{4}{d(p-1)}s^{p-1}\right)^{\frac{d}{4}(p-1)-1} N_2 \quad \text{for all } s \in (0,1), \tag{2.22}$$

$$\widetilde{\mathcal{N}}_2(\Gamma_\omega(0)) = 0, \qquad \widetilde{\mathcal{N}}_2(\Gamma_\omega(1)) = \widetilde{N}_2 \tag{2.23}$$

(see (2.16) for (2.22)). Using (2.22), we find that $\widetilde{\mathcal{N}}_2(\Gamma_\omega(s))$ is continuous and monotone increasing with respect to s on $[0,1]$. Hence, the intermediate value theorem, together with (2.23), shows that for any $\eta \in (0, \widetilde{N}_2)$, there exists $s_\eta \in (0,1)$ such that $\widetilde{\mathcal{N}}_2(\Gamma_\omega(s_\eta)) = \eta$. Moreover, (2.21) gives us that $\Gamma_\omega(s_\eta) \in PW_+$. We put $f = \Gamma_\omega(s_\eta)$ and consider the scaled function $f_\lambda := \lambda^{\frac{2}{p-1}} f(\lambda \cdot)$ for $\lambda > 0$. Then, it is easy to see that

$$f_\lambda \in PW_+, \quad \widetilde{\mathcal{N}}_2(f_\lambda) = \widetilde{\mathcal{N}}_2(f) = \eta, \quad \|f_\lambda\|_{L^2} = \lambda^{-\frac{d(p-1)-4}{2(p-1)}} \|f\|_{L^2} \quad \text{for all } \lambda > 0. \tag{2.24}$$

Hence, for any $\alpha > 0$, f_λ with $\lambda = \left(\frac{\|f\|_{L^2}}{\alpha}\right)^{\frac{2(p-1)}{d(p-1)-4}}$ is what we want. □

Finally, we give the invariance results of the sets PW, PW_+ and PW_- under the flow defined by the equation (1.1):

Lemma 2.4 (Invariance of PW). *Let $\psi_0 \in PW$ and ψ be the corresponding solution to the equation (1.1). Then, we have that*

$$\psi(t) \in PW \quad \textit{for all } t \in I_{\max}.$$

Proof of Lemma 2.4. This lemma immediately follows from the mass and energy conservation laws (1.5) and (1.6). □

Proposition 2.5 (Invariance of PW_+). *Let $\psi_0 \in PW_+$ and ψ be the corresponding solution to the equation (1.1). Then, ψ exists globally in time and satisfies the followings:*

$$\psi(t) \in PW_+ \quad \textit{for all } t \in \mathbb{R}, \tag{2.25}$$

$$\|\nabla \psi(t)\|_{L^2}^2 < \frac{d(p-1)}{d(p-1)-4} \mathcal{H}(\psi_0) \quad \textit{for all } t \in \mathbb{R}, \tag{2.26}$$

$$\mathcal{K}(\psi(t)) > \left(1 - \frac{\widetilde{N}_2(\psi_0)}{\widetilde{N}_2}\right) \mathcal{H}(\psi_0) \quad \textit{for all } t \in \mathbb{R}. \tag{2.27}$$

Proof of Proposition 2.5. We first prove the invariance of PW_+ under the flow defined by (1.1). With the help of Lemma 2.4, it suffices to show that

$$\mathcal{K}(\psi(t)) > 0 \quad \text{for all } t \in I_{\max}. \tag{2.28}$$

Supposing the contrary that (2.28) fails, we can take $t_0 \in I_{\max}$ such that

$$\mathcal{K}(\psi(t_0)) = 0. \tag{2.29}$$

Then, (2.29) and the energy conservation law (1.6) yield that

$$\begin{aligned} 0 &= \mathcal{K}(\psi(t_0)) \\ &= \mathcal{H}(\psi(t_0)) - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\} \|\psi(t_0)\|_{L^{p+1}}^{p+1} \\ &= \mathcal{H}(\psi_0) - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\} \frac{2(p+1)}{d(p-1)} \|\nabla \psi(t_0)\|_{L^2}^2. \end{aligned} \tag{2.30}$$

Since $\psi_0 \in PW_+ \subset PW$, we have $\mathcal{H}(\psi_0) < \mathcal{B}(\psi_0)$. This inequality and (2.30) lead us to that

$$0 < \mathcal{B}(\psi_0) - \frac{d(p-1)-4}{d(p-1)} \|\nabla\psi(t_0)\|_{L^2}^2, \tag{2.31}$$

which is equivalent to

$$\|\nabla\psi(t_0)\|_{L^2}^2 < \frac{d(p-1)}{d(p-1)-4}\mathcal{B}(\psi_0) = \left(\frac{N_2}{\|\psi(t_0)\|_{L^2}^{p+1-\frac{d}{2}(p-1)}} \right)^{\frac{4}{d(p-1)-4}}. \tag{2.32}$$

Dividing the both sides of (2.32) by $\|\nabla\psi(t_0)\|_{L^2}^2$, we obtain that

$$1 < \left(\frac{N_2}{\mathcal{N}_2(\psi(t_0))} \right)^{\frac{4}{d(p-1)-4}}. \tag{2.33}$$

On the other hand, (2.29), together with the definition of N_2, leads us to that $\mathcal{N}_2(\psi(t_0)) \geq N_2$, so that

$$\left(\frac{N_2}{\mathcal{N}_2(\psi(t_0))} \right)^{\frac{4}{d(p-1)-4}} \leq 1. \tag{2.34}$$

This inequality (2.34) contradicts (2.33). Thus, (2.28) must hold.

Once we obtain (2.28), we can easily obtain the following uniform bound:

$$\|\nabla\psi(t)\|_{L^2}^2 < \frac{d(p-1)}{d(p-1)-4}\mathcal{H}(\psi_0) \quad \text{for all } t \in I_{\max}. \tag{2.35}$$

Indeed, it follows from the energy conservation law (1.6) and (2.28) that

$$\begin{aligned} \mathcal{H}(\psi_0) = \mathcal{H}(\psi(t)) &= \|\nabla\psi(t)\|_{L^2}^2 - \frac{2}{p+1} \|\psi(t)\|_{L^{p+1}}^{p+1} \\ &> \frac{d(p-1)-4}{d(p-1)} \|\nabla\psi(t)\|_{L^2}^2 \quad \text{for all } t \in I_{\max}. \end{aligned} \tag{2.36}$$

The estimate (2.35), together with the sufficient condition for the blowup (1.4), leads us to that $I_{\max} = \mathbb{R}$. Hence, (2.25) and (2.26) follow from (2.28) and (2.35), respectively.

It remains to prove (2.27). The Gagliardo-Nirenberg inequality (1.25) gives us that

$$\begin{aligned} \mathcal{K}(\psi(t)) &= \|\nabla\psi(t)\|_{L^2}^2 - \frac{d(p-1)}{2(p+1)} \|\psi(t)\|_{L^{p+1}}^{p+1} \\ &\geq \|\nabla\psi(t)\|_{L^2}^2 - \frac{d(p-1)}{2(p+1)} \frac{1}{N_3}\mathcal{N}_2(\psi(t)) \|\nabla\psi(t)\|_{L^2}^2. \end{aligned} \tag{2.37}$$

Moreover, this inequality (2.37) and the relation $N_3 = \frac{d(p-1)}{2(p+1)}N_2$ (see (1.28)) yield that

$$\mathcal{K}(\psi(t)) \geq \|\nabla\psi(t)\|_{L^2}^2 - \frac{\mathcal{N}_2(\psi(t))}{N_2} \|\nabla\psi(t)\|_{L^2}^2. \tag{2.38}$$

Here, it follows from (2.26) that

$$\begin{aligned} \mathcal{N}_2(\psi(t)) &< \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \|\psi_0\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}(\psi_0)}^{\frac{d}{2}(p-1)-2} \\ &= \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \widetilde{\mathcal{N}_2}(\psi_0). \end{aligned} \tag{2.39}$$

Hence, combining (2.38), (2.39) and $\mathcal{H}(\psi(t)) > \mathcal{K}(\psi(t)) > 0$, we obtain that

$$\begin{aligned}\mathcal{K}(\psi(t)) &> \|\nabla\psi(t)\|_{L^2}^2 - \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \frac{\widetilde{\mathcal{N}}_2(\psi_0)}{N_2} \|\nabla\psi(t)\|_{L^2}^2 \\ &= \left(1 - \frac{\widetilde{\mathcal{N}}_2(\psi_0)}{\widetilde{N}_2}\right) \|\nabla\psi(t)\|_{L^2}^2 \\ &\geq \left(1 - \frac{\widetilde{\mathcal{N}}_2(\psi_0)}{\widetilde{N}_2}\right) \mathcal{H}(\psi(t)) = \left(1 - \frac{\widetilde{\mathcal{N}}_2(\psi_0)}{\widetilde{N}_2}\right) \mathcal{H}(\psi_0).\end{aligned} \tag{2.40}$$

□

Proposition 2.6 (Invariance of PW_-)**.** *Let $\psi_0 \in PW_-$ and ψ be the corresponding solution to the equation (1.1). Then, we have*

$$\psi(t) \in PW-, \quad \mathcal{K}(\psi(t)) < -\varepsilon_0 \qquad \textit{for all } t \in I_{\max}, \tag{2.41}$$

where $\varepsilon_0 = \mathcal{B}(\psi_0) - \mathcal{H}(\psi_0) > 0$.

Proof of Proposition 2.6. By the virtue of Lemma 2.4, it suffices to show that

$$\mathcal{K}(\psi(t)) < -\varepsilon_0 \qquad \text{for all } t \in I_{\max}. \tag{2.42}$$

We first prove that

$$\mathcal{K}(\psi_0) < -\varepsilon_0. \tag{2.43}$$

Supposing the contrary that $-\varepsilon_0 \leq \mathcal{K}(\psi_0)$, we have

$$\begin{aligned}0 &\leq \mathcal{K}(\psi_0) + \varepsilon_0 \\ &= \mathcal{H}(\psi_0) + \varepsilon_0 - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\} \|\psi_0\|_{L^{p+1}}^{p+1} \\ &= \mathcal{B}(\psi_0) - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\} \|\psi_0\|_{L^{p+1}}^{p+1}.\end{aligned} \tag{2.44}$$

Moreover, it follows from (2.44) and $\mathcal{K}(\psi_0) < 0$ that

$$\begin{aligned}0 &< \mathcal{B}(\psi_0) - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\} \frac{2(p+1)}{d(p-1)} \|\nabla\psi_0\|_{L^2}^2 \\ &= \mathcal{B}(\psi_0) - \frac{d(p-1)-4}{d(p-1)} \|\nabla\psi_0\|_{L^2}^2,\end{aligned} \tag{2.45}$$

so that

$$\|\nabla\psi_0\|_{L^2}^2 < \frac{d(p-1)}{d(p-1)-4} \mathcal{B}(\psi_0) = \left(\frac{N_2}{\|\psi_0\|_{L^2}^{p+1-\frac{d}{2}(p-1)}}\right)^{\frac{4}{d(p-1)-4}}. \tag{2.46}$$

Dividing the both sides of (2.46) by $\|\nabla\psi_0\|_{L^2}^2$, we obtain that

$$1 < \left(\frac{N_2}{\mathcal{N}_2(\psi_0)}\right)^{\frac{4}{d(p-1)-4}}. \tag{2.47}$$

On the other hand, since $\mathcal{K}(\psi_0) < 0$, the definition of N_2 (see (1.20)) implies that

$$\left(\frac{N_2}{\mathcal{N}_2(\psi_0)}\right)^{\frac{4}{d(p-1)-4}} \le 1, \tag{2.48}$$

which contradicts (2.47). Hence, we have proved (2.43).

Next, we prove (2.42). Suppose the contrary that (2.42) fails. Then, it follows from (2.43) and $\psi \in C(I_{\max}; H^1(\mathbb{R}^d))$ that there exists $t_1 \in I_{\max} \setminus \{0\}$ such that $-\varepsilon_0 = \mathcal{K}(\psi(t_1))$. This relation and the energy conservation law (1.6) lead us to that

$$\begin{aligned} 0 &= \mathcal{K}(\psi(t_1)) + \varepsilon_0 \\ &= \mathcal{H}(\psi_0) + \varepsilon_0 - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\}\|\psi(t_1)\|_{L^{p+1}}^{p+1} \\ &= \mathcal{B}(\psi_0) - \frac{2}{p+1}\left\{\frac{d}{4}(p-1) - 1\right\}\frac{2(p+1)}{d(p-1)}\left(\|\nabla\psi(t_1)\|_{L^2}^2 + \varepsilon_0\right) \\ &< \mathcal{B}(\psi_0) - \frac{d(p-1)-4}{d(p-1)}\|\nabla\psi(t_1)\|_{L^2}^2. \end{aligned} \tag{2.49}$$

Hence, we have that

$$\begin{aligned} \|\nabla\psi(t_1)\|_{L^2}^2 < \frac{d(p-1)}{d(p-1)-4}\mathcal{B}(\psi_0) &= \left(\frac{N_2}{\|\psi_0\|_{L^2}^{p+1-\frac{d}{2}(p-1)}}\right)^{\frac{4}{d(p-1)-4}} \\ &= \left(\frac{N_2}{\|\psi(t_1)\|_{L^2}^{p+1-\frac{d}{2}(p-1)}}\right)^{\frac{4}{d(p-1)-4}}, \end{aligned} \tag{2.50}$$

where we have used the mass conservation law (1.5) to derive the last equality. Then, an argument similar to the above yields a contradiction: Thus, we completed the proof. ☐

3 Strichartz type estimate and scattering

In this section, we introduce a certain space-time function space in addition to the usual Strichartz spaces, which enable us to control long-time behavior of solutions. Using this function space, we prepare two important propositions: Proposition 3.7 in Section 3.2 (small data theory) and Proposition 3.8 in Section 3.3 (long time perturbation theory). The former is used to avoid the vanishing and the latter to avoid the dichotomy in our concentration compactness like argument in Section 4.2. In the end of this section, we show the existence of the wave operators on PW_+.

3.1 Auxiliary function space X

In order to prove the scattering result ((1.50) in Theorem 1.1), we need to handle the inhomogeneous term of the integral equation associated with (1.1) in a suitable function space. Therefore, we will prepare a function space $X(I)$, $I \subset \mathbb{R}$, in which Strichartz type estimate works well.

Our equation (1.1) is invariant under the scaling

$$\psi(x,t) \mapsto \psi_\lambda(x,t) := \lambda^{\frac{2}{p-1}}\psi(\lambda x, \lambda^2 t), \tag{3.1}$$

which determines a critical regularity

$$s_p := \frac{d}{2} - \frac{2}{p-1}. \tag{3.2}$$

The condition (1.2) implies that $0 < s_p < 1$.

Throughout this paper, we fix a number q_1 with $p+1 < q_1 < 2^*$. Then, we define indices r_0, r_1 and $\widetilde{r}_1$ by

$$\frac{1}{r_0} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1}\right), \tag{3.3}$$

$$\frac{1}{r_1} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1} - \frac{s_p}{d}\right), \tag{3.4}$$

$$\frac{1}{\widetilde{r}_1} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_1} + \frac{s_p}{d}\right). \tag{3.5}$$

Here, the pair (q_1, r_0) is admissible. Besides these indices, we define a pair (q_2, r_2) by

$$\frac{p-1}{q_2} = 1 - \frac{2}{q_1}, \qquad \frac{1}{r_2} := \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_2} - \frac{s_p}{d}\right). \tag{3.6}$$

It is worth while noting that the Sobolev embedding and the Strichartz estimate lead us to the following estimate: For any pair (q, r) satisfying

$$\frac{d}{2}(p-1) \le q < 2^*, \qquad \frac{1}{r} = \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q} - \frac{s_p}{d}\right), \tag{3.7}$$

we have

$$\left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^r(I;L^q)} \lesssim \left\| (-\Delta)^{\frac{s_p}{2}} f \right\|_{L^2} \quad \text{for all } f \in \dot{H}^{s_p}(\mathbb{R}^d) \text{ and interval } I, \tag{3.8}$$

where the implicit constant depends only on d, p and q. The pairs (q_1, r_1) and (q_2, r_2) satisfy the condition (3.7), so that the estimate (3.8) is valid for these pairs.

Now, for any interval I, we put

$$X(I) = L^{r_1}(I; L^{q_1}) \cap L^{r_2}(I; L^{q_2}), \tag{3.9}$$

$$S(I) = L^\infty(I; L^2) \cap L^{r_0}(I; L^{q_1}). \tag{3.10}$$

We find that Strichartz type estimates work well in the space $X(I)$:

Lemma 3.1. *Assume that $d \ge 1$ and $2 + \frac{4}{d} < p+1 < 2^*$. Let $t_0 \in \mathbb{R}$ and I be an interval whose closure contains t_0. Then, we have*

$$\left\| \int_{t_0}^{t} e^{i(t-t')\Delta} v(t')\,dt' \right\|_{X(I)} \lesssim \|v\|_{L^{\widetilde{r}_1'}(I;L^{q_1'})}, \tag{3.11}$$

$$\left\| \int_{t_0}^{t} e^{i(t-t')\Delta} (v_1 v_2)(t')\,dt' \right\|_{X(I)} \lesssim \|v_1\|_{L^{r_1}(I;L^{q_1})} \|v_2\|_{L^{\frac{r_2}{p-1}}(I;L^{\frac{q_2}{p-1}})}, \tag{3.12}$$

where the implicit constants depend only on d, p and q_1.



Figure 1: Strichartz type estimates: $Q_0 : (\frac{1}{q_1}, \frac{1}{r_0})$, $Q_1 : (\frac{1}{q_1}, \frac{1}{r_1})$, $Q_2 : (\frac{1}{q_2}, \frac{1}{r_2})$, $\widetilde{Q}_1 : (\frac{1}{q_1}, \frac{1}{\widetilde{r}_1})$.

The estimate (3.11) in Lemma 3.1 is due to Foschi (see [17], Theorem 1.4). The estimate (3.12) is an immediate consequence of (3.11) and the Hölder inequality.

The following lemma is frequently used in the next section (Section 4); It is fundamental and easily obtained from the Hölder inequality and the chain rule:

Lemma 3.2. *Assume that* $d \ge 1$ *and* $2 + \frac{4}{d} < p + 1 < 2^*$. *Let* $t_0 \in \mathbb{R}$ *and* I *be an interval whose closure contains* t_0. *Then, we have*

$$\left\| |v|^{p-1} v \right\|_{L^{r_0'}(I;L^{q_1'})} \le \|v\|_{L^{r_0}(I;L^{q_1})} \|v\|_{L^{r_2}(I;L^{q_2})}^{p-1}, \tag{3.13}$$

$$\left\| \nabla \left(|v|^{p-1} v \right) \right\|_{L^{r_0'}(I;L^{q_1'})} \lesssim \|\nabla v\|_{L^{r_0}(I;L^{q_1})} \|v\|_{L^{r_2}(I;L^{q_2})}^{p-1}, \tag{3.14}$$

where the implicit constant depends only on d, p *and* q_1.

We also need the following interpolation estimate in the next section (Section 4.2):

Lemma 3.3. *For* $j \in \{1, 2\}$, *there exist a constant* $\theta_j \in (0, 1)$ *such that*

$$\left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^{r_j}(I;L^{q_j})} \lesssim \left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^\infty(I;L^{\frac{d}{2}(p-1)})}^{1-\theta_j} \left\| (-\Delta)^{\frac{s_p}{2}} f \right\|_{L^2}^{\theta_j} \quad \text{for all } f \in \dot{H}^{s_p}(\mathbb{R}^d),$$

where the implicit constant depends only on d, p *and* q_1.

Proof of Lemma 3.3. Fix a pair (q, r) satisfying (3.7) and $q_1 < q < 2^*$. Applying the

Hölder inequality first and (3.8) afterword, we obtain that

$$\begin{aligned}\left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^{r_j}(I;L^{q_j})} &\le \left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^\infty(I;L^{\frac{d}{2}(p-1)})}^{1-\theta_j} \left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^r(I;L^q)}^{\theta_j} \\ &\lesssim \left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^\infty(I;L^{\frac{d}{2}(p-1)})}^{1-\theta_j} \left\| (-\Delta)^{\frac{s_p}{2}} f \right\|_{L^2}^{\theta_j}, \quad j=1,2\end{aligned} \tag{3.15}$$

where

$$\theta_j := \frac{q}{q_j}\frac{2q_j - d(p-1)}{2q - d(p-1)},$$

and the implicit constant depends only on d, p and q_1 (we may ignore the dependence of q). ☐

At the end of this subsection, we record a decay estimate for the free solution:

Lemma 3.4. *Assume that $d \ge 1$. Then, we have*

$$\lim_{t\to\pm\infty} \left\| e^{\frac{i}{2}t\Delta} f \right\|_{L^q} = 0 \quad \text{for all } q \in (2, 2^*) \text{ and } f \in H^1(\mathbb{R}^d). \tag{3.16}$$

Proof of Lemma 3.16. This lemma is easily follows from the L^p-L^q-estimate for the free solution and the density of the Schwartz space $\mathcal{S}(\mathbb{R}^d)$ in $H^1(\mathbb{R}^d)$. ☐

3.2 Sufficient conditions for scattering

We shall give two sufficient conditions for solutions to have asymptotic states in the energy space $H^1(\mathbb{R}^d)$. One of them is the small data theory (see Proposition 3.7 and Remark 3.1). We also give the proof of Proposition 1.4 here.

We begin with the following proposition:

Proposition 3.5 (Scattering in the energy space)**.** *Assume that $d \ge 1$ and $2 + \frac{4}{d} < p + 1 < 2^*$. Let ψ be a solution to the equation (1.1).*
(i) *Suppose that ψ exists on $[0,\infty)$ and satisfies that*

$$\|\psi\|_{X([0,\infty))} < \infty, \qquad \|\psi\|_{L^\infty([0,\infty);H^1)} < \infty. \tag{3.17}$$

Then, there exists a unique asymptotic state $\phi_+ \in H^1(\mathbb{R}^d)$ such that

$$\lim_{t\to\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{H^1} = 0. \tag{3.18}$$

(ii) *Suppose that ψ exists on $(-\infty,0]$ and satisfies that*

$$\|\psi\|_{X((-\infty,0])} < \infty, \qquad \|\psi\|_{L^\infty((-\infty,0];H^1)} < \infty. \tag{3.19}$$

Then, there exists a unique asymptotic state $\phi_- \in H^1(\mathbb{R}^d)$ such that

$$\lim_{t\to-\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_- \right\|_{H^1} = 0. \tag{3.20}$$

For the proof of Proposition 3.5, we need the following lemma.

Lemma 3.6. *Assume that* $d \geq 1$ *and* $2 + \frac{4}{d} < p + 1 < 2^*$*. Let* I *be an interval and* ψ *be a solution to the equation (1.1) on* I*. Suppose that*

$$\|\psi\|_{X(I)} < \infty, \quad \|\psi\|_{L^\infty(I;H^1)} < \infty. \tag{3.21}$$

Then, we have that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{S(I)} < \infty.$$

Proof of Lemma 3.6. For the desired result, it suffices to show that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(I;L^q)} < \infty \tag{3.22}$$

for all pair (q,r) with $q_1 < q < 2^*$ and $\frac{1}{r} = \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q}\right)$ ((q,r) is an admissible pair). Indeed, the Hölder inequality, together with (3.21) and (3.22), gives us that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^{r_0}(I;L^{q_1})} \leq \|\psi\|_{L^\infty(I;H^1)}^{1-\frac{q(q_1-2)}{q_1(q-2)}} \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(I;L^q)}^{\frac{q(q_1-2)}{q_1(q-2)}} < \infty. \tag{3.23}$$

We shall prove (3.22). Let J be a subinterval of I with the property that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(J;L^q)} < \infty. \tag{3.24}$$

Then, applying the Strichartz estimate to the integral equation associated with (1.1), we obtain that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(J;L^q)} \lesssim \|\psi(t_0)\|_{H^1} + \left\|(1-\Delta)^{\frac{1}{2}}\left(|\psi|^{p-1}\psi\right)\right\|_{L^{r_0'}(J;L^{q_1'})}, \tag{3.25}$$

where the implicit constant depends only on d, p, q_1 and q. Moreover, it follows from Lemma 3.2 and the Hölder inequality that

$$\begin{aligned}\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(J;L^q)} &\lesssim \|\psi(t_0)\|_{H^1} + \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^{r_0}(J;L^{q_1})} \|\psi\|_{L^{r_2}(J;L^{q_2})}^{p-1} \\ &\leq \|\psi\|_{L^\infty(I;H^1)} + \|\psi\|_{L^\infty(J;H^1)}^{1-\theta} \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(J;L^q)}^{\theta} \|\psi\|_{X(J)}^{p-1},\end{aligned} \tag{3.26}$$

where

$$\theta := \frac{q(q_1-2)}{q_1(q-2)} \in (0,1)$$

and the implicit constant depends only on d, p, q_1 and q. This estimate (3.26) and the Young inequality show that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^r(J;L^q)} \lesssim \|\psi\|_{L^\infty(I;H^1)} + \|\psi\|_{L^\infty(I;H^1)} \|\psi\|_{X(I)}^{\frac{p-1}{1-\theta}}, \tag{3.27}$$

where the implicit constant depends only on d, p, q_1 and q. Since the right-hand side of (3.27) is independent of J, the condition (3.21) leads us to (3.22). ☐

Now, we give the proof of Proposition 3.5.

Proof of Proposition 3.5. Since the proofs of (i) and (ii) are very similar, we consider (i) only. A starting point is the following formula derived from the integral equation associated to (1.1):

$$e^{-\frac{i}{2}t\Delta}\psi(t) - e^{-\frac{i}{2}s\Delta}\psi(s) = \frac{i}{2}\int_{s}^{t} e^{-\frac{i}{2}t'\Delta}\left(|\psi|^{p-1}\psi\right)(t')dt' \quad \text{for all } s, t \in [0,\infty). \tag{3.28}$$

Applying the Strichartz estimate and Lemma 3.2 to this formula, we obtain that

$$\begin{aligned}\left\| e^{-\frac{i}{2}t\Delta}\psi(t) - e^{-\frac{i}{2}s\Delta}\psi(s)\right\|_{H^1} &\le \sup_{s'\in[s,t]}\left\| \int_{s}^{s'} e^{-\frac{i}{2}t'\Delta}\left(|\psi|^{p-1}\psi\right)(t')dt'\right\|_{H^1} \\ &\lesssim \left\|(1-\Delta)^{\frac{1}{2}}\left(|\psi|^{p-1}\psi\right)\right\|_{L^{r_0'}([s,t];L^{q_1'})} \\ &\lesssim \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{S([0,\infty))}\|\psi\|_{X([s,t])}^{p-1},\end{aligned} \tag{3.29}$$

where the implicit constant depends only on d, p and q_1. Then, it follows from Lemma 3.6 and the condition (3.17) that the right-hand side of (3.29) vanishes as $s, t \to \infty$, so that the completeness of $H^1(\mathbb{R}^d)$ leads us to that there exists $\phi_+ \in H^1(\mathbb{R}^d)$ such that

$$\lim_{t\to\infty}\left\|\psi(t) - e^{\frac{i}{2}t\Delta}\phi_+\right\|_{H^1} = \lim_{t\to\infty}\left\|e^{-\frac{i}{2}t\Delta}\psi(t) - \phi_+\right\|_{H^1} = 0. \tag{3.30}$$

To complete the proof, we shall show the uniqueness of a function ϕ_+ satisfying (3.30). Let ϕ_+ and ϕ_+' be functions in $H^1(\mathbb{R}^d)$ satisfying (3.30). Then, we easily verify that

$$\left\|\phi_+ - \phi_+'\right\|_{H^1} \le \left\|e^{-\frac{i}{2}t\Delta}\psi(t) - \phi_+\right\|_{H^1} + \left\|e^{-\frac{i}{2}t\Delta}\psi(t) - \phi_+'\right\|_{H^1} \quad \text{for all } t > 0. \tag{3.31}$$

Hence, taking $t \to \infty$, we have by (3.30) that $\phi_+ = \phi_+'$ in $H^1(\mathbb{R}^d)$. ☐

The following proposition gives us another sufficient condition for the boundedness of $X(I)$ and $S(I)$-norms.

Proposition 3.7 (Small data theory). *Assume that $d \ge 1$ and $2 + \frac{4}{d} < p + 1 < 2^*$. Let $t_0 \in \mathbb{R}$ and I be an interval whose closure contains t_0. Then, there exists a positive constant δ, depending only on d, p and q_1, with the following property: for any $\psi_0 \in H^1(\mathbb{R}^d)$ satisfying that*

$$\left\|e^{\frac{i}{2}(t-t_0)\Delta}\psi_0\right\|_{X(I)} \le \delta, \tag{3.32}$$

there exists a unique solution $\psi \in C(I; H^1(\mathbb{R}^d))$ to the equation (1.1) with $\psi(t_0) = \psi_0$ such that

$$\|\psi\|_{X(I)} < 2\left\|e^{\frac{i}{2}(t-t_0)\Delta}\psi_0\right\|_{X(I)}, \quad \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{S(I)} \lesssim \|\psi_0\|_{H^1}, \tag{3.33}$$

where the implicit constant depends only on d, p and q_1.

Remark 3.1. *Proposition 3.7, with the help of (3.8) and Proposition 3.5, shows a small data scattering, i.e., there exists $\varepsilon > 0$ such that for any $\psi_0 \in H^1(\mathbb{R}^d)$ with $\|\psi_0\|_{H^1} < \varepsilon$, the corresponding solution has an asymptotic state in $H^1(\mathbb{R}^d)$. For: It follows from (3.8) that there exists $\varepsilon > 0$ such that if $\|\psi_0\|_{H^1} < \varepsilon$, then $\left\|e^{\frac{i}{2}(t-t_0)\Delta}\psi_0\right\|_{X(\mathbb{R})} < \delta$ for δ found in Proposition 3.7. This fact, together with Proposition 3.5, yields the desired result.*

Proof of Proposition 3.7. We prove this lemma by the standard contraction mapping principle.

It follows from the Strichartz estimate that there exists a constant $C_0 > 0$, depending only on d, p and q_1, such that

$$\left\| e^{\frac{i}{2}t\Delta} f \right\|_{S(I)} \le C_0 \| f \|_{L^2} \quad \text{for all } f \in L^2(\mathbb{R}^d). \tag{3.34}$$

Using this constant, we define a set $Y(I)$ and a metric ρ there by

$$Y(I) := \left\{ u \in C(I; H^1(\mathbb{R}^d)) \;\middle|\; \begin{array}{l} \| u \|_{X(I)} \le 2 \left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)}, \\ \left\| (1-\Delta)^{\frac{1}{2}} u \right\|_{S(I)} \le 2C_0 \| \psi_0 \|_{H^1} \end{array} \right\}, \tag{3.35}$$

and

$$\rho(u,v) = \| u - v \|_{X(I) \cap S(I)}, \quad u, v \in Y(I). \tag{3.36}$$

We can verify that $(Y(I), \rho)$ is a complete metric space. Moreover, we define a map $\mathcal{T}$ on this space by

$$\mathcal{T}(u) := e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 + \frac{i}{2} \int_{t_0}^{t} e^{\frac{i}{2}(t-t')\Delta} |u(t')|^{p-1} u(t') \, dt', \quad u \in Y(I). \tag{3.37}$$

Now, let $\delta > 0$ be a constant to be specified later, and suppose that

$$\left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)} < \delta, \tag{3.38}$$

so that

$$\| u \|_{X(I)} \le 2\delta \quad \text{for all } u \in Y(I). \tag{3.39}$$

We shall show that $\mathcal{T}$ maps $Y(I)$ into itself for sufficiently small $\delta > 0$. Take any $u \in Y(I)$. Then, we have $\mathcal{T}(u) \in C(I; H^1(\mathbb{R}^d))$ (see [27]). Moreover, Lemma 3.1 and (3.39) yield that

$$\begin{aligned} \| \mathcal{T}(u) \|_{X(I)} &\le \left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)} + C_1 \| u \|_{X(I)}^{p} \\ &\le \left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)} + 2^p C_1 \delta^{p-1} \left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)} \end{aligned} \tag{3.40}$$

for some constant $C_1 > 0$ depending only on d, p and q_1. Hence, if we take δ so small that

$$2^p C_1 \delta^{p-1} \le 1, \tag{3.41}$$

then

$$\| \mathcal{T}(u) \|_{X(I)} \le 2 \left\| e^{\frac{i}{2}(t-t_0)\Delta} \psi_0 \right\|_{X(I)}. \tag{3.42}$$

On the other hand, it follows from the Strichartz estimate and Lemma 3.2 that

$$\begin{aligned} \left\| (1-\Delta)^{\frac{1}{2}} \mathcal{T}(u) \right\|_{S(I)} &\le C_0 \| \psi_0 \|_{H^1} + C_2 \left\| (1-\Delta)^{\frac{1}{2}} u \right\|_{S(I)} \| u \|_{X(I)}^{p-1} \\ &\le C_0 \| \psi_0 \|_{H^1} + 2^p C_2 C_0 \delta^{p-1} \| \psi_0 \|_{H^1} \end{aligned} \tag{3.43}$$

for some constant $C_2 > 0$ depending only on d, p and q_1. If δ satisfies that

$$2^p C_2 \delta^{p-1} \le 1, \tag{3.44}$$

then we have from (3.43) that

$$\left\| (1-\Delta)^{\frac{1}{2}} \mathcal{T}(u) \right\|_{S(I)} \le 2C_0 \left\| \psi_0 \right\|_{H^1}. \tag{3.45}$$

Thus, if we take δ satisfying (3.41) and (3.44), then $\mathcal{T}$ becomes a self-map on $Y(I)$.

Next, we shall show that $\mathcal{T}$ is a contraction map on $(Y(I), \rho)$ for sufficiently small δ. Lemma 3.1, together with (D.2) and (3.39), gives us that

$$\begin{aligned} \left\| \mathcal{T}(u) - \mathcal{T}(v) \right\|_{X(I)} &\le \left\| \int_{t_0}^{t} e^{\frac{i}{2}(t-t')\Delta} \left(|u|^{p-1}u - |v|^{p-1}v \right)(t')dt' \right\|_{X(I)} \\ &\le C_3 \left(\|u\|_{X(I)}^{p-1} + \|v\|_{X(I)}^{p-1} \right) \|u-v\|_{X(I)} \\ &\le 2^p C_3 \delta^{p-1} \|u-v\|_{X(I)} \quad \text{for all } u, v \in Y(I), \end{aligned} \tag{3.46}$$

where C_3 is some positive constant depending only on d, p and q_1. Similarly, we have by Lemma 3.2 that

$$\left\| \mathcal{T}(u) - \mathcal{T}(v) \right\|_{S(I)} \le 2^p C_4 \delta^{p-1} \|u-v\|_{S(I)} \tag{3.47}$$

for some constant $C_4 > 0$ depending only on d, p and q_1. Hence, we find that $\mathcal{T}$ becomes a contraction map in $(Y(I), \rho)$ for sufficiently small δ. Then, the contraction mapping principle shows that there exists a solution $\psi \in Y(I)$ to the equation (1.1) with $\psi(t_0) = \psi_0$, which, together with the uniqueness of solutions in $C(I; H^1(\mathbb{R}^d))$ (see [28]), completes the proof. □

At the end of this subsection, we give the proof of Proposition 1.4.

Proof of Theorem 1.4. Let ψ be a global solution to (1.1) with an initial datum $\psi_0 \in H^1(\mathbb{R}^d)$ at $t = 0$.
We shall prove that (i) implies (ii): Suppose that (i) holds. We first show the uniform boundedness:

$$\sup_{t \in [0,\infty)} \|\psi(t)\|_{H^1} < \infty. \tag{3.48}$$

The condition (i) yields that

$$\sup_{t \ge T} \|\psi(t)\|_{L^{p+1}} \le 1 \quad \text{for some } T > 0. \tag{3.49}$$

Combining this with the energy conservation law (1.6), we obtain that

$$\|\nabla \psi(t)\|_{L^2}^2 = \mathcal{H}(\psi(t)) - \|\psi(t)\|_{L^{p+1}}^{p+1} \le \mathcal{H}(\psi_0) + 1 \quad \text{for all } t \ge T. \tag{3.50}$$

Hence, this estimate and the mass conservation law (1.5) give (3.48).

Now, we shall prove (ii). We employ the Hölder inequality and the Sobolev embedding to obtain that

$$\|\psi(t)\|_{L^q} \lesssim \|\psi(t)\|_{L^{p+1}}^{1-\theta} \|\psi(t)\|_{H^1}^{\theta} \quad \text{for all } t \in [0,\infty) \text{ and } q \in (p+1, 2^*), \tag{3.51}$$

where θ is some constant in $(0,1)$, and the implicit constant is independent of t. Hence, the condition (i), together with (3.48), shows that

$$\lim_{t\to\infty} \|\psi(t)\|_{L^q} = 0 \quad \text{for all } q \in (p+1, 2^*). \tag{3.52}$$

On the other hand, we have by the Hölder inequality and the mass conservation law (1.5) that

$$\lim_{t\to\infty} \|\psi(t)\|_{L^q} \le \|\psi_0\|_{L^2}^{1-\frac{(q-2)(p+1)}{q(p-1)}} \lim_{t\to\infty} \|\psi(t)\|_{L^{p+1}}^{\frac{(q-2)(p+1)}{q(p-1)}} = 0 \quad \text{for all } q \in (2, p+1). \tag{3.53}$$

We shall prove (ii) implies (iii): We introduce a number $r_{2,0}$ such that $(q_2, r_{2,0})$ is admissible, i.e.,

$$\frac{1}{r_{2,0}} = \frac{d}{2}\left(\frac{1}{2} - \frac{1}{q_2}\right), \tag{3.54}$$

where q_2 is the number defined in (3.6). Then, we have that

$$2 < r_{2,0} < r_2 < \infty \tag{3.55}$$

for r_2 defined in (3.6). Moreover, we define a number q_e by $q_e = 2^*$ if $d \ge 3$ and $q_e = 2q_1 (> q_2)$ if $d = 1, 2$, and a space $\bar{S}(I)$ by

$$\bar{S}(I) = \bigcap_{\substack{(q,r):admissible \\ 2\le q\le q_e}} L^r(I; L^q) \quad \text{for an interval } I. \tag{3.56}$$

Then, it follows from the Strichartz estimate and Lemma 3.2 that

$$\begin{aligned}
&\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{\bar{S}([t_0,t_1))} \\
&\lesssim \|\psi(t_0)\|_{H^1} + \left\|(1-\Delta)^{\frac{1}{2}}|\psi|^{p-1}\psi\right\|_{L^{r_0'}([t_0,t_1);L^{q_1'})} \\
&\le \|\psi\|_{L^\infty([0,\infty);H^1)} + \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^{r_0}([t_0,t_1);L^{q_1})} \|\psi\|_{L^{r_2}([t_0,t_1);L^{q_2})}^{p-1} \\
&\le \|\psi\|_{L^\infty([0,\infty);H^1)} + \sup_{t\ge t_0} \|\psi(t)\|_{L^{q_2}}^{\frac{p-1}{r_2}(r_2-r_{2,0})} \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{L^{r_0}([t_0,t_1);L^{q_1})} \|\psi\|_{L^{r_{2,0}}([t_0,t_1);L^{q_2})}^{\frac{p-1}{r_2}r_{2,0}} \\
&\le \|\psi\|_{L^\infty([0,\infty);H^1)} + \sup_{t\ge t_0} \|\psi(t)\|_{L^{q_2}}^{\frac{p-1}{r_2}(r_2-r_{2,0})} \left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{\bar{S}([t_0,t_1);L^{q_1})}^{1+\frac{p-1}{r_2}r_{2,0}},
\end{aligned} \tag{3.57}$$

where the implicit constant is independent of t_0 and t_1. Note here that since the condition (ii) includes (i), we have that $\|\psi\|_{L^\infty([0,\infty);H^1)} < \infty$ (see (3.48) above). Hence, the estimate (3.57), together with the condition (ii), shows that

$$\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{\bar{S}([t_0,+\infty))} < \infty \quad \text{for all sufficiently large } t_0 > 0, \tag{3.58}$$

so that (iii) holds.

We shall prove that (iii) implies (iv): The estimate (3.8) immediately gives us the desired result.

We shall prove that (iv) implies (v): This follows from Proposition 3.5.

We shall prove that (v) implies (i): We define r by

$$\frac{1}{r} = \frac{d}{2}\left(\frac{1}{2} - \frac{1}{p+1}\right), \tag{3.59}$$

so that $(p+1, r)$ is an admissible pair. Then, the Strichartz estimate yields that

$$\left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{L^r(\mathbb{R};L^{p+1})} \lesssim \|\phi_+\|_{L^2}, \tag{3.60}$$

where the implicit constant depends only on d and p. Hence, we have that

$$\liminf_{t\to\infty} \left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{L^{p+1}} = 0. \tag{3.61}$$

Moreover, it follows from the existence of an asymptotic state ϕ_+ and (3.61) that

$$\liminf_{t\to\infty} \|\psi(t)\|_{L^{p+1}} \le \lim_{t\to\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{L^{p+1}} + \liminf_{t\to\infty} \left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{L^{p+1}} = 0. \tag{3.62}$$

Using the existence of an asymptotic state again, we also obtain that

$$\lim_{t\to\infty} \|\nabla\psi(t)\|_{L^2} = \lim_{t\to\infty} \left\| \nabla e^{-\frac{i}{2}t\Delta}\psi(t) \right\|_{L^2} = \|\nabla\phi_+\|_{L^2}. \tag{3.63}$$

This formula (3.63) and the energy conservation law (1.6) lead us to that

$$\lim_{t\to\infty} \|\psi(t)\|_{L^{p+1}}^{p+1} = -\frac{p+1}{2}\mathcal{H}(\psi(0)) + \frac{p+1}{2}\|\nabla\phi_+\|_{L^2}^2. \tag{3.64}$$

In particular, $\lim_{t\to\infty} \|\psi(t)\|_{L^{p+1}}^{p+1}$ exists. Hence, the desired result (i) follows from (3.62). $\square$

3.3 Long time perturbation theory

We will employ the so-called "Born type approximation" to prove that the solutions starting from PW_+ have asymptotic states (see Section 4). The following proposition plays a crucial role there.

Proposition 3.8 (Long time perturbation theory)**.** *Assume that $d \ge 1$ and $2 + \frac{4}{d} < p + 1 < 2^*$. Then, for any $A > 1$, there exists $\varepsilon > 0$, depending only on A, d, p and q_1, such that the following holds: Let I be an interval, and u be a function in $C(I; H^1(\mathbb{R}^d))$ such that*

$$\|u\|_{X(I)} \le A, \tag{3.65}$$

$$\left\| 2i\partial_t u + \Delta u + |u|^{p-1}u \right\|_{L^{\tilde{r}_1'}(I;L^{q_1'})} \le \varepsilon. \tag{3.66}$$

If $\psi \in C(\mathbb{R}; H^1)$ is a global solution to the equation (1.1) and satisfies that

$$\left\| e^{\frac{i}{2}(t-t_1)\Delta}(\psi(t_1) - u(t_1)) \right\|_{X(I)} \le \varepsilon \quad \textit{for some } t_1 \in I, \tag{3.67}$$

then we have

$$\|\psi\|_{X(I)} \lesssim 1, \tag{3.68}$$

where the implicit constant depends only on d, p, q_1 and A.

Remark 3.2. *We can find from the proof below that Proposition 3.8 still holds valid if we replace* ψ *with a function* $\widetilde{\psi} \in C(\mathbb{R}; H^1(\mathbb{R}^d))$ *satisfying that*

$$\left\| 2i\partial_t \widetilde{\psi} + \Delta \widetilde{\psi} + |\widetilde{\psi}|^{p-1}\widetilde{\psi} \right\|_{L^{\widetilde{r}_1'}(I;L^{q_1'})} \le \varepsilon. \tag{3.69}$$

Proof of Proposition 3.8. Let u be a function in $C(I; H^1(\mathbb{R}^d)$ satisfying (3.65) and (3.66) for $\varepsilon > 0$ to be chosen later. Moreover, let ψ be a global solution to the equation (1.1) satisfying (3.67).

For simplicity, we suppose that $I = [t_1, \infty)$ in what follows. The other cases are treated in a way similar to this case.

We have from the condition (3.65) that: for any $\delta > 0$, there exist disjoint intervals $I_1, \dots, I_N$, with a form $I_j = [t_j, t_{j+1})$ $(t_{N+1} = \infty)$, such that

$$I = \bigcup_{j=1}^{N} I_j, \tag{3.70}$$

and

$$\|u\|_{X(I_j)} \le \delta \quad \text{for all } j = 1, \dots, N, \tag{3.71}$$

where N is some number depending only on δ, A, d, p and q_1.

We put

$$w := \psi - u, \qquad e := 2i\partial_t u + \Delta u + |u|^{p-1}u, \tag{3.72}$$

Then, w satisfies that

$$2i\partial_t w + \Delta w + |w+u|^{p-1}(w+u) - |u|^{p-1}u + e = 0, \tag{3.73}$$

$$\left\| e^{\frac{i}{2}(t-t_1)\Delta} w(t_1) \right\|_{X(I)} \le \varepsilon. \tag{3.74}$$

We consider the integral equations associated with (3.73):

$$w(t) = e^{\frac{i}{2}(t-t_j)\Delta} w(t_j) + \frac{i}{2}\int_{t_j}^{t} e^{\frac{i}{2}(t-t')\Delta} W(t')\,dt', \quad j = 1, \dots, N, \tag{3.75}$$

where

$$W := |w+u|^{p-1}(w+u) - |u|^{p-1}u - e. \tag{3.76}$$

It follows from the inhomogeneous Strichartz estimate (3.11), the elementary inequality (D.2) and the Hölder inequality that

$$\begin{aligned} &\left\| \int_{t_j}^{t} e^{\frac{i}{2}(t-t')\Delta} W(t')\,dt' \right\|_{X([t_j,t_j'))} \\ &\le \mathcal{C}\Big\{ \|w\|_{L^{r_1}([t_j,t_j');L^{q_1})} \|u\|_{L^{r_2}([t_j,t_j');L^{q_2})}^{p-1} + \|w\|_{L^{r_1}([t_j,t_j');L_x^{q_1})} \|w\|_{L^{r_2}([t_j,t_j');L^{q_2})}^{p-1} \\ &\qquad\qquad + \|e\|_{L^{\widetilde{r}_1'}([t_j,t_j');L^{q_1'})} \Big\} \qquad \text{for all } j = 1, \dots, N \text{ and } t_j' \in I_j, \end{aligned} \tag{3.77}$$

where $\mathcal{C}$ is some constant depending only on d, p and q_1. Hence, combining this estimate with (3.66) and (3.71), we obtain that

$$\|w\|_{X([t_j,t_j'))} \le \left\| e^{\frac{i}{2}(t-t_j)\Delta} w(t_j) \right\|_{X(I_j)} + \mathcal{C}\left\{ \delta^{p-1} \|w\|_{X([t_j,t_j'))} + \|w\|^p_{X([t_j,t_j'))} + \varepsilon \right\} \tag{3.78}$$
$$\text{for all } j = 1, \dots, N \text{ and } t_j' \in I_j.$$

Now, we fix a constant δ such that

$$\delta < \left(\frac{1}{4(1+2\mathcal{C})} \right)^{\frac{1}{p-1}}, \tag{3.79}$$

so that the number N is also fixed. Then, we shall show that

$$\left\| e^{\frac{i}{2}(t-t_j)\Delta} w(t_j) \right\|_{X(I)} \le (1+2^j\mathcal{C})\varepsilon \qquad \text{for all } j = 1, \dots, N, \tag{3.80}$$

$$\|w\|_{X(I_j)} \le \left(1+2^{j+1}\mathcal{C}\right)\varepsilon \qquad \text{for all } j = 1, \dots, N, \tag{3.81}$$

provided that

$$\varepsilon < \left(\frac{1}{4(1+2^{N+1}\mathcal{C})^p} \right)^{\frac{1}{p-1}}. \tag{3.82}$$

We first consider the case $j = 1$. The estimate (3.80) for $j = 1$ obviously follows from (3.74). We put

$$\bar{t}_1 = \sup\left\{ t_1 \le t \le t_2 \mid \|w\|_{X([t_1,t])} \le (1+4\mathcal{C})\varepsilon \right\}. \tag{3.83}$$

Then, the estimate (3.78), together with (3.74), shows that $\bar{t}_1 > t_1$. Supposing the contrary that $\bar{t}_1 < t_2$, we have from the continuity of w that

$$\|w\|_{X([t_1,\bar{t}_1])} = (1+4\mathcal{C})\varepsilon. \tag{3.84}$$

However, (3.78), together with (3.80) with $j = 1$ and (3.84), leads us to that

$$\begin{aligned} \|w\|_{X([t_1,\bar{t}_1])} &\le (1+2\mathcal{C})\,\varepsilon + \mathcal{C}\left\{ \delta^{p-1}(1+4\mathcal{C})\varepsilon + (1+4\mathcal{C})^p\varepsilon^p + \varepsilon \right\} \\ &< \varepsilon + 4\mathcal{C}\varepsilon. \end{aligned} \tag{3.85}$$

This contradicts (3.84). Thus, it must hold that $\bar{t}_1 = t_2$ and therefore (3.81) holds for $j = 1$.

We shall prove the general case $j = n$ ($2 \le n \le N$), provided that both (3.80) and (3.81) hold for all $j = 1, \dots, n-1$.

Multiplying the integral equation (3.75) with $j = n$ and $t = t_n$ by $e^{\frac{i}{2}(t-t_n)\Delta}$, we obtain that

$$e^{\frac{i}{2}(t-t_n)\Delta} w(t_n) = e^{\frac{i}{2}(t-t_{n-1})\Delta} w(t_{n-1}) + \frac{i}{2}\int_{t_{n-1}}^{t_n} e^{\frac{i}{2}(t-t')\Delta} \chi_{I_{n-1}}(t') W(t')\, dt', \tag{3.86}$$

where $\chi_{I_{n-1}}$ is the characteristic function on I_{n-1}. Then, the same estimate as (3.78) yields that

$$\begin{aligned} \left\| e^{\frac{i}{2}(t-t_n)\Delta} w(t_n) \right\|_{X(I)} &\le \left\| e^{\frac{i}{2}(t-t_{n-1})\Delta} w(t_{n-1}) \right\|_{X(I)} \\ &\quad + \mathcal{C}\left\{ \delta^{p-1} \|w\|_{X(I_{n-1})} + \|w\|^p_{X(I_{n-1})} + \varepsilon \right\} \end{aligned} \tag{3.87}$$

for the same constant $\mathcal{C}$ found in (3.78). This estimate, together with the inductive hypothesis, gives us that

$$\begin{aligned}\left\|e^{\frac{i}{2}(t-t_n)\Delta}w(t_n)\right\|_{X(I)} &< (1+2^{n-1}\mathcal{C})\varepsilon + \mathcal{C}\left\{2^{n-3}\varepsilon + \frac{1}{4}\varepsilon + \varepsilon\right\} \\ &\le (1+2^{n}\mathcal{C})\varepsilon,\end{aligned} \tag{3.88}$$

which shows that (3.80) holds for $j = n$.

Next, we consider (3.81) for $j = n$. As well as the case $j = 1$, we put

$$\bar{t}_n = \sup\left\{t_n \le t \le t_{n+1} \;\middle|\; \|w\|_{X([t_n,t])} \le (1+2^{n+1}\mathcal{C})\varepsilon\right\}. \tag{3.89}$$

The estimate (3.78), together with (3.88), shows that $t_n < \bar{t}_n$. We suppose the contrary that $\bar{t}_n < t_{n+1}$, so that

$$\|w\|_{X([t_n,\bar{t}_n])} = \left(1+2^{n+1}\mathcal{C}\right)\varepsilon. \tag{3.90}$$

Then, combining (3.78) with (3.88) and (3.90), we obtain that

$$\|w\|_{X([t_n,\bar{t}_n])} < (1+2^{n}\mathcal{C})\,\varepsilon + \mathcal{C}\left\{2^{n-2}\varepsilon + \frac{1}{4}\varepsilon + \varepsilon\right\} \le \left(1+2^{n+1}\mathcal{C}\right)\varepsilon, \tag{3.91}$$

which contradicts (3.90). Hence, we have proved (3.80) and (3.81).

Now, it follows from (3.81) and (3.82) that

$$\|w\|^{r_j}_{L^{r_j}(I;L^{q_j})} = \sum_{n=1}^{N} \|w\|^{r_j}_{L^{r_j}(I_n;L^{q_j})} \le \sum_{n=1}^{N} \left(1+2^{N+1}\mathcal{C}\right)^{r_j}\varepsilon^{r_j} \le \frac{N}{4^{r_j}} \le N^{r_j} \quad \text{for } j = 1,2. \tag{3.92}$$

Hence, we have from (3.65) and (3.92) that

$$\|\psi\|_{X(I)} \le \|u\|_{X(I)} + \|w\|_{X(I)} \le A + N, \tag{3.93}$$

which completes the proof. □

3.4 Wave operators

The following proposition tells us that the wave operators are well-defined on Ω.

Proposition 3.9 (Existence of wave operators). *Assume $d \ge 1$ and $2 + \frac{4}{d} \le p+1 < 2^*$.*
(i) *For any $\phi_+ \in \Omega$, there exists a unique $\psi_0 \in PW_+$ such that the corresponding solution ψ to the equation (1.1) with $\psi(0) = \psi_0$ exists globally in time and satisfies the followings:*

$$\psi \in X([0,+\infty)), \tag{3.94}$$

$$\lim_{t\to+\infty}\left\|\psi(t) - e^{\frac{i}{2}t\Delta}\phi_+\right\|_{H^1} = 0, \tag{3.95}$$

$$\mathcal{H}(\psi(t)) = \|\nabla\phi_+\|^2_{L^2} \quad \textit{for all } t \in \mathbb{R}. \tag{3.96}$$

Furthermore, if $\|\phi_+\|_{H^1}$ is sufficiently small, then we have

$$\|\psi\|_{X(\mathbb{R})} \lesssim \|\phi_+\|_{H^1}, \tag{3.97}$$

where the implicit constant depends only on d, p *and* q_1.
The map defined by $\phi_+ \mapsto \psi_0$ *is continuous from* Ω *into* PW_+ *in the* $H^1(\mathbb{R}^d)$*-topology.*

(ii) *For any* $\phi_- \in \Omega$, *there exists a unique* $\psi_0 \in PW_+$ *such that the corresponding solution* ψ *to the equation (1.1) with* $\psi(0) = \psi_0$ *exists globally in time and satisfies that*

$$\psi \in X((-\infty, 0]), \tag{3.98}$$

$$\lim_{t \to -\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_- \right\|_{H^1} = 0, \tag{3.99}$$

$$\mathcal{H}(\psi(t)) = \|\nabla \phi_-\|_{L^2}^2 \quad \textit{for all } t \in \mathbb{R}. \tag{3.100}$$

Furthermore, if $\|\phi_-\|_{H^1}$ *is sufficiently small, then we have*

$$\|\psi\|_{X(\mathbb{R})} \lesssim \|\phi_-\|_{H^1}, \tag{3.101}$$

where the implicit constant depends only on d, p *and* q_1.
The map defined by $\phi_- \mapsto \psi_0$ *is continuous from* Ω *into* PW_+ *in the* $H^1(\mathbb{R}^d)$*-topology.*

Proof of Proposition 3.9. Since the proofs of (i) and (ii) are very similar, we prove (i) only. We look for a solution to the following integral equation in $X([0,\infty))$ for all $\phi_+ \in \Omega$:

$$\psi(t) = e^{\frac{i}{2}t\Delta}\phi_+ - \frac{i}{2}\int_t^{+\infty} e^{\frac{i}{2}(t-t')\Delta}\left\{|\psi(t')|^{p-1}\psi(t')\right\} dt'. \tag{3.102}$$

We first note that the estimate (3.8) shows that: For any $\delta > 0$, there exists $T_\delta > 0$ such that

$$\left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{X([T_\delta, +\infty))} < \delta. \tag{3.103}$$

Moreover, it follows from the Strichartz estimate that: There exists $C_0 > 0$, depending only on d, p and q_1, such that

$$\left\| (1-\Delta)^{\frac{1}{2}} e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{S(\mathbb{R})} \le C_0 \|\phi_+\|_{H^1}. \tag{3.104}$$

Using these constants T_δ and C_0, we define a set $Y_\delta[\phi_+]$ by

$$Y_\delta[\phi_+] := \left\{ u \in C(I_\delta; H^1(\mathbb{R}^d)) \;\middle|\; \begin{array}{l} \|u\|_{X(I_\delta)} \le 2\left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{X(I_\delta)}, \\ \left\| (1-\Delta)^{\frac{1}{2}} u \right\|_{S(I_\delta)} \le 2C_0 \|\phi_+\|_{H^1} \end{array} \right\}, \tag{3.105}$$

where $I_\delta = [T_\delta, \infty)$. We can verify that $Y_\delta[\phi_+]$ becomes a complete metric space with a metric ρ defined by

$$\rho(u,v) := \max\left\{ \|u - v\|_{X(I_\delta)}, \; \|u - v\|_{S(I_\delta)} \right\} \quad \text{for all } u, v \in Y_\delta[\phi_+]. \tag{3.106}$$

Then, an argument similar to the proof of Proposition 3.7 (small data theory) leads us to that: If $\delta > 0$ is sufficiently small, then there exists a unique solution $\psi \in Y_\delta[\phi_+]$ to the integral equation (3.102). In particular, we find that such a solution exists globally in time and satisfies (3.101), provided that $\|\phi_+\|_{H^1}$ is sufficiently small.

We shall show that the function ψ obtained above becomes a solution to the equation (1.1). Multiplying the both sides of (3.102) by the free operator, we have

$$e^{\frac{i}{2}(t-T)\Delta}\psi(T) = e^{\frac{i}{2}t\Delta}\phi_{+} - \frac{i}{2}\int_{T}^{+\infty} e^{\frac{i}{2}(t-t')\Delta}\left\{|\psi(t')|^{p-1}\psi(t')\right\}dt' \quad \text{for all } t, T \in I_{\delta}. \tag{3.107}$$

Subtraction (3.107) from (3.102) yields further that

$$\psi(t) = e^{\frac{i}{2}(t-T)\Delta}\psi(T) + \frac{i}{2}\int_{T}^{t} e^{\frac{i}{2}(t-t')\Delta}\left\{|\psi(t')|^{p-1}\psi(t')\right\}dt' \quad \text{for all } t, T \in I_{\delta}. \tag{3.108}$$

Hence, we find by this formula (3.108) that ψ is a solution to the equation (1.1) on I_{δ}.

Next, we shall extend the solution ψ to the whole interval $\mathbb{R}$. In view of Proposition 2.5, it suffices to show that $\psi(t) \in PW_{+}$ for some $t \in I_{\delta}$. Applying the usual inhomogeneous Strichartz estimate to (3.102), and then applying Lemma 3.2, we obtain that

$$\lim_{t\to+\infty}\left\|\psi(t) - e^{\frac{i}{2}t\Delta}\phi_{+}\right\|_{H^1} \lesssim \lim_{t\to+\infty}\left\|(1-\Delta)^{\frac{1}{2}}\psi\right\|_{S([t,\infty))}\|\psi\|_{X([t,\infty))}^{p-1} = 0. \tag{3.109}$$

Using this estimate (3.109) and Lemma 3.4, we conclude that

$$\begin{aligned}\lim_{t\to+\infty}\mathcal{K}(\psi(t)) &= \lim_{t\to+\infty}\mathcal{K}(e^{\frac{i}{2}t\Delta}\phi_{+}) \\ &= \lim_{t\to+\infty}\left\{\mathcal{K}(\phi_{+}) - \frac{d(p-1)}{2(p+1)}\left\|e^{\frac{i}{2}t\Delta}\phi_{+}\right\|_{L^{p+1}}^{p+1} + \frac{d(p-1)}{2(p+1)}\|\phi_{+}\|_{L^{p+1}}^{p+1}\right\} \\ &= \mathcal{K}(\phi_{+}) + \frac{d(p-1)}{2(p+1)}\|\phi_{+}\|_{L^{p+1}}^{p+1} > 0.\end{aligned} \tag{3.110}$$

Hence, it holds that

$$0 < \mathcal{K}(\psi(t)) < \mathcal{H}(\psi(t)) \quad \text{for all sufficiently large } t > T_{\delta}. \tag{3.111}$$

Moreover, using (3.109) and Lemma 3.4 again, and employing the condition $\phi_{+} \in \Omega$, we obtain that

$$\begin{aligned}&\lim_{t\to+\infty}\widetilde{\mathcal{N}}_2(\psi(t)) = \lim_{t\to+\infty}\widetilde{\mathcal{N}}_2(e^{\frac{i}{2}t\Delta}\phi_{+}) \\ &= \lim_{t\to+\infty}\|\phi_{+}\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\sqrt{\|\nabla\phi_{+}\|_{L^2}^2 - \frac{2}{p+1}\left\|e^{\frac{i}{2}t\Delta}\phi_{+}\right\|_{L^{p+1}}^{p+1}}^{\frac{d}{2}(p-1)-2} \\ &= \|\phi_{+}\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla\phi_{+}\|_{L^2}^{\frac{d}{2}(p-1)-2} = \mathcal{N}_2(\phi_{+}) < \widetilde{N}_2.\end{aligned} \tag{3.112}$$

This estimate, together with (1.43), shows that

$$\psi(t) \in PW_{+} \quad \text{for all sufficiently large } t > T_{\delta}. \tag{3.113}$$

Thus, we have proved the global existence of ψ. Since $\psi \in X([T,+\infty))$ for sufficiently large $T > 0$, we also have $\psi \in X([0,+\infty))$.

Now, we put $\psi_0 = \psi(0)$. Then, ψ_0 is what we want. Indeed, the desired properties (3.94) and (3.95) have been obtained. Moreover, (3.95) and Lemma 3.4 immediately give (3.96).

We shall prove the uniqueness here. Let ψ_0 and $\widetilde{\psi}_0$ be functions in PW_+ such that the solutions ψ and $\widetilde{\psi}$ to the equation (1.1) with $\psi(0) = \psi_0$ and $\widetilde{\psi}(0) = \widetilde{\psi}_0$ satisfy that

$$\lim_{t\to\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{H^1} = \lim_{t\to\infty} \left\| \widetilde{\psi}(t) - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{H^1} = 0. \tag{3.114}$$

Using (3.114), we find that

$$\lim_{t\to\infty} \left\| \psi(t) - \widetilde{\psi}(t) \right\|_{H^1} = 0. \tag{3.115}$$

Then, supposing the contrary that $\psi_0 \neq \widetilde{\psi}_0$, we have by the standard uniqueness result (see [28]) that $\psi(t) \neq \widetilde{\psi}(t)$ for all $t \in \mathbb{R}$, which contradicts (3.115): Thus, $\psi_0 = \widetilde{\psi}_0$.

Finally, we prove the continuity of the map defined by $\phi_+ \in \Omega \mapsto \psi_0 \in PW_+$. We use W_+ to denote this map. Let $\phi_+ \in \Omega$, and let $\{\phi_{+,n}\}_{n\in\mathbb{N}}$ be a sequence in Ω satisfying that

$$\lim_{n\to\infty} \|\phi_{+,n} - \phi_+\|_{H^1} = 0. \tag{3.116}$$

The estimate (3.8), together with (3.116), shows that

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\phi_{+,n} - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{X(\mathbb{R})} = 0. \tag{3.117}$$

Let ψ and ψ_n be the solutions to (1.1) with $\psi(0) = W_+\phi_+$ and $\psi_n(0) = W_+\phi_{+,n}$, respectively. Then, ψ and ψ_n exist globally in time (see Proposition 2.5) and have the following properties as proved above:

$$\psi(t), \psi_n(t) \in PW_+ \quad \text{for all } t \in \mathbb{R}, \tag{3.118}$$

$$\psi, \psi_n \in X([0,\infty)), \tag{3.119}$$

$$\lim_{t\to+\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{H^1} = \lim_{t\to+\infty} \left\| \psi_n(t) - e^{\frac{i}{2}t\Delta}\phi_{+,n} \right\|_{H^1} = 0, \tag{3.120}$$

$$\mathcal{H}(\psi(t)) = \|\nabla\phi_+\|_{L^2}^2, \quad \mathcal{H}(\psi_n(t)) = \|\nabla\phi_{n,+}\|_{L^2}^2 \quad \text{for all } t \in \mathbb{R}. \tag{3.121}$$

Moreover, we find by (3.120) that ψ and ψ_n satisfy that

$$\psi(t) = e^{\frac{i}{2}t\Delta}\phi_+ - \frac{i}{2}\int_t^{+\infty} e^{\frac{i}{2}(t-t')\Delta} \left\{ |\psi(t')|^{p-1}\psi(t') \right\} dt', \tag{3.122}$$

$$\psi_n(t) = e^{\frac{i}{2}t\Delta}\phi_{+,n} - \frac{i}{2}\int_t^{+\infty} e^{\frac{i}{2}(t-t')\Delta} \left\{ |\psi_n(t')|^{p-1}\psi_n(t') \right\} dt'. \tag{3.123}$$

Note here that it follows from (3.116) and (3.117) that there exists a number $n_0 \in \mathbb{N}$ such that

$$\left\| e^{\frac{i}{2}t\Delta}\phi_{+,n} \right\|_{X(I)} \leq 2 \left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{X(I)} \quad \text{for all interval } I \text{ and } n \geq n_0, \tag{3.124}$$

$$\|\phi_{+,n}\|_{H^1} \leq 2\|\phi_+\|_{H^1} \quad \text{for all } n \geq n_0. \tag{3.125}$$

We find by (3.124) that: For any $\delta > 0$, there exists $T_\delta > 0$, independent of n, such that

$$\left\| e^{\frac{i}{2}t\Delta}\phi_+ \right\|_{X([T_\delta,\infty))} < \delta, \qquad \left\| e^{\frac{i}{2}}t\Delta\phi_{+,n} \right\|_{X([T_\delta,\infty))} \le \delta \quad \text{for all } n \ge n_0. \tag{3.126}$$

Using (3.126) and the formulas (3.122) and (3.123), we also find that: There exists a constant $\delta_0 > 0$ with the following property: for any $\delta \in (0, \delta_0]$, there exists $T_\delta > 0$ such that

$$\|\psi\|_{X([T_\delta,\infty))} < 2\delta, \qquad \|\psi_n\|_{X([T_\delta,\infty))} \le 2\delta \quad \text{for all } n \ge n_0. \tag{3.127}$$

Now, we consider an estimate for the difference $\psi_n - \psi$. The formulas (3.122) and (3.123) shows that

$$\psi_n(t) - \psi(t) = e^{\frac{i}{2}t\Delta}\left(\phi_{+,n} - \phi_+\right) - \frac{i}{2}\int_t^{+\infty} e^{\frac{i}{2}(t-t')\Delta}\left\{|\psi_n|^{p-1}\psi_n - |\psi|^{p-1}\psi\right\}(t')\,dt'. \tag{3.128}$$

Applying the Strichartz estimate to (3.128), and using (D.2) and (3.127), we obtain that

$$\begin{aligned}
&\|\psi_n - \psi\|_{S([T_\delta,\infty))} \\
&\lesssim \|\phi_{+,n} - \phi_+\|_{L^2} + \left(\|\psi_n\|_{X([T_\delta,\infty))}^{p-1} + \|\psi\|_{X([T_\delta,\infty))}^{p-1}\right)\|\psi_n - \psi\|_{S([T_\delta,\infty))} \\
&\le \|\phi_{+,n} - \phi_+\|_{L^2} + 2^p\delta^{p-1}\|\psi_n - \psi\|_{S([T_\delta,\infty))} \quad \text{for all } n \ge n_0,
\end{aligned} \tag{3.129}$$

where the implicit constant depends only on d, p and q_1. This estimate, together with (3.116), shows that

$$\lim_{n\to\infty}\|\psi_n - \psi\|_{S([T_\delta,\infty))} = 0 \quad \text{for all sufficiently small } \delta > 0. \tag{3.130}$$

Similarly, we can obtain that

$$\lim_{n\to\infty}\|\psi_n - \psi\|_{X([T_\delta,\infty))} = 0 \quad \text{for all sufficiently small } \delta > 0. \tag{3.131}$$

Moreover, considering the integral equations of $\partial_j\psi$ and $\partial_j\psi_n$ for $1 \le j \le d$, we obtain by the Strichartz estimate that

$$\begin{aligned}
\|\partial_j\psi_n - \partial_j\psi\|_{S([T_\delta,\infty))} &\lesssim \|\partial_j\phi_{+,n} - \partial_j\phi_+\|_{L^2} + \left\||\psi_n|^{p-1}(\partial_j\psi_n - \partial_j\psi)\right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\
&\quad + \left\|\left(|\psi_n|^{p-1} - |\psi|^{p-1}\right)\partial_j\psi\right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\
&\quad + \left\||\psi_n|^{p-3}\psi_n^2\left(\partial_j\overline{\psi_n} - \partial_j\overline{\psi}\right)\right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\
&\quad + \left\|\left(|\psi_n|^{p-3}\psi_n^2 - |\psi|^{p-3}\psi^2\right)\partial_j\overline{\psi}\right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\
&\lesssim \|\partial_j\phi_{+,n} - \partial_j\phi_+\|_{L^2} + \|\psi_n\|_{X([T_\delta,\infty)}^{p-1}\|\partial_j\psi_n - \partial_j\psi\|_{S([T_\delta,\infty))} \\
&\quad + \left\||\psi_n|^{p-1} - |\psi|^{p-1}\right\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})}\|\partial_j\psi\|_{S([T_\delta,\infty))} \\
&\quad + \|\psi_n\|_{X([T_\delta,\infty))}^{p-1}\|\partial_j\psi_n - \partial_j\psi\|_{S([T_\delta,\infty))} \\
&\quad + \left\||\psi_n|^{p-3}\psi_n^2 - |\psi|^{p-3}\psi^2\right\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})}\|\partial_j\psi\|_{S([T_\delta,\infty))},
\end{aligned} \tag{3.132}$$

where the implicit constant depends only on d, p and q_1. Note here that, Lemma 3.6, together with (3.119), gives us that $\|(1-\Delta)^{\frac{1}{2}}\psi\|_{S([0,\infty))} < \infty$. We also note that (3.131) implies that

$$\begin{aligned}&\lim_{n\to\infty} \big\| |\psi_n|^{p-1} - |\psi|^{p-1} \big\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})} \\ &= \lim_{n\to\infty} \big\| |\psi_n|^{p-3}\psi_n^2 - |\psi|^{p-3}\psi^2 \big\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})} = 0 \quad \text{for all sufficiently small } \delta > 0. \end{aligned} \tag{3.133}$$

Hence, we have by (3.132) that

$$\lim_{n\to\infty} \|\partial_j\psi_n - \partial_j\psi\|_{S([T_\delta,\infty))} = 0 \quad \text{for all sufficiently small } \delta > 0. \tag{3.134}$$

Thus, we obtain from (3.130) and (3.134) that

$$\lim_{n\to\infty} \|\psi_n - \psi\|_{L^\infty([T_\delta,\infty);H^1)} = 0 \quad \text{for all sufficiently small } \delta > 0, \tag{3.135}$$

so that it follows from the continuous dependence of solutions on initial data that

$$\lim_{n\to\infty} \|W_+\phi_{+,n} - W_+\phi_+\|_{H^1} = \lim_{n\to\infty} \|\psi_n(0) - \psi(0)\|_{H^1} = 0, \tag{3.136}$$

which completes the proof. ☐

4 Analysis on PW_+

Our aim here is to prove Theorem 1.1. Obviously, Proposition 2.5 provides (1.48) and (1.49). Therefore, it remains to prove the asymptotic completeness in PW_+.

In order to prove the existence of asymptotic states for a solution ψ, it suffices to show that $\|\psi\|_{X(\mathbb{R})} < \infty$ by virtue of Proposition 3.5 and Proposition 2.5. To this end, we introduce a subset of PW_+:

$$PW_+(\delta) := \left\{ f \in PW_+ \;\middle|\; \widetilde{\mathcal{N}}_2(f) := \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}(f)}^{\frac{d}{2}(p-1)-2} < \delta \right\}, \quad \delta > 0. \tag{4.1}$$

Moreover, we define a number N_c by

$$\begin{aligned}\widetilde{N}_c &:= \sup\big\{ \delta > 0 \bigm| \|\psi\|_{X(\mathbb{R})} < \infty \quad \text{for all } \psi_0 \in PW_+(\delta) \big\} \\ &= \inf\big\{ \delta > 0 \bigm| \|\psi\|_{X(\mathbb{R})} = \infty \quad \text{for some } \psi_0 \in PW_+(\delta) \big\},\end{aligned} \tag{4.2}$$

where ψ denotes the solution to (1.1) with $\psi(0) = \psi_0$.

Since we have

$$PW_+ = PW_+(\widetilde{N}_2), \tag{4.3}$$

our task is to prove $\widetilde{N}_c = \widetilde{N}_2$.

It is worth while noting here that $\widetilde{N}_c > 0$: For, it follows from (3.8), the interpolation estimate and Proposition 2.5 that

$$\begin{aligned}
\left\| e^{\frac{i}{2}t\Delta}\psi_0 \right\|_{X(\mathbb{R})}^{p-1} &\lesssim \|(-\Delta)^{\frac{s_p}{2}}\psi_0\|_{L^2}^{p-1} \\
&\le \|\psi_0\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \|\nabla\psi_0\|_{L^2}^{\frac{d}{2}(p-1)-2} \\
&< \|\psi_0\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\frac{d(p-1)}{d(p-1)-4}\mathcal{H}(\psi_0)}^{\,\frac{d}{2}(p-1)-2} \\
&= \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\,\frac{d}{2}(p-1)-2} \widetilde{\mathcal{N}}_2(\psi_0) \\
&< \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\,\frac{d}{2}(p-1)-2} \delta \qquad \text{for all } \delta > 0 \text{ and } \psi_0 \in PW_+(\delta),
\end{aligned} \tag{4.4}$$

where the implicit constant depends only on d, p and q_1. This estimate shows that there exists $\delta_0 > 0$, depending only on d, p and q_1, such that

$$\left\| e^{\frac{i}{2}t\Delta}\psi_0 \right\|_{X(\mathbb{R})} < \delta_{Pr.3.7} \quad \text{for all } \psi_0 \in PW_+(\delta_0), \tag{4.5}$$

where $\delta_{Pr.3.7}$ is a constant found in Proposition 3.7. Hence, we have by the small data theory (Proposition 3.7) that $\widetilde{N}_2 \ge \delta_0 > 0$.

4.1 One soliton vs. Virial identity

In this section, we present our strategy to prove $\widetilde{N}_c = \widetilde{N}_2$. We suppose the contrary that $\widetilde{N}_c < \widetilde{N}_2$. In this undesired situation, we can find a one-soliton-like solution to our equation (1.1) in PW_+ (see Proposition 4.1 below). Then, the soliton-like behavior contradicts the one described by the generalized virial identity (Lemma A.3), so that we conclude that $\widetilde{N}_c = \widetilde{N}_2$. At the end of this Section 4.1, we actually show this, provided that such a soliton-like solution exists.

The construction of a soliton-like solution is rather long. We divide it into two parts; In Section 4.2, one finds a candidate for the soliton, and in Section 4.3, one sees that the candidate actually behaves like a one-soliton-like solution.

We here briefly explain how to find a candidate for the one-soliton-like solution. If $\widetilde{N}_c < \widetilde{N}_2$, then we can take a sequence $\{\psi_n\}$ of solutions to (1.1) with the property that

$$\psi_n(t) \in PW_+ \text{ for all } t \in \mathbb{R}, \quad \|\psi_n\|_{X(\mathbb{R})} = \infty, \quad \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(\psi_n(0)) = \widetilde{N}_c. \tag{4.6}$$

We consider the integral equation for ψ_n:

$$\psi_n(t) = e^{\frac{i}{2}t\Delta} f_n + \frac{i}{2}\int_0^t e^{\frac{i}{2}(t-t')\Delta} \left\{ |\psi_n(t')|^{p-1}\psi_n(t') \right\} dt', \tag{4.7}$$

where we put $f_n = \psi_n(0)$. We first observe that the linear part of this integral equation possibly behaves like as follows[1]:

$$e^{\frac{i}{2}t\Delta} f_n(x) \sim \sum_{l\ge 1} e^{\frac{i}{2}(t-\tau_n^l)\Delta} e^{-\eta_n^l\cdot\nabla} f^l(x) \tag{4.8}$$

---

[1] $e^{-\eta_n^l\cdot\nabla}$ denotes the space-translation by $-\eta_n$. We may expect the number of summands f^l is finite.

for some nontrivial functions $f^l \in PW_+$, $\tau_n^l \in \mathbb{R}$ and $\eta_n^l \in \mathbb{R}^d$. Of course, this is not a good approximation to ψ_n. So, putting $\tau_\infty^l = \lim_{n\to\infty} \tau_n^l$ (possibly $\tau_\infty^l = \pm\infty$), we solve our equation (1.1) with the initial datum $e^{-\frac{i}{2}\tau_\infty^l\Delta} f^l$ at $t = -\tau_\infty^l$:

$$\psi^l(t) = e^{\frac{i}{2}(t+\tau_\infty^l)\Delta} e^{-\frac{i}{2}\tau_\infty^l\Delta} f^l + \frac{i}{2}\int_{-\tau_\infty^l}^{t} e^{\frac{i}{2}(t-t')\Delta} \left\{ |\psi^l(t')|^{p-1}\psi^l(t') \right\} dt'. \tag{4.9}$$

Here, in case of $\tau_\infty^l = \pm\infty$, we are regarding this as the final value problem:

$$e^{-\frac{i}{2}t\Delta}\psi^l(t) = f^l + \frac{i}{2}\int_{\mp\infty}^{t} e^{-\frac{i}{2}t'\Delta} \left\{ |\psi^l(t')|^{p-1}\psi^l(t') \right\} dt'. \tag{4.10}$$

Then, instead of (4.8), we consider the superposition of these solutions with the space-time translations:

$$\psi_n^{app}(x,t) := \sum_{l\geq 1} (e^{-\tau_n^l \frac{\partial}{\partial t} - \eta_n^l\cdot\nabla}\psi^l)(x,t) = \sum_{l\geq 1} \psi^l(x-\eta_n^l, t-\tau_n^l). \tag{4.11}$$

By Lemma 4.5 below, we will see that this formal object ψ_n^{app} is an "almost" solution to our equation (1.1) with the initial datum $\sum_{l\geq 1} e^{-\frac{i}{2}\tau_n^l\Delta} e^{-\eta_n^l\cdot\nabla} f^l$, and supposed to be a good approximation to ψ_n. In other words, a kind of superposition principle holds valid in an asymptotic sense as $n \to \infty$. By virtue of the long time perturbation theory (Proposition 3.8), the sum in ψ_n^{app} consists of a finite number of solutions. Actually, as a consequence of the minimizing property of the sequence $\{\psi_n\}$ (4.6), the summand is just one: put $\Psi := \psi^1$. Then, it turns out that Ψ is a one-soliton-like solution which we are looking for. In fact, we can prove:

Proposition 4.1 (One-soliton-like solution in PW_+)**.** *Suppose that* $\widetilde{N}_c < \widetilde{N}_2$. *Then, there exists a global solution* $\Psi \in C(\mathbb{R}; H^1(\mathbb{R}^d))$ *to the equation (1.1) with the following properties:* $\{\Psi(t)\}_{t\in\mathbb{R}}$ *is;*
(i) *a minimizer such that*

$$\|\Psi\|_{X(\mathbb{R})} = \infty, \qquad \widetilde{\mathcal{N}}_2(\Psi(t)) = \widetilde{N}_c, \quad \Psi(t) \in PW_+ \quad \textit{for all } t \in \mathbb{R}, \tag{4.12}$$

(ii) *uniformly bounded in* $H^1(\mathbb{R}^d)$ *with*

$$\|\Psi(t)\|_{L^2} = \|\Psi(0)\|_{L^2} = 1 \quad \textit{for all } t \in \mathbb{R}, \tag{4.13}$$

and

$$\sup_{t\in\mathbb{R}} \|\nabla\Psi(t)\|_{L^2} \leq N_c^{\frac{1}{\frac{d}{2}(p-1)-2}}, \tag{4.14}$$

(iii) *a family of functions with zero momentum, i.e.,*

$$\Im \int_{\mathbb{R}^d} \overline{\Psi}(x,t)\nabla\Psi(x,t)\, dx = 0 \quad \textit{for all } t \in \mathbb{R}, \tag{4.15}$$

(iv) *tight in* $H^1(\mathbb{R}^d)$ *in the following sense: For any* $\varepsilon > 0$, *there exists* $R_\varepsilon > 0$ *and a continuous path* $\gamma_\varepsilon \in C([0,\infty); \mathbb{R}^d)$ *with* $\gamma_\varepsilon(0) = 0$ *such that*

$$\int_{|x-\gamma_\varepsilon(t)|<R_\varepsilon} |\Psi(x,t)|^2\, dx > 1 - \varepsilon \quad \textit{for all } t \in [0,\infty), \tag{4.16}$$

and

$$\int_{|x-\gamma_\varepsilon(t)|<R_\varepsilon} |\nabla\Psi(x,t)|^2\, dx > \|\nabla\Psi(t)\|_{L^2}^2 - \varepsilon \quad \textit{for all } t \in [0,\infty). \tag{4.17}$$

We will give the proof of Proposition 4.1 in Sections 4.2 and 4.3 later, as mentioned before. The properties (i), (ii) and (iii) are easy matters. The most important part is the tightness (iv). To prove the tightness, we introduce a concentrate function for Ψ (see (4.214) below) and consider a sequence $\{\Psi(\cdot+t_n)\}_{n\in\mathbb{N}}$ for an appropriate sequence $\{t_n\}$ with $t_n\to+\infty$. Analogous arguments with finding Ψ work on $\{\Psi(\cdot+t_n)\}_{n\in\mathbb{N}}$ as well, so that we can show the tightness. Once we get the tightness, Lemma E.2 immediately gives us the continuous path γ_ε described in (iv) of Proposition 4.1. In order to prove $\widetilde{N}_c=\widetilde{N}_2$, we need to know more subtle behavior of the path however. For a sufficiently long time, we can take γ_ε as the almost center of mass, say γ_ε^{ac}:

Lemma 4.2 (Almost center of mass)**.** *Let Ψ be a global solution to the equation (1.1) satisfying the properties (4.12), (4.13), (4.14), (4.15), and (4.16), (4.17). Let R_ε be a radius found in* (iv) *of Proposition 4.1 for $\varepsilon>0$. We define an "almost center of mass" by*

$$\gamma_{\varepsilon,R}^{ac}(t):=(\vec{w}_{20R},|\Psi(t)|^2)\quad \text{for all } \varepsilon\in(0,\tfrac{1}{100}) \text{ and } R>R_\varepsilon, \tag{4.18}$$

where $\vec{w}_R$ is the function defined by (A.3). Then, we have:

$$\gamma_{\varepsilon,R}^{ac}\in C^1([0,\infty);\mathbb{R}^d), \tag{4.19}$$

and there exists a constant $\alpha>0$, depending only on d and p, such that

$$\left|\gamma_{\varepsilon,R}^{ac}(t)\right|\le 20R\quad \text{for all } t\in\left[0,\alpha\tfrac{R}{\sqrt{\varepsilon}}\right], \tag{4.20}$$

$$\int_{|x-\gamma_{\varepsilon,R}^{ac}(t)|\le 4R}|\Psi(x,t)|^2+|\nabla\Psi(x,t)|^2\,dx\ge\|\Psi(t)\|_{H^1}^2-\varepsilon\quad \text{for all } t\in\left[0,\alpha\tfrac{R}{\sqrt{\varepsilon}}\right]. \tag{4.21}$$

Remark 4.1. *In the proof below, we find that the following estimate holds (see (4.33)):*

$$\left|\frac{d\gamma_{\varepsilon,R}^{ac}}{dt}(t)\right|\lesssim\sqrt{\varepsilon}\quad \text{for all } t\in\left[0,\alpha\tfrac{R}{\sqrt{\varepsilon}}\right],$$

where the implicit constant depends only on d and q.

Proof of Lemma 4.2. We have from Proposition B.1 in [46] that

$$\gamma_{\varepsilon,R}^{ac}(t)=(\vec{w}_{20R},|\Psi(0)|^2)+\left(2\Im\int_0^t\int_{\mathbb{R}^d}\nabla\vec{w}_{20R}^{\,j}(x)\cdot\nabla\Psi(x,s)\overline{\Psi(x,s)}\,dx\,ds\right)_{j=1,\dots,d}. \tag{4.22}$$

This formula, with the help of (4.13), (4.14) and (A.8), immediately shows (4.19): $\gamma_{\varepsilon,R}^{ac}\in C^1([0,\infty);\mathbb{R}^d)$.

Next, we shall prove the properties (4.20) and (4.21). Let γ_ε be a path found in Proposition 4.1, and t_ε be the first time such that the size of γ_ε reaches $10R$, i.e.,

$$t_\varepsilon:=\inf\left\{t\ge0\;\middle|\;|\gamma_\varepsilon(t)|=10R\right\}. \tag{4.23}$$

Since $\gamma_\varepsilon\in C([0,\infty);\mathbb{R}^d)$ with $\gamma_\varepsilon(0)=0$, we have that $t_\varepsilon>0$ and

$$|\gamma_\varepsilon(t)|\le 10R\quad \text{for all } t\in[0,t_\varepsilon]. \tag{4.24}$$

We claim that

$$|\gamma_{\varepsilon,R}^{ac}(t)-\gamma_\varepsilon(t)|<2R\quad \text{for all } t\in[0,t_\varepsilon]. \tag{4.25}$$

It follows from the property (4.13) that

$$\begin{aligned}
\left|\gamma_{\varepsilon,R}^{ac}(t)-\gamma_\varepsilon(t)\right| &= \left|(\vec{w}_{20R},|\Psi(t)|^2)-\gamma_\varepsilon(t)\|\Psi(t)\|_{L^2}^2\right| \\
&= \left|\int_{\mathbb{R}^d}\{\vec{w}_{20R}(x)-\gamma_\varepsilon(t)\}\,|\Psi(x,t)|^2\,dx\right| \\
&\le \int_{|x-\gamma_\varepsilon(t)|\le R}|x-\gamma_\varepsilon(t)|\,|\Psi(x,t)|^2\,dx \\
&\qquad + \int_{|x-\gamma_\varepsilon(t)|\ge R}|\vec{w}_{20R}-\gamma_\varepsilon(t)|\,|\Psi(x,t)|^2\,dx.
\end{aligned} \tag{4.26}$$

Moreover, applying (4.24) and (A.6) to the second term on the right-hand side above, we obtain that

$$\left|\gamma_{\varepsilon,R}^{ac}(t)-\gamma_\varepsilon(t)\right| \le R\|\Psi(t)\|_{L^2}^2+50R\int_{|x-\gamma_\varepsilon(t)|\ge R}|\Psi(x,t)|^2\,dx \quad \text{for all } t\in[0,t_\varepsilon]. \tag{4.27}$$

Hence, this inequality (4.27), together with (4.13) and the tightness (4.16), yields that

$$\left|\gamma_{\varepsilon,R}^{ac}(t)-\gamma_\varepsilon(t)\right| \le R+50R\varepsilon<2R \quad \text{for all } \varepsilon<\tfrac{1}{100} \text{ and } t\in[0,t_\varepsilon]. \tag{4.28}$$

Now, we have by (4.24) and (4.25) that

$$\left|\gamma_{\varepsilon,R}^{ac}(t)\right| \le 12R \quad \text{for all } t\in[0,t_\varepsilon]. \tag{4.29}$$

Moreover, (4.25) also gives us that

$$B_R(\gamma_\varepsilon(t))\subset B_{4R}(\gamma_\varepsilon^{ac}(t)) \quad \text{for all } t\in[0,t_\varepsilon], \tag{4.30}$$

so that the tightness of $\{\Psi(t)\}$ in $H^1(\mathbb{R}^d)$ (see (4.16) and (4.17)) gives us that

$$\int_{|x-\gamma_{\varepsilon,R}^{ac}(t)|\le 4R}|\Psi(x,t)|^2+|\nabla\Psi(t)|^2\,dx\ge\|\Psi(t)\|_{H^1}^2-\varepsilon \quad \text{for all } t\in[0,t_\varepsilon]. \tag{4.31}$$

Therefore, for the desired results (4.20) and (4.21), it suffices to show that there exists a constant $\alpha>0$, depending only on d and p, such that

$$\alpha\frac{R}{\sqrt{\varepsilon}}\le t_\varepsilon. \tag{4.32}$$

To this end, we prove that

$$\left|\frac{d\gamma_{\varepsilon,R}^{ac}}{dt}(t)\right|\lesssim\sqrt{\varepsilon} \quad \text{for all } t\in[0,t_\varepsilon], \tag{4.33}$$

where the implicit constant depends only on d and p.

Before proving (4.33), we describe how it yields (4.32). We easily verify by (4.33) that

$$|\gamma_{\varepsilon,R}^{ac}(t_\varepsilon)|-|\gamma_{\varepsilon,R}^{ac}(0)|\le\int_0^{t_\varepsilon}\left|\frac{d\gamma_{\varepsilon,R}^{ac}}{dt}(t)\right|dt\lesssim\sqrt{\varepsilon}t_\varepsilon, \tag{4.34}$$

where the implicit constant depends only on d and p. Then, it follows from (4.25) and $\gamma_\varepsilon(0) = 0$ that

$$\begin{aligned}\sqrt{\varepsilon} t_\varepsilon &\gtrsim |\gamma_{\varepsilon,R}^{ac}(t_\varepsilon)| - 2R \\ &\geq |\gamma_\varepsilon(t_\varepsilon)| - |\gamma_{\varepsilon,R}^{ac}(t_\varepsilon) - \gamma_\varepsilon(t_\varepsilon)| - 2R \\ &\geq 10R - 2R - 2R = 6R,\end{aligned} \tag{4.35}$$

which gives (4.32).

Finally, we prove (4.33). Using the formula (4.22) and the property (4.15), we obtain that

$$\begin{aligned}\left|\frac{d\gamma_{\varepsilon,R}^{ac}}{dt}(t)\right|^2 &= \sum_{j=1}^{d} \left|2\Im(\nabla \vec{w}_{20R}^{j} \cdot \nabla\Psi(t), \Psi(t))\right|^2 \\ &\leq 4\sum_{j=1}^{d} \left\|\nabla \vec{w}_{20R}^{j}\right\|_{L^\infty}^2 \|\Psi(t)\|_{L^2}^2 \int_{|x|\geq 20R} |\nabla\Psi(x,t)|^2 \, dx \quad \text{for all } t \geq 0.\end{aligned} \tag{4.36}$$

Applying (4.13) and (A.8) to the right-hand side above, we further obtain that

$$\left|\frac{d\gamma_{\varepsilon,R}^{ac}}{dt}(t)\right|^2 \lesssim \int_{|x|\geq 20R} |\nabla\Psi(x,t)|^2 \, dx \quad \text{for all } t \geq 0, \tag{4.37}$$

where the implicit constant depends only on d and p. Since the estimate (4.24) shows that

$$B_R(\gamma_\varepsilon(t)) \subset B_{20R}(0) \quad \text{for all } t \in [0, t_\varepsilon], \tag{4.38}$$

the estimate (4.37), together with the tightness (4.17), leads to (4.33). ☐

Lemma 4.2 implies that Ψ found in Proposition 4.1 is in a bound motion, rather, a standing wave. On the other hand, the generalized virial identity ((A.20) in Lemma A.3) suggests that Ψ is in a scattering motion. As we already mentioned at the beginning of this Section 4.1, these two facts contradict each other; Thus, we see that $\widetilde{N}_2 = \widetilde{N}_c$. Here, we show this point precisely:

The generalized virial identity (A.20), together with (A.27) and (A.28), yields that

$$\begin{aligned}&(W_R, |\Psi(t)|^2) \\ &\geq (W_R, |\Psi(0)|^2) + 2t\Im(\vec{w}_R \cdot \nabla\Psi(0), \Psi(0)) + 2\int_0^t \int_0^{t'} \mathcal{K}(\Psi(t''))\, dt''dt' \\ &\quad - 2\int_0^t \int_0^{t'} \int_{|x|\geq R} \rho_1(x)|\nabla\Psi(x,t'')|^2 + \rho_2(x)\left|\frac{x}{|x|}\cdot\nabla\Psi(x,t'')\right|^2 dx dt'' dt' \\ &\quad - \frac{1}{2}\int_0^t \int_0^{t'} \|\Delta(\operatorname{div}\vec{w}_R)\|_{L^\infty} \left\|\Psi(t'')\right\|_{L^2}^2 dt''dt' \qquad \text{for all } R > 0.\end{aligned} \tag{4.39}$$

Applying the estimates (4.13), (A.9) and (A.31) to the right-hand side above, we obtain that

$$\begin{aligned}&(W_R, |\Psi(t)|^2)\\&\ge (W_R, |\Psi(0)|^2) + 2t\Im(\vec{w}_R\cdot\nabla\Psi(0), \Psi(0)) + 2\int_0^t\int_0^{t'}\mathcal{K}(\Psi(t''))\,dt''dt'\\&\quad - 2\,(K_1+K_2)\int_0^t\int_0^{t'}\int_{|x|\ge R}|\nabla\Psi(x,t'')|^2\,dxdt''dt' - \frac{10d^2K}{R^2}t^2 \quad \text{for all } R>0,\end{aligned} \tag{4.40}$$

where K is the constant defined in (A.1), and K_1 and K_2 are the constants found in Lemma A.4. Moreover, it follows from the estimate (2.27) in Proposition 2.5 that

$$\begin{aligned}&(W_R, |\Psi(t)|^2)\\&\ge (W_R, |\Psi(0)|^2) + 2t\Im(\vec{w}_R\cdot\nabla\Psi(0), \Psi(0)) + t^2\omega_0\mathcal{H}(\Psi(0))\\&\quad - 2(K_1+K_2)\int_0^t\int_0^{t'}\int_{|x|\ge R}|\nabla\Psi(x,t'')|^2\,dxdt''dt' - \frac{10d^2K}{R^2}t^2 \quad \text{for all } R>0,\end{aligned} \tag{4.41}$$

where we put

$$\omega_0 := 1 - \frac{\widetilde{N}_2(\Psi(0))}{\widetilde{N}_2}. \tag{4.42}$$

Here, we have by Lemma 4.2 that: For any $\varepsilon \in (0, \frac{1}{100})$, there exists $R_\varepsilon > 0$ with the following property: for any $R \ge R_\varepsilon$, there exists $\gamma_{\varepsilon,R}^{ac} \in C^1([0,\infty);\mathbb{R}^d)$ such that

$$\left|\gamma_{\varepsilon,R}^{ac}(t)\right| \le 20R \quad \text{for all } t \in \big[\,0,\ \alpha\tfrac{R}{\sqrt{\varepsilon}}\,\big], \tag{4.43}$$

$$\int_{|x-\gamma_{\varepsilon,R}^{ac}(t)|\ge 4R}|\nabla\Psi(x,t)|^2dx < \varepsilon \quad \text{for all } t \in \big[\,0,\ \alpha\tfrac{R}{\sqrt{\varepsilon}}\,\big], \tag{4.44}$$

where α is some constant depending only on d and p.

We employ (4.43) to obtain that

$$|x-\gamma_{\varepsilon,R}^{ac}(t)| \ge 4R \quad \text{for all } R \ge R_\varepsilon,\ t \in \big[\,0,\ \alpha\tfrac{R}{\sqrt{\varepsilon}}\,\big] \text{ and } x \in \mathbb{R}^d \text{ with } |x| \ge 24R. \tag{4.45}$$

Hence, (4.41), together with the tightness (4.44), leads to that

$$\begin{aligned}(W_{50R}, |\Psi(t)|^2) &\ge (W_{50R}, |\Psi(0)|^2) + 2t\Im(\vec{w}_{50R}\cdot\nabla\Psi(0), \Psi(0)) + t^2\omega_0\mathcal{H}(\Psi(0))\\&\quad - (K_1+K_2)t^2\varepsilon - \frac{10d^2K}{(50R)^2}t^2 \quad \text{for all } R \ge R_\varepsilon \text{ and } t \in \big[\,0,\ \alpha\tfrac{R}{\sqrt{\varepsilon}}\,\big]\,.\end{aligned} \tag{4.46}$$

We choose ε so small that

$$0 < \varepsilon < \min\left\{\frac{1}{100},\ \frac{\omega_0}{4(K_1+K_2)}\mathcal{H}(\Psi(0))\right\}, \tag{4.47}$$

and R so large that

$$R \ge \max\left\{R_\varepsilon,\ \frac{d\sqrt{K}}{\sqrt{\omega_0\mathcal{H}(\Psi(0))}}\right\}. \tag{4.48}$$

Then, it follows from (4.46) that

$$
\begin{aligned}
(W_{50R}, |\Psi(t)|^2) \ge (W_{50R}, |\Psi(0)|^2) + 2t\Im(\vec{w}_{50R}\cdot\nabla\Psi(0), \Psi(0)) + \frac{t^2}{2}\omega_0\mathcal{H}(\Psi(0))& \\
\text{for all } t \in \left[\, 0,\, \alpha\tfrac{R}{\sqrt{\varepsilon}}\, \right].&
\end{aligned}
\tag{4.49}
$$

Dividing the both sides of (4.49) by t^2 and applying the estimates (4.13) and (A.7), we obtain that

$$
\begin{aligned}
\frac{8(50R)^2}{t^2} \ge \frac{1}{t^2}(W_{50R}, |\Psi(0)|^2) + \frac{2}{t}\Im(\vec{w}_{50R}\cdot\nabla\Psi(0), \Psi(0)) + \frac{\omega_0}{2}\mathcal{H}(\Psi(0))& \\
\text{for all } t \in \left[\, 0,\, \alpha\tfrac{R}{\sqrt{\varepsilon}}\, \right].&
\end{aligned}
\tag{4.50}
$$

In particular, when $t = \alpha\frac{R}{\sqrt{\varepsilon}}$, we have by (4.13), (A.6) and (A.7) that

$$
\begin{aligned}
\frac{8(50)^2\varepsilon}{\alpha^2} &\ge \frac{\varepsilon}{\alpha^2 R^2}(W_{50R}, |\Psi(0)|^2) + \frac{2\sqrt{\varepsilon}}{\alpha R}\Im(\vec{w}_{50R}\cdot\nabla\Psi(0), \Psi(0)) + \frac{\omega_0}{2}\mathcal{H}(\Psi(0)) \\
&\ge -\frac{8(50)^2\varepsilon}{\alpha^2} - \frac{200\sqrt{\varepsilon}}{\alpha}\|\nabla\Psi(0)\|_{L^2} + \frac{\omega_0}{2}\mathcal{H}(\Psi(0)),
\end{aligned}
\tag{4.51}
$$

so that

$$
\frac{8(50)^2\varepsilon}{\alpha^2} + \frac{8(50)^2\varepsilon}{\alpha^2} + \frac{200\sqrt{\varepsilon}}{\alpha}\|\nabla\Psi(0)\|_{L^2} \ge \frac{\omega_0}{2}\mathcal{H}(\Psi(0)). \tag{4.52}
$$

However, taking $\varepsilon \to 0$ in (4.52), we obtain a contradiction. This absurd conclusion comes from the existence of one-soliton-like solution Ψ (see Proposition 4.1). Thus, it must hold that $\widetilde{N}_c = \widetilde{N}_2$, provided that Proposition 4.2 is valid.

4.2 Solving the variational problem for $\widetilde{N}_c$

In this section, we construct a candidate for the one-soliton-like solution, considering the variational problem for $\widetilde{N}_c$.

Supposing that $\widetilde{N}_c < \widetilde{N}_2$, we can take a minimizing sequence $\{\delta_n\}_{n\in\mathbb{N}}$ such that

$$
\widetilde{N}_c < \delta_n < \widetilde{N}_2 \quad \text{for all } n \in \mathbb{N}, \qquad \lim_{n\to\infty}\delta_n = \widetilde{N}_c. \tag{4.53}
$$

Moreover, Lemma 2.3 enables us to take a sequence $\{\psi_{0,n}\}_{n\in\mathbb{N}}$ in PW_+ such that

$$
\|\psi_{0,n}\|_{L^2} = 1 \quad \text{for all } n \in \mathbb{N}, \tag{4.54}
$$

$$
\widetilde{N}_c < \widetilde{\mathcal{N}}_2(\psi_{0,n}) < \delta_n \quad \text{for all } n \in \mathbb{N}. \tag{4.55}
$$

Note that (4.55), together with (4.53), leads to that

$$
\lim_{n\to\infty}\widetilde{\mathcal{N}}_2(\psi_{0,n}) = \widetilde{N}_c. \tag{4.56}
$$

Let ψ_n be the solution to (1.1) with $\psi_n(0) = \psi_{0,n}$. Then, (4.55), together with the definition of $\widetilde{N}_c$ (see (4.2)), implies that

$$
\|\psi_n\|_{X(\mathbb{R})} = \infty \quad \text{for all } n \in \mathbb{N}. \tag{4.57}
$$

We also find that

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\psi_{0,n} \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} > 0. \tag{4.58}$$

Indeed, if (4.58) fails, then the small data theory (Proposition 3.7) concludes that

$$\|\psi_n\|_{X(\mathbb{R})} < \infty \quad \text{for sufficiently large } n \in \mathbb{N},$$

which contradicts (4.57).

The following lemma gives us a candidate for the one-soliton-like solution in Proposition 4.1.

Lemma 4.3. *Assume that* $d \ge 1$ *and* $2 + \frac{4}{d} < p+1 < 2^*$. *Suppose that* $\widetilde{N}_c < \widetilde{N}_2$. *Let* $\{\psi_n\}$ *be a sequence of global solutions to the equation (1.1) in* $C(\mathbb{R}; H^1(\mathbb{R}^d))$ *such that*

$$\psi_n(t) \in PW_+ \quad \text{for all } t \in \mathbb{R} \text{ and } n \in \mathbb{N}, \tag{4.59}$$

$$\|\psi_n(t)\|_{L^2} = 1 \quad \text{for all } n \in \mathbb{N} \text{ and } t \in \mathbb{R}, \tag{4.60}$$

$$\sup_{n\in\mathbb{N}} \|\psi_n(0)\|_{H^1} < \infty, \tag{4.61}$$

$$\lim_{n\to\infty} \widetilde{\mathcal{N}}_2(\psi_n(t)) = \widetilde{N}_c \quad \text{for all } t \in \mathbb{R}, \tag{4.62}$$

$$\|\psi_n\|_{X(\mathbb{R})} = \infty \quad \text{for all } n \in \mathbb{N}. \tag{4.63}$$

Furthermore, we suppose that

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\psi_n(0) \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} > 0. \tag{4.64}$$

Then, there exists a subsequence of $\{\psi_n\}$ *(still denoted by the same symbol), which satisfies the following property: There exist*
(i) *a nontrivial global solution* $\Psi \in C(\mathbb{R}; H^1(\mathbb{R}^d))$ *to the equation (1.1) with*

$$\|\Psi\|_{X(\mathbb{R})} = \infty, \tag{4.65}$$

$$\Psi(t) \in PW_+ \quad \text{for all } t \in \mathbb{R}, \tag{4.66}$$

$$\|\Psi(t)\|_{L^2} = 1 \quad \text{for all } t \in \mathbb{R}, \tag{4.67}$$

$$\widetilde{\mathcal{N}}_2(\Psi(t)) = \widetilde{N}_c \quad \text{for all } t \in \mathbb{R}, \tag{4.68}$$

and
(ii) *a nontrivial function* $f \in PW_+$, *a sequence* $\{\tau_n\}$ *in* $\mathbb{R}$ *with* $\lim_{n\to\infty} \tau_n = \tau_\infty$ *for some* $\tau_\infty \in \mathbb{R} \cup \{\pm\infty\}$, *and a sequence* $\{\eta_n\}$ *in* $\mathbb{R}^d$ *such that*

$$\lim_{n\to\infty} e^{\frac{i}{2}\tau_n\Delta}e^{\eta_n\cdot\nabla}\psi_n(0) = f \quad \text{weakly in } H^1(\mathbb{R}^d), \text{ and a.e. in } \mathbb{R}^d, \tag{4.69}$$

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}\tau_n\Delta}e^{\eta_n\cdot\nabla}\psi_n(0) - f \right\|_{L^2(\mathbb{R}^d)} = 0, \tag{4.70}$$

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}(t+\tau_n)\Delta}e^{\eta_n\cdot\nabla}\psi_n(0) - e^{\frac{i}{2}t\Delta}f \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})\cap X(\mathbb{R})} = 0, \tag{4.71}$$

$$\lim_{n\to\infty} \left\| \Psi(-\tau_n) - e^{-\frac{i}{2}\tau_n\Delta}f \right\|_{H^1} = 0, \tag{4.72}$$

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\psi_n(0) - e^{\frac{i}{2}t\Delta}\left(e^{-\tau_n\frac{\partial}{\partial t} - \eta_n\cdot\nabla}\Psi\right)(0) \right\|_{X(\mathbb{R})} = 0. \tag{4.73}$$

Especially, we have

$$\|f\|_{L^2} = \|\Psi(t)\|_{L^2} \quad \text{for all } t \in \mathbb{R}, \qquad \|\nabla f\|_{L^2} = \lim_{n\to\infty} \|\nabla\Psi(-\tau_n)\|_{L^2}. \tag{4.74}$$

Remark 4.2. *In this Lemma 4.3, Ψ is actually a solution to the integral equation (4.9).*

For the proof of Lemma 4.3, we prepare the following Lemmata 4.4 and 4.5. The former is a compactness lemma and the latter is a key ingredient to prove the superposition principle (4.11) as mentioned in Section 4.1.

Lemma 4.4. *Assume that $d \ge 1$ and $2 + \frac{4}{d} < p + 1 < 2^*$. Let $\{f_n\}$ be a uniformly bounded sequence in $H^1(\mathbb{R}^d)$. Suppose that*

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta} f_n \right\|_{L^\infty(\mathbb{R}; L^{\frac{d}{2}(p-1)})} > 0. \tag{4.75}$$

Then, there exists a subsequence of $\{f_n\}$ (still denoted by the same symbol), which satisfies the following property: There exist a nontrivial function $f \in H^1(\mathbb{R}^d)$, a sequence $\{t_n\}$ in $\mathbb{R}$ with $\lim_{n\to\infty} t_n = t_\infty$ for some $t_\infty \in \mathbb{R} \cup \{\pm\infty\}$, and a sequence $\{y_n\}$ in $\mathbb{R}^d$ such that, putting $f_n^*(x) := e^{\frac{i}{2}t_n\Delta} f_n(x + y_n)$, we have that

$$\lim_{n\to\infty} f_n^* = f \quad \text{weakly in } H^1(\mathbb{R}^d), \text{ and strongly in } L^q_{loc}(\mathbb{R}^d) \text{ for all } q \in [2, 2^*), \tag{4.76}$$

$$\lim_{n\to\infty} \left\{ \||\nabla|^s f_n\|_{L^2}^2 - \||\nabla|^s (f_n^* - f)\|_{L^2}^2 \right\} = \||\nabla|^s f\|_{L^2}^2 \quad \text{for all } s \in [0, 1], \tag{4.77}$$

$$\lim_{n\to\infty} \left\{ \|f_n\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f) \right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f \right\|_{L^q}^q \right\} = 0 \quad \text{for all } q \in [2, 2^*), \tag{4.78}$$

$$\lim_{n\to\infty} \left\{ \mathcal{H}(f_n) - \mathcal{H}\left(e^{-\frac{i}{2}t_n\Delta}(f_n^* - f) \right) - \mathcal{H}\left(e^{-\frac{i}{2}t_n\Delta} f \right) \right\} = 0, \tag{4.79}$$

$$\sup_{n\in\mathbb{N}} \|f_n^* - f\|_{H^1} \le \sup_{n\in\mathbb{N}} \|f_n\|_{H^1} < \infty. \tag{4.80}$$

Proof of Lemma 4.4. Put

$$A = \frac{1}{2} \limsup_{n\to\infty} \left\| e^{\frac{i}{2}t_n\Delta} f_n \right\|_{L^{\frac{d}{2}(p-1)}}^{\frac{d}{2}(p-1)}. \tag{4.81}$$

Note that $A > 0$ by (4.75).

We take a subsequence of $\{f_n\}$ (still denoted by the same symbol) and a sequence $\{t_n\}$ in $\mathbb{R}$ such that

$$\inf_{n\in\mathbb{N}} \left\| e^{\frac{i}{2}t_n\Delta} f_n \right\|_{L^{\frac{d}{2}(p-1)}}^{\frac{d(p-1)}{2}} \ge A. \tag{4.82}$$

Here, extracting some subsequence of $\{t_n\}$, we may assume that

$$\lim_{n\to\infty} t_n = t_\infty \quad \text{for some } t_\infty \in \mathbb{R} \text{ or } t_\infty \in \{\pm\infty\}. \tag{4.83}$$

In addition to (4.82), we have by the Sobolev embedding that

$$\sup_{n\in\mathbb{N}} \left\| e^{\frac{i}{2}t_n\Delta} f_n \right\|_{L^{p+1}}^{p+1} \le C_p B^{p+1} \tag{4.84}$$

for some constant C_p depending only on d and p, where we put

$$B = \sup_{n\in\mathbb{N}} \|f_n\|_{H^1} < \infty. \tag{4.85}$$

Then, Lemma B.1 (with $\alpha = 2$, $\beta = \frac{d(p-1)}{2}$ and $\gamma = p+1$) shows that

$$\begin{aligned}\mathcal{L}^d\left(\left[\left|e^{\frac{i}{2}t_n\Delta}f_n\right| > \eta\right]\right) &> \frac{A}{2}\eta^{\frac{d}{2}(p-1)} \\ &\text{for all } n\in\mathbb{N} \text{ and } 0<\eta<\min\left\{1, \left(\tfrac{A}{4B^2}\right)^{\frac{1}{\frac{d(p-1)}{2}-2}}, \left(\tfrac{A}{4C_pB^{p+1}}\right)^{\frac{1}{p+1-\frac{d(p-1)}{2}}}\right\}.\end{aligned} \tag{4.86}$$

Moreover, Lemma B.2 implies that: There exists a sequence $\{y_n\}$ in $\mathbb{R}^d$ such that

$$\mathcal{L}^d\left(\left[\left|e^{\frac{i}{2}t_n\Delta}f_n(\cdot+y_n)\right| > \frac{\eta}{2}\right]\cap B_1(0)\right) \gtrsim \left(\frac{1+A\eta^{\frac{d}{2}(p-1)+2}}{1+B}\right)^{\frac{2}{2^\dagger-2}} \quad \text{for all } n\in\mathbb{N}, \tag{4.87}$$

where $2^\dagger = 4$ if $d = 1, 2$ and $2^\dagger = 2^*$ if $d \geq 3$, and the implicit constant depends only on d and p. For the sequence $\{y_n\}$ found above, we put

$$f_n^*(x) = e^{\frac{i}{2}t_n\Delta}f(x+y_n). \tag{4.88}$$

Then, $\{f_n^*\}$ is uniformly bounded in $H^1(\mathbb{R}^d)$ as well as $\{f_n\}$. Hence, there exist a subsequence of $\{f_n^*\}$ (still denoted by the same symbol), a function $f \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n\to\infty} f_n^* = f \quad \text{weakly in } H^1(\mathbb{R}^d). \tag{4.89}$$

Here, (4.89), with the help of the Sobolev embedding, also yields that

$$\lim_{n\to\infty} f_n^* = f \quad \text{strongly in } L^q_{loc}(\mathbb{R}^d) \quad \text{for all } q\in[2,2^*). \tag{4.90}$$

Therefore, we have by (4.87) that

$$\begin{aligned}\|f\|_{L^q}^q \geq \|f\|_{L^q(B_1(0))}^q = \lim_{n\to\infty}\|f_n^*\|_{L^q(B_1(0))}^q &\gtrsim \eta^q\left(\frac{1+A\eta^{\frac{d}{2}(p-1)+2}}{1+B}\right)^{\frac{2}{2^\dagger-2}} \\ &\text{for all } q\in[2,2^*) \text{ and } 0<\eta<\min\left\{1, \left(\tfrac{A}{4B^2}\right)^{\frac{1}{\frac{d(p-1)}{2}-2}}, \left(\tfrac{A}{4C_pB^{p+1}}\right)^{\frac{1}{p+1-\frac{d(p-1)}{2}}}\right\},\end{aligned} \tag{4.91}$$

where the implicit constant depends only on d and q. Thus, f is nontrivial.

Now, we shall show that the sequence $\{f_n^*\}$ satisfies the property (4.77). Indeed, it follows from the weak convergence (4.89) that

$$\begin{aligned}&\||\nabla|^s(f_n^*-f)\|_{L^2}^2 \\ &= \||\nabla|^s f_n^*\|_{L^2}^2 - \||\nabla|^s f\|_{L^2}^2 - 2\Re\int_{\mathbb{R}^d}|\nabla|^s(f_n^*(x)-f(x))\,|\nabla|^s f(x)\,dx \\ &= \||\nabla|^s f_n\|_{L^2}^2 - \||\nabla|^s f\|_{L^2}^2 + o_n(1) \qquad \text{for all } s\in[0,1],\end{aligned} \tag{4.92}$$

so that (4.77) holds.

Next, we shall show (4.78). We first consider the case $t_\infty = \lim_{n\to\infty} t_n \in \mathbb{R}$. Then, we can easily verify that

$$\lim_{n\to\infty} e^{-\frac{i}{2}t_n\Delta} f_n^* = e^{-\frac{i}{2}t_\infty\Delta} f \quad \text{weakly in } H^1(\mathbb{R}^d)\text{, and a.e. in } \mathbb{R}^d, \tag{4.93}$$

$$\lim_{n\to\infty} e^{-\frac{i}{2}t_n\Delta} f = e^{-\frac{i}{2}t_\infty\Delta} f \quad \text{strongly in } H^1(\mathbb{R}^d). \tag{4.94}$$

Moreover, the triangle inequality gives us that

$$\begin{aligned} &\left| \|f_n\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q \right| \\ &\le \left| \left\| e^{-\frac{i}{2}t_n\Delta} f_n^*\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f_n^* - e^{\frac{i}{2}t_\infty\Delta} f\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_\infty\Delta} f\right\|_{L^q}^q \right| \\ &\quad + \left| \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f_n^* - e^{-\frac{i}{2}t_\infty\Delta} f\right\|_{L^q}^q \right| \\ &\quad + \left| \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_\infty\Delta} f\right\|_{L^q}^q \right| \qquad \text{for all } q \in [2, 2^*). \end{aligned} \tag{4.95}$$

Here, we have by the Sobolev embedding that

$$\sup_{n\in\mathbb{N}} \left\| e^{-\frac{i}{2}t_n\Delta} f_n^*\right\|_{L^q} \lesssim \sup_{n\in\mathbb{N}} \left\| e^{-\frac{i}{2}t_n\Delta} f_n^*\right\|_{H^1} = \sup_{n\in\mathbb{N}} \|f_n\|_{H^1} < \infty \quad \text{for all } q \in [2, 2^*). \tag{4.96}$$

Therefore, Lemma B.4, together with (4.93) and (4.96), implies that the first term on the right-hand side of (4.95) vanishes as $n \to \infty$. The second term also vanishes as $n \to \infty$. Indeed, it follows from (D.1) and (4.94) that

$$\begin{aligned} &\lim_{n\to\infty} \left| \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f_n^* - e^{-\frac{i}{2}t_\infty\Delta} f\right\|_{L^q}^q \right| \\ &\lesssim \lim_{n\to\infty} \left(\|f_n\|_{H^1}^{q-1} + \|f\|_{H^1}^{q-1} \right) \left\| e^{-\frac{i}{2}t_n\Delta} f - e^{-\frac{i}{2}t_\infty\Delta} f\right\|_{L^q} = 0. \end{aligned} \tag{4.97}$$

Moreover, (4.94) immediately yields that the third term vanishes as $n \to \infty$. Thus, we have proved the property (4.78) when $t_\infty \in \mathbb{R}$.

We next suppose that $t_\infty \in \{\pm\infty\}$. In this case, (D.1) and Lemma 3.4 yield that

$$\begin{aligned} &\lim_{n\to\infty} \left| \|f_n\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q \right| \\ &= \lim_{n\to\infty} \left| \left\| e^{-\frac{i}{2}t_n\Delta} f_n^*\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q \right| \\ &\le \lim_{n\to\infty} \left| \left\| e^{-\frac{i}{2}t_n\Delta} f_n^*\right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n\Delta}(f_n^* - f)\right\|_{L^q}^q \right| + \lim_{n\to\infty} \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q \\ &\lesssim \lim_{n\to\infty} \left(\|f_n\|_{H^1}^{q-1} + \|f\|_{H^1}^{q-1} \right) \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q} + \lim_{n\to\infty} \left\| e^{-\frac{i}{2}t_n\Delta} f\right\|_{L^q}^q \\ &= 0 \qquad \text{for all } q \in (2, 2^*). \end{aligned} \tag{4.98}$$

Since (4.78) with $q = 2$ follows from (4.77), we have proved (4.78).

The property (4.79) immediately follows form (4.77) and (4.78).

Finally, we shall prove (4.80). It follows from (4.77) that

$$\|f_n^* - f\|_{H^1} \le \|f_n\|_{H^1} \quad \text{for sufficiently large } n \in \mathbb{N}. \tag{4.99}$$

Hence, extracting further some subsequence of $\{f_n^*\}$, we obtain (4.80). ☐

Lemma 4.5 (Dichotomy)**.** *Let u be a function in $X(\mathbb{R})$ and let $\{(\eta_n^1, \tau_n^1)\}$, …, $\{(\eta_n^L, \tau_n^L)\}$ be sequences in $\mathbb{R}^d \times \mathbb{R}$ with*

$$\lim_{n\to\infty} \Big(|\tau_n^k - \tau_n^l| + |\eta_n^k - \eta_n^l| \Big) = \infty \quad \textit{for all } 1 \le k < l \le L. \tag{4.100}$$

Then, putting $u_n^l(x,t) := u(x - \eta_n^l, t - \tau_n^l)$, we have

$$\lim_{n\to\infty} \left\| \left| \sum_{k=1}^{L} u_n^k \right|^{p-1} \sum_{l=1}^{L} u_n^l - \sum_{l=1}^{L} |u_n^l|^{p-1} u_n^l \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} = 0 \tag{4.101}$$

and

$$\lim_{n\to\infty} \left\| \left| u_n^k \right|^{q_j - 1} u_n^l \right\|_{L^{\frac{r_j}{q_j}}(\mathbb{R};L^1)} = 0 \quad \textit{for all } j = 1,2 \textit{ and } k \ne l, \tag{4.102}$$

where q_j and r_j ($j = 1,2$) are the exponents introduced in Section 3.1.

Proof of Lemma 4.5. Put

$$r_n = \frac{1}{2} \inf_{1 \le k < l \le L} \Big(\left|\tau_n^k - \tau_n^l\right| + \left|\eta_n^k - \eta_n^l\right| \Big) \tag{4.103}$$

and define functions χ_n and χ_n^l on $\mathbb{R}^d \times \mathbb{R}$ by

$$\chi_n(x,t) := \begin{cases} 1 & \text{if} \quad |(x,t)| < r_n, \\ 0 & \text{if} \quad |(x,t)| \ge r_n, \end{cases} \qquad \chi_n^l(x,t) := \chi_n(x - \eta_n^l, t - \tau_n^l). \tag{4.104}$$

Now, we shall prove (4.101). The triangle inequality and Lemma D.2 show that

$$\begin{aligned} &\left\| \left| \sum_{k=1}^{L} u_n^k \right|^{p-1} \sum_{l=1}^{L} u_n^l - \sum_{l=1}^{L} |u_n^l|^{p-1} u_n^l \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &\le \sum_{l=1}^{L} \left\| \left| \sum_{k=1}^{L} u_n^k \right|^{p-1} u_n^l - |u_n^l|^{p-1} u_n^l \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &\lesssim \sum_{l=1}^{L} \sum_{\substack{k=1 \\ k \ne l}}^{L} \left\| \left| u_n^k \right|^{p-1} |u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})}, \end{aligned} \tag{4.105}$$

where the implicit constant depends only on p and L. Hence, for (4.101), it suffices to show the following estimate:

$$\lim_{n\to\infty} \left\| \left| u_n^k \right|^{p-1} |u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} = 0 \quad \text{for all } k \ne l. \tag{4.106}$$

We begin with the following estimate:

$$\begin{aligned}\left\| \left|u_n^k\right|^{p-1} |u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \le{}& \left\| \left|u_n^k\right|^{p-1} |u_n^l| - \left|\chi_n^k u_n^k\right|^{p-1} |\chi_n^l u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &+ \left\| \left|\chi_n^k u_n^k\right|^{p-1} |\chi_n^l u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} .\end{aligned} \tag{4.107}$$

The Hölder inequality gives an estimate for the first term on the right-hand side of (4.107):

$$\begin{aligned}&\left\| \left|u_n^k\right|^{p-1} |u_n^l| - \left|\chi_n^k u_n^k\right|^{p-1} |\chi_n^l u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &\le \left\| \left|u_n^k\right|^{p-1} |u_n^l| - \left|\chi_n^k u_n^k\right|^{p-1} |u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &\qquad + \left\| \left|\chi_n^k u_n^k\right|^{p-1} |u_n^l| - \left|\chi_n^k u_n^k\right|^{p-1} |\chi_n^l u_n^l| \right\|_{L^{\widetilde{r}_1'}(\mathbb{R};L^{q_1'})} \\ &\lesssim \left\|(1-\chi_n^k)u_n^k\right\|_{X(\mathbb{R})}^{p-1} \left\|u_n^l\right\|_{X(\mathbb{R})} + \left\|u_n^k\right\|_{X(\mathbb{R})}^{p-1} \left\|(1-\chi_n^l)u_n^l\right\|_{X(\mathbb{R})} \\ &= \|(1-\chi_n)u\|_{X(\mathbb{R})}^{p-1} \|u\|_{X(\mathbb{R})} + \|u\|_{X(\mathbb{R})}^{p-1} \|(1-\chi_n)u\|_{X(\mathbb{R})} ,\end{aligned} \tag{4.108}$$

where the implicit constant depends only on d, p and q_1. Note here that

$$\lim_{n\to\infty} (1-\chi_n(x,t))u(x,t) = 0 \quad \text{a.a. } (x,t) \in \mathbb{R}^d \times \mathbb{R} \text{ and all } l = 1, \dots, L, \tag{4.109}$$

$$|(1-\chi_n(x,t))u(x,t)| \le |u(x,t)| \quad \text{a.a. } (x,t) \in \mathbb{R}^d \times \mathbb{R} \text{ and all } l = 1, \dots, L, \tag{4.110}$$

so that the Lebesgue dominated convergence theorem gives us that

$$\lim_{n\to\infty} \|(1-\chi_n)u\|_{X(\mathbb{R})} = 0. \tag{4.111}$$

Hence, the first term on the right-hand side of (4.107) vanishes as $n \to \infty$.

On the other hand, it follows from

$$\operatorname{supp} \chi_n^k \cap \operatorname{supp} \chi_n^l = \emptyset \quad \text{for } k \neq l \tag{4.112}$$

that the second term on the right-hand side of (4.107) is zero. Thus, we have proved (4.106).

In a way similar to the proof of (4.106), we can obtain (4.102). □

Now, we are in a position to prove Lemma 4.3.

Proof of Lemma 4.3. Put $f_n = \psi_n(0)$. Since $\psi_n(0) \in PW_+$, Proposition 2.5 and (4.62) give us that

$$\begin{aligned}\limsup_{n\to\infty}\mathcal{N}_2(f_n) &\le \limsup_{n\to\infty}\sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2}\widetilde{\mathcal{N}}_2(f_n)\\ &= \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2}\widetilde{N}_c \\ &< \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2}\widetilde{N}_2 = N_2.\end{aligned}\tag{4.113}$$

Now, we apply Lemma 4.4 to the sequence $\{f_n\}$ and obtain that: There exist a subsequence of $\{f_n\}$ (still denoted by the same symbol), a nontrivial function $f^1 \in H^1(\mathbb{R}^d)$, a sequence $\{t_n^1\}$ in $\mathbb{R}$ with $t_n^1 \to t_\infty^1 \in \mathbb{R}\cup\{\pm\infty\}$, and a sequence $\{y_n^1\}$ in $\mathbb{R}^d$ such that, putting

$$f_n^1(x) := \left(e^{\frac{i}{2}t_n^1\Delta}e^{y_n^1\cdot\nabla}f_n\right)(x) = e^{\frac{i}{2}t_n^1\Delta}f_n(x+y_n^1), \tag{4.114}$$

we have:

$$\lim_{n\to\infty} f_n^1 = f^1 \quad \text{weakly in } H^1(\mathbb{R}^d) \quad \text{for all } s\in[0,1], \tag{4.115}$$

$$\lim_{n\to\infty}\left\{\||\nabla|^s f_n\|_{L^2}^2 - \left\||\nabla|^s(f_n^1 - f^1)\right\|_{L^2}^2\right\} = \left\||\nabla|^s f^1\right\|_{L^2}^2 \quad \text{for all } s\in[0,1], \tag{4.116}$$

$$\lim_{n\to\infty}\left\{\|f_n\|_{L^q}^q - \left\|e^{-\frac{i}{2}t_n^1\Delta}(f_n^1-f^1)\right\|_{L^q}^q - \left\|e^{-\frac{i}{2}t_n^1\Delta}f^1\right\|_{L^q}^q\right\} = 0 \quad \text{for all } q\in[2,2^*), \tag{4.117}$$

$$\lim_{n\to\infty}\left\{\mathcal{H}(f_n) - \mathcal{H}(e^{-\frac{i}{2}t_n^1\Delta}(f_n^1-f^1)) - \mathcal{H}(e^{-\frac{i}{2}t_n^1\Delta}f^1)\right\} = 0, \tag{4.118}$$

$$\sup_{n\in\mathbb{N}}\left\|f_n^1 - f^1\right\|_{H^1} < \infty. \tag{4.119}$$

Besides, it follows from (4.113) and (4.116) that

$$\mathcal{N}_2(f^1) \le \limsup_{n\to\infty}\mathcal{N}_2(f_n) < N_2, \tag{4.120}$$

$$\limsup_{n\to\infty}\mathcal{N}_2(e^{-\frac{i}{2}t_n^1\Delta}(f_n^1-f^1)) = \limsup_{n\to\infty}\mathcal{N}_2(f_n^1-f^1) \le \limsup_{n\to\infty}\mathcal{N}_2(f_n) < N_2, \tag{4.121}$$

so that we have by the definition of N_2 (see (1.20)) that

$$0 < \mathcal{K}(f^1) < \mathcal{H}(f^1), \tag{4.122}$$

$$0 < \mathcal{K}(e^{-\frac{i}{2}t_n^1\Delta}(f_n^1-f^1)) < \mathcal{H}(e^{-\frac{i}{2}t_n^1\Delta}(f_n^1-f^1)) \quad \text{for sufficiently large } n\in\mathbb{N}. \tag{4.123}$$

We shall show that

$$\begin{cases} e^{-\frac{i}{2}t_\infty^1\Delta}f^1 \in PW_+ & \text{if} \quad t_\infty^1\in\mathbb{R}, \\ \qquad f^1\in\Omega & \text{if} \quad t_\infty^1 = \pm\infty. \end{cases}\tag{4.124}$$

Suppose first that $t_{\infty}^{1} \in \mathbb{R}$. Then, the estimate (4.113) gives us that

$$\mathcal{N}_2(e^{-\frac{i}{2}t_{\infty}^{1}\Delta} f^1) = \mathcal{N}_2(f^1) < N_2, \tag{4.125}$$

which, together with the definition of N_2, yields that

$$0 < \mathcal{K}(e^{-\frac{i}{2}t_{\infty}^{1}\Delta} f^1) < \mathcal{H}(e^{-\frac{i}{2}t_{\infty}^{1}\Delta} f^1). \tag{4.126}$$

Moreover, (4.116) and (4.118), with the help of (4.123), show that

$$\widetilde{\mathcal{N}}_2(e^{-\frac{i}{2}t_{\infty}^{1}\Delta} f^1) = \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(e^{-\frac{i}{2}t_{n}^{1}\Delta} f^1) \le \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(f_n). \tag{4.127}$$

Combining (4.127) with (4.62), we obtain that

$$\widetilde{\mathcal{N}}_2(e^{-\frac{i}{2}t_{\infty}^{1}\Delta} f^1) \le \widetilde{N}_c < \widetilde{N}_2. \tag{4.128}$$

Hence, (4.126) and (4.128), with the help of the relation (1.43), lead to that $e^{-t_{\infty}^{1}\Delta} f^1 \in PW_{+}$.

We next suppose that $t_{\infty}^{1} \in \{\pm\infty\}$. In this case , the formula (4.118), together with (4.123), yields that

$$\begin{aligned}\left\|\nabla f^1\right\|_{L^2}^{2} &= \mathcal{H}(e^{-\frac{i}{2}t_{n}^{1}\Delta} f^1) + \frac{2}{p+1}\left\|e^{-\frac{i}{2}t_{n}^{1}\Delta} f^1\right\|_{L^{p+1}}^{p+1} \\ &\le \mathcal{H}(f_n) + o_n(1) + \frac{2}{p+1}\left\|e^{-\frac{i}{2}t_{n}^{1}\Delta} f^1\right\|_{L^{p+1}}^{p+1},\end{aligned} \tag{4.129}$$

so that we have by Lemma 3.4 that

$$\left\|\nabla f^1\right\|_{L^2}^{2} \le \lim_{n\to\infty} \mathcal{H}(f_n). \tag{4.130}$$

This estimate and (4.116) with $s = 0$, together with (4.62), show that

$$\mathcal{N}_2(f^1) \le \lim_{n\to\infty} \|f_n\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}(f_n)}^{\frac{d}{2}(p-1)-2} = \widetilde{N}_c < \widetilde{N}_2. \tag{4.131}$$

Hence, we have that $f^1 \in \Omega$.

Now, we suppose that

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\left(f_n^1 - f^1\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} > 0. \tag{4.132}$$

Then, we can apply Lemma 4.4 to the sequence $\{f_n^1 - f^1\}$, so that we find that: There exist a subsequence of $\{f_n^1 - f^1\}$ (still denoted by the same symbol), a nontrivial function $f^2 \in H^1(\mathbb{R}^d)$, a sequence $\{t_n^2\}$ in $\mathbb{R}$ with $\lim_{n\to\infty} t_n^2 = t_\infty^2 \in \mathbb{R}\cup\{\pm\infty\}$ and a sequence $\{y_n^2\}$ in $\mathbb{R}^d$ such that, putting

$$f_n^2 := e^{\frac{i}{2}t_n^2\Delta} e^{y_n^2\cdot\nabla}\left(f_n^1 - f^1\right), \tag{4.133}$$

we have:

$$\lim_{n\to\infty} f_n^2 = f^2 \quad \text{weakly in } H^1(\mathbb{R}^d), \tag{4.134}$$

$$\lim_{n\to\infty} \left\{ \left\| |\nabla|^s \left(f_n^1 - f^1\right) \right\|_{L^2}^2 - \left\| |\nabla|^s \left(f_n^2 - f^2\right) \right\|_{L^2}^2 \right\} = \left\| |\nabla|^s f^2 \right\|_{L^2}^2 \quad \text{for all } s \in [0,1], \tag{4.135}$$

$$\lim_{n\to\infty} \left\{ \|f_n^1 - f^1\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n^2\Delta}(f_n^2 - f^2) \right\|_{L^q}^q - \left\| e^{-\frac{i}{2}t_n^2\Delta} f^2 \right\|_{L^q}^q \right\} = 0 \quad \text{for all } q \in [2, 2^*), \tag{4.136}$$

$$\lim_{n\to\infty} \left\{ \mathcal{H}(f_n^1 - f^1) - \mathcal{H}(e^{-\frac{i}{2}t_n^2\Delta}(f_n^2 - f^2)) - \mathcal{H}(e^{-\frac{i}{2}t_n^2\Delta} f^2) \right\} = 0, \tag{4.137}$$

$$\sup_{n\in\mathbb{N}} \left\| |\nabla|^s (f_n^2 - f^2) \right\|_{L^2} < \infty. \tag{4.138}$$

Note here that f_n is represented in the form

$$f_n = e^{-\frac{i}{2}\tau_n^1\Delta} e^{-\eta_n^1\cdot\nabla} f^1 + e^{-\frac{i}{2}\tau_n^2\Delta} e^{-\eta_n^2\cdot\nabla} f^2 + e^{-\frac{i}{2}\tau_n^2\Delta} e^{-\eta_n^2\cdot\nabla} \left(f_n^2 - f^2\right) \quad \text{for all } n \in \mathbb{N}, \tag{4.139}$$

where

$$\tau_n^1 := t_n^1, \quad \eta_n^1 := y_n^1, \qquad \tau_n^2 := \tau_n^1 + t_n^2, \quad \eta_n^2 := \eta_n^1 + y_n^2. \tag{4.140}$$

We also note that the following formulas hold:

$$\lim_{n\to\infty} \left\{ \| |\nabla|^s f_n \|_{L^2}^2 - \left\| |\nabla|^s \left(f_n^2 - f^2\right) \right\|_{L^2}^2 \right\} = \sum_{k=1}^{2} \left\| |\nabla|^s f^k \right\|_{L^2}^2 \quad \text{for all } s \in [0,1], \tag{4.141}$$

$$\lim_{n\to\infty} \left\{ \|f_n\|_{L^q}^q - \left\| e^{-\frac{i}{2}\tau_n^2\Delta}(f_n^2 - f^2) \right\|_{L^q}^q - \sum_{k=1}^{2} \left\| e^{-\frac{i}{2}\tau_n^k\Delta} f^k \right\|_{L^q}^q \right\} = 0 \quad \text{for all } q \in [2, 2^*), \tag{4.142}$$

$$\lim_{n\to\infty} \left\{ \mathcal{H}(f_n) - \mathcal{H}(e^{-\frac{i}{2}\tau_n^2\Delta}(f_n^2 - f^2)) - \sum_{k=1}^{2} \mathcal{H}(e^{-\frac{i}{2}\tau_n^k\Delta} f^k) \right\} = 0. \tag{4.143}$$

Moreover, in a similar way to the proofs of (4.123) and (4.124), we obtain

$$0 < \mathcal{K}\left(e^{-\frac{i}{2}\tau_n^2\Delta}\left(f_n^2 - f^2\right)\right) < \mathcal{H}\left(e^{-\frac{i}{2}\tau_n^2\Delta}\left(f_n^2 - f^2\right)\right) \tag{4.144}$$

and

$$\begin{cases} e^{-\frac{i}{2}\tau_\infty^2} f^2 \in PW_+ & \text{if } \tau_\infty^2 \in \mathbb{R}, \\ \qquad f^2 \in \Omega & \text{if } \tau_\infty^2 = \pm\infty. \end{cases} \tag{4.145}$$

We shall show that

$$\lim_{n\to\infty} |\tau_n^2 - \tau_n^1| + |\eta_n^2 - \eta_n^1| = \infty. \tag{4.146}$$

Supposing that (4.146) fails, we can take convergent subsequences of $\{\tau_n^2 - \tau_n^1\}$ and $\{\eta_n^2 - \eta_n^1\}$. Then, (4.115), together with the unitarity of the operators $e^{-\frac{i}{2}(\tau_n^2 - \tau_n^1)\Delta}$ and $e^{-(\eta_n^2 - \eta_n^1)\cdot\nabla}$ on $H^1(\mathbb{R}^d)$, shows that

$$\lim_{n\to\infty} f_n^2 = 0 \quad \text{weakly in } H^1(\mathbb{R}^d), \tag{4.147}$$

which contradicts the fact that f^2 is nontrivial. Thus, (4.146) holds.

Now, we suppose that

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\left(f_n^2 - f^2\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} > 0. \tag{4.148}$$

Then, we can repeat the above procedure; Iterative use of Lemma 4.4 implies the following Lemma:

Lemma 4.6. *For some subsequence of $\{f_n\}$ (still denoted by the same symbol), there exists*
(i) a family of nontrivial functions in $H^1(\mathbb{R}^d)$, $\{f^1, f^2, f^3, \ldots\}$, and
(ii) a family of sequences in $\mathbb{R}^d\times\mathbb{R}$, $\left\{\{(\eta_n^1,\tau_n^1)\},\{(\eta_n^2,\tau_n^2)\},\{(\eta_n^3,\tau_n^3)\},\ldots\right\}$ with

$$\lim_{n\to\infty}\tau_n^l = \tau_\infty^l \in \mathbb{R}\cup\{\pm\infty\} \quad \textit{for all } l\ge 1, \tag{4.149}$$

$$\left\{\begin{array}{ll} e^{-\frac{i}{2}\tau_\infty^l\Delta}f^l \in PW_+ & \textit{if} \quad \tau_\infty^l\in\mathbb{R}, \\ f^l\in\Omega & \textit{if} \quad \tau_\infty^l = \pm\infty \end{array}\right. \quad \textit{for all } l\ge 1, \tag{4.150}$$

and

$$\lim_{n\to\infty}|\tau_n^l-\tau_n^k| + |\eta_n^l - \eta_n^k| = \infty \quad \textit{for all } 1\le k<l, \tag{4.151}$$

such that, putting

$$f_n^0 := f_n, \quad f^0 := 0, \quad \tau_n^0 := 0, \quad \eta_n^0 := 0,$$

$$f_n^l := e^{\frac{i}{2}(\tau_n^l-\tau_n^{l-1})\Delta}e^{(\eta_n^l-\eta_n^{l-1})\cdot\nabla}(f_n^{l-1}-f^{l-1}) \quad \textit{for } l\ge 1,$$

we have, for all $l\ge 1$:

$$\lim_{n\to\infty} f_n^l = f^l \quad \textit{weakly in } H^1(\mathbb{R}^d)\textit{, and strongly in } L^q_{loc}(\mathbb{R}^d) \textit{ for all } q\in[2,2^*), \tag{4.152}$$

$$\lim_{n\to\infty}\left\{\||\nabla|^s f_n\|_{L^2}^2 - \left\||\nabla|^s\left(f_n^l - f^l\right)\right\|_{L^2}^2\right\} = \sum_{k=1}^{l}\left\||\nabla|^s f^k\right\|_{L^2}^2 \quad \textit{for all } s\in[0,1], \tag{4.153}$$

$$\lim_{n\to\infty}\left\{\|f_n\|_{L^q}^q - \left\|e^{-\frac{i}{2}\tau_n^l\Delta}\left(f_n^l-f^l\right)\right\|_{L^q}^q - \sum_{k=1}^{l}\left\|e^{-\frac{i}{2}\tau_n^k\Delta}f^k\right\|_{L^q}^q\right\} = 0 \quad \textit{for all } q\in[2,2^*), \tag{4.154}$$

$$\lim_{n\to\infty}\left\{\mathcal{H}(f_n) - \mathcal{H}(e^{-\frac{i}{2}\tau_n^l\Delta}\left(f_n^l-f^l\right)) - \sum_{k=1}^{l}\mathcal{H}(e^{-\frac{i}{2}\tau_n^k\Delta}f^k)\right\} = 0, \tag{4.155}$$

$$\mathcal{K}(e^{-\frac{i}{2}\tau_n^l\Delta}\left(f_n^l-f^l\right)) > 0. \tag{4.156}$$

Furthermore, putting $N := \#\{f^1, f^2, f^3, \ldots\}$, we have the alternatives: if N is finite, then

$$\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^N - f^N\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})\cap X(\mathbb{R})} = 0; \tag{4.157}$$

if $N=\infty$, then

$$\lim_{l\to\infty}\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^l - f^l\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})\cap X(\mathbb{R})} = 0. \tag{4.158}$$

Proof of Lemma 4.6. The last assertion (4.158) is nontrivial. We prove this, provided that the other properties are proved. We first show that

$$\lim_{l\to\infty}\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^l-f^l\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})}=0. \tag{4.159}$$

Suppose the contrary that there exists a constant $\varepsilon_0>0$ such that

$$\limsup_{l\to\infty}\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^l-f^l\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})}\ge\varepsilon_0. \tag{4.160}$$

Then, we can take an increasing sequence of indices $\{j(L)\}_{L\in\mathbb{N}}$ such that

$$\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^{j(L)}-f^{j(L)}\right)\right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})}\ge\frac{\varepsilon_0}{2}. \tag{4.161}$$

Here, it follows from the construction of the family $\{f^1,f^2,f^3,\ldots\}$ (see also (4.91) in the proof of Lemma 4.4) that there exists a constant $C(\varepsilon_0)>0$ depending only on d, p, ε_0 and $\sup_{n\in\mathbb{N}}\|f_n\|_{L^2}$ such that

$$\left\|f^{j(L)+1}\right\|_{L^2}\ge C(\varepsilon_0)\quad\text{for all } L\in\mathbb{N}. \tag{4.162}$$

Then, the uniform bound (4.61) (recall that $f_n=\psi_n(0)$), (4.153) and (4.162), yield that

$$\begin{aligned}\sup_{n\in\mathbb{N}}\|f_n\|_{L^2}^2&\ge\lim_{n\to\infty}\left\{\|f_n\|_{L^2}^2-\left\|f_n^{j(M)+1}-f^{j(M)+1}\right\|_{L^2}^2\right\}=\sum_{k=1}^{j(M)+1}\left\|f^k\right\|_{L^2}^2\\&\ge\sum_{L=1}^{M}\left\|f^{j(L)+1}\right\|_{L^2}^2\ge MC(\varepsilon_0)\quad\text{for all } M\in\mathbb{N}.\end{aligned} \tag{4.163}$$

Since $\sup_{n\in\mathbb{N}}\|f_n\|_{L^2}<\infty$, taking $M\to\infty$ in (4.163), we have a contradiction. Thus, (4.159) holds.

It remains to prove that

$$\lim_{l\to\infty}\lim_{n\to\infty}\left\|e^{\frac{i}{2}t\Delta}\left(f_n^l-f^l\right)\right\|_{X(\mathbb{R})}=0. \tag{4.164}$$

This estimate follows from Lemma 3.3 and (4.159). □

Proof of Lemma 4.3 (continued) We are back to the proof of Lemma 4.3.

We begin with proving that that $N=1$ in Lemma 4.6. To this end, we define an approximate solution ψ_n^{app} of ψ_n: Putting $L=N$ if $N<\infty$; L sufficiently large number specified later exactly if $N=\infty$, we define

$$\psi_n^{app}(x,t):=\sum_{l=1}^{L}\psi^l(x-\eta_n^l,t-\tau_n^l). \tag{4.165}$$

Here, each ψ^l is the solution to (4.9) (or (4.10)) with f^l just found in Lemma 4.6, and each (η_n^l,τ_n^l) is the sequence found in Lemma 4.6, so that we find that

$$\lim_{n\to\infty}\left\|\psi^l(-\tau_n^l)-e^{-\frac{i}{2}\tau_n^l\Delta}f^l\right\|_{H^1}=0\quad\text{for all } 1\le l\le L, \tag{4.166}$$

$$\psi^l(t)\in PW_+\quad\text{for all } 1\le l\le L \text{ and } t\in\mathbb{R}. \tag{4.167}$$

We note again that if $\tau_\infty^l = \pm\infty$, then ψ^l is the solution to the final value problem (4.10); since $f^l \in \Omega$ if $\tau_\infty^l = \pm\infty$ (see (4.150)), we actually obtain the desired solution ψ^l by Proposition 3.9.

Now, we shall show that

$$\|\psi^l\|_{X(\mathbb{R})} < \infty \quad \text{for all } 1 \leq l \leq L, \quad \text{if } N \geq 2. \tag{4.168}$$

For (4.168), it suffices to show that $\widetilde{\mathcal{N}}_2(\psi^l(0)) < \widetilde{N}_c$ by the definition of $\widetilde{N}_c$ (see (4.2)). Suppose $N \geq 2$, so that $L \geq 2$. Then, we employ (4.153), (4.155), (4.156) and obtain that

$$\begin{aligned}
\widetilde{\mathcal{N}}_2(f_n) &= \sqrt{\|f_n^L - f^L\|_{L^2}^2 + \sum_{k=1}^{L} \|f^k\|_{L^2}^2 + o_n(1)}^{\,p+1-\frac{d}{2}(p-1)} \\
&\quad \times \sqrt{\mathcal{H}\left(e^{-\frac{i}{2}\tau_n^L\Delta}\left(f_n^L - f^L\right)\right) + \sum_{k=1}^{L} \mathcal{H}\left(e^{-\frac{i}{2}\tau_n^k\Delta} f^k\right) + o_n(1)}^{\,\frac{d}{2}(p-1)-2} \\
&\geq \sqrt{\sum_{k=1}^{L} \|f^k\|_{L^2}^2}^{\,p+1-\frac{d}{2}(p-1)} \sqrt{\sum_{k=1}^{L} \mathcal{H}\left(e^{-\frac{i}{2}\tau_n^k\Delta} f^k\right)}^{\,\frac{d}{2}(p-1)-2} + o_n(1).
\end{aligned} \tag{4.169}$$

Since (4.150) and Lemma 3.4 imply that

$$\mathcal{H}\left(e^{-\frac{i}{2}\tau_n^k\Delta} f^k\right) > 0 \quad \text{for all } k \geq 1 \text{ and sufficiently large } n, \tag{4.170}$$

and since ψ^l is the solution to (4.9) or (4.10), we have from (4.62) and (4.169) that

$$\begin{aligned}
\widetilde{N}_c &= \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(f_n) \\
&> \lim_{n\to\infty} \left\|f^l\right\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}\left(e^{-\frac{i}{2}\tau_n^l\Delta} f^l\right)}^{\,\frac{d}{2}(p-1)-2} \\
&= \lim_{n\to\infty} \left\|\psi^l(-\tau_n)\right\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}\left(\psi^l(-\tau_n)\right)}^{\,\frac{d}{2}(p-1)-2} \\
&= \lim_{n\to\infty} \left\|\psi^l(0)\right\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \sqrt{\mathcal{H}\left(\psi^l(0)\right)}^{\,\frac{d}{2}(p-1)-2} \\
&= \widetilde{\mathcal{N}}_2(\psi^l(0)) \qquad \text{for all } 1 \leq l \leq L \text{ and sufficiently large } n,
\end{aligned} \tag{4.171}$$

Thus, (4.168) holds.

We know by (4.168) that

$$\sup_{n\in\mathbb{N}} \|\psi_n^{app}\|_{X(\mathbb{R})} < \infty. \tag{4.172}$$

Furthermore, we will see that: When $N = \infty$, there exists $A > 0$ with the property that for any $L \in \mathbb{N}$ (the number of components of ψ_n^{app}, see (4.165)), there exists $n_L \in \mathbb{N}$ such that

$$\sup_{n \geq n_L} \|\psi_n^{app}\|_{X(\mathbb{R})} \leq A. \tag{4.173}$$

We shall prove this fact. Recall that $X(\mathbb{R}) = L^{r_1}(\mathbb{R}; L^{q_1}) \cap L^{r_2}(I; L^{q_2})$. Lemma D.2 yields that

$$\begin{aligned}
&\|\psi_n^{app}(t)\|_{L^{q_j}}^{q_j} = \left\| \sum_{l=1}^{L} \psi^l(\cdot - \eta_n^l, t - \tau_n^l) \right\|_{L^{q_j}}^{q_j} \\
&\le \sum_{l=1}^{L} \left\| \psi^l(t - \tau_n^l) \right\|_{L^{q_j}}^{q_j} \\
&\quad + C_L \sum_{l=1}^{L} \sum_{\substack{k=1 \\ k \neq l}}^{L} \int_{\mathbb{R}^d} \left| \psi^k(x - \eta_n^k, t - \tau_n^k) \right| \left| \psi^l(x - \eta_n^l, t - \tau_n^l) \right|^{q_j - 1} dx \quad \text{for } j = 1, 2
\end{aligned} \tag{4.174}$$

for some constant $C_L > 0$ depending only on d, p, q_1 and L. Therefore, we have by the triangle inequality that

$$\begin{aligned}
&\|\psi_n^{app}\|_{L^{r_j}(\mathbb{R}; L^{q_j})}^{q_j} = \left\| \|\psi_n^{app}\|_{L^{q_j}}^{q_j} \right\|_{L^{\frac{r_j}{q_j}}(\mathbb{R})} \\
&\le \sum_{l=1}^{L} \left\| \psi^l \right\|_{L^{r_j}(\mathbb{R}; L^{q_j})}^{q_j} \\
&\quad + C_L \sum_{l=1}^{L} \sum_{\substack{k=1 \\ k \neq l}}^{L} \left\| \left| \psi^k(\cdot - \eta_n^k, \cdot - \tau_n^k) \right| \left| \psi^l(\cdot - \eta_n^l, \cdot - \tau_n^l) \right|^{q_j - 1} \right\|_{L^{\frac{r_j}{q_j}}(\mathbb{R}; L^1)} \\
&=: I_j + II_j \qquad \text{for } j = 1, 2.
\end{aligned} \tag{4.175}$$

We first consider the term I_j. The formula (4.153) and the uniform bound (4.61) show that

$$\sum_{l=1}^{\infty} \left\| f^l \right\|_{H^1}^2 < \infty, \tag{4.176}$$

so that

$$\lim_{l \to \infty} \left\| f^l \right\|_{H^1} = 0. \tag{4.177}$$

Hence, it follows from Proposition 3.7 (when $\tau_\infty^l \in \mathbb{R}$) and Proposition 3.9 (when $\tau_n^l = \pm\infty$) that: There exists $l_0 \in \mathbb{N}$, independent of L (the number of components of ψ_n^{app}), such that

$$\|\psi^l\|_{X(\mathbb{R})} \lesssim \|f^l\|_{H^1} \le 1 \quad \text{for all } l \ge l_0, \tag{4.178}$$

where the implicit constant depends only on d, p and q_1. Then, (4.168), (4.176) and (4.178), together with the fact $q_j > 2$ for $j = 1, 2$, lead to that

$$\begin{aligned}
I_j &= \sum_{l=1}^{l_0} \left\| \psi^l \right\|_{L^{r_j}(\mathbb{R}; L^{q_j})}^{q_j} + \sum_{l=l_0+1}^{\infty} \left\| \psi^l \right\|_{L^{r_j}(\mathbb{R}; L^{q_j})}^{q_j} \\
&\lesssim \sum_{l=1}^{l_0} \left\| \psi^l \right\|_{X(\mathbb{R})}^{q_j} + \sum_{l=l_0+1}^{\infty} \left\| f^l \right\|_{H^1}^2 < \infty,
\end{aligned} \tag{4.179}$$

where the implicit constant depends only on d, p and q_1.

Next, we consider the term II_j. Lemma 4.5, together with the condition (4.151), implies that: There exists $n_L \in \mathbb{N}$ such that

$$\left\| \left| \psi^k(\cdot - \eta_n^k, \cdot - \tau_n^k) \right| \left| \psi^l(\cdot - \eta_n^l, \cdot - \tau_n^l) \right|^{q_j - 1} \right\|_{L^{\frac{r_j}{q_j}}(\mathbb{R};L^1)} \le \frac{1}{C_L L^2} \quad \text{for all } n \ge n_L \text{ and } k \neq l. \tag{4.180}$$

Hence, we see that

$$II_j \le 1 \quad \text{for all } n \ge n_L. \tag{4.181}$$

Thus, we have proved (4.173).

We shall show that the case $N \ge 2$ can not occur. Note that ψ_n^{app} solves the following equation:

$$2i\frac{\partial}{\partial t}\psi_n^{app} + \Delta\psi_n^{app} + |\psi_n^{app}|^{p-1}\psi_n^{app} = e_n, \tag{4.182}$$

where

$$e_n(x,t) := |\psi_n^{app}(x,t)|^{p-1}\psi_n^{app}(x,t) - \sum_{l=1}^{L} \left| \psi^l(x - \eta_n^l, t - \tau_n^l) \right|^{p-1} \psi^l(x - \eta_n^l, t - \tau_n^l). \tag{4.183}$$

Proposition 3.8 (long time perturbation theory), with the help of (4.173), tells us that there exists $\varepsilon_1 > 0$, independent of L when $N = \infty$, with the following property: If there exists $n \ge n_L$ (n_L is the number found in (4.173) if $N = \infty$, $n_L = 1$ if $N < \infty$) such that

$$\left\| e^{\frac{i}{2}t\Delta} \left(\psi_n(0) - \psi_n^{app}(0) \right) \right\|_{X(\mathbb{R})} \le \varepsilon_1 \tag{4.184}$$

and

$$\|e_n\|_{L^{\tilde{r}'}(\mathbb{R};L^{q_1'})} \le \varepsilon_1, \tag{4.185}$$

then

$$\|\psi_n\|_{X(\mathbb{R})} < \infty. \tag{4.186}$$

In the sequel, we show that if $N \ge 2$, then (4.184) and (4.185) hold valid for some L, which shows $N = 1$ since (4.186) contradicts (4.63). It is worth while noting here that

$$f_n = \sum_{l=1}^{L} e^{-\frac{i}{2}\tau_n^l\Delta} e^{-\eta_n^l\cdot\nabla} f^l + e^{-\frac{i}{2}\tau_n^L\Delta} e^{-\eta_n^L\cdot\nabla} \left(f_n^L - f^L \right), \tag{4.187}$$

in other words,

$$e^{\frac{i}{2}t\Delta} f_n = \sum_{l=1}^{L} e^{\frac{i}{2}(t-\tau_n^l)\Delta} e^{-\eta_n^l\cdot\nabla} f^l + e^{\frac{i}{2}(t-\tau_n^L)\Delta} e^{-\eta_n^L\cdot\nabla} \left(f_n^L - f^L \right). \tag{4.188}$$

Invoking Lemma 4.5, we find that (4.185) holds for all sufficiently large n. We next consider (4.184). The formula (4.188), with the help of (3.8), shows that

$$\begin{aligned}
&\left\| e^{\frac{i}{2}t\Delta} \left(\psi_n(0) - \psi_n^{app}(0) \right) \right\|_{X(\mathbb{R})} = \left\| e^{\frac{i}{2}t\Delta} \left(f_n - \sum_{l=1}^{L} e^{-\eta_n^l \cdot \nabla} \psi^l(-\tau_n^l) \right) \right\|_{X(\mathbb{R})} \\
&\le \left\| e^{\frac{i}{2}(t-\tau_n^L)\Delta} e^{-\eta_n^L \cdot \nabla} \left(f_n^L - f^L \right) \right\|_{X(\mathbb{R})} \\
&\quad + \sum_{l=1}^{L} \left\| e^{\frac{i}{2}t\Delta} \left(e^{-\frac{i}{2}\tau_n^l \Delta} e^{-\eta_n^l \cdot \nabla} f^l - e^{-\eta_n^l \cdot \nabla} \psi^l(-\tau_n^l) \right) \right\|_{X(\mathbb{R})} \\
&\le \left\| e^{\frac{i}{2}t\Delta} \left(f_n^L - f^L \right) \right\|_{X(\mathbb{R})} + C \sum_{l=1}^{L} \left\| e^{-\frac{i}{2}\tau_n^l \Delta} f^l - \psi^l(-\tau_n^l) \right\|_{H^1},
\end{aligned} \tag{4.189}$$

where C is some constant depending only on d, p and q_1. Here, we have by (4.157) and (4.158) that

$$\begin{aligned}
&\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta} \left(f_n^L - f^L \right) \right\|_{X(\mathbb{R})} \le \frac{\varepsilon_1}{4} \\
&\qquad\qquad \text{for } L = N \text{ if } N < \infty, \text{ and sufficiently large } L \text{ if } N = \infty.
\end{aligned} \tag{4.190}$$

Hence, for all $L \in \mathbb{N}$ satisfying (4.190), there exists $n_{L,1} \in \mathbb{N}$ such that

$$\left\| e^{\frac{i}{2}t\Delta} \left(f_n^L - f^L \right) \right\|_{X(\mathbb{R})} \le \frac{\varepsilon_1}{2} \quad \text{for all } n \ge n_{L,1}. \tag{4.191}$$

Moreover, (4.166) shows that for all $L \le N$, there exists $n_{L,2} \in \mathbb{N}$ such that

$$\left\| e^{-\frac{i}{2}\tau_n^l \Delta} f^l - \psi^l(-\tau_n^l) \right\|_{H^1} \le \frac{\varepsilon_1}{2CL} \quad \text{for all } n \ge n_{L,2} \text{ and } 1 \le l \le L, \tag{4.192}$$

where C is the constant found in (4.189). Combining (4.189) with (4.191) and (4.192), we see that for all $L \in \mathbb{N}$ satisfying (4.190), there exists $n_{L,3} \in \mathbb{N}$ such that

$$\left\| e^{\frac{i}{2}t\Delta} \left(\psi_n(0) - \psi_n^{app}(0) \right) \right\|_{X(\mathbb{R})} \le \varepsilon_1 \quad \text{for all } n \ge n_{L,3}, \tag{4.193}$$

which gives (4.184).

We have just proved $N = 1$, and therefore L should be one;

$$\psi_n^{app}(x,t) = \psi^1(x - \eta_n^1, t - \tau_n^1) = \left(e^{-\tau_n^1 \frac{\partial}{\partial t} - \eta_n^1 \cdot \nabla} \psi^1 \right)(x,t).$$

Put $\Psi = \psi^1$, $f = f^1$, $(\gamma_n, \tau_n) = (\gamma_n^1, \tau_n^1)$ and $\tau_\infty = \tau_\infty^1$. Then, these are what we want. Indeed, we have already shown that these satisfy the properties (4.66), (4.69), (4.71), (4.72); see (4.167) for (4.66), (4.152) for (4.69), (4.157) for (4.71), and (4.166) for (4.72). Moreover, the property (4.74) immediately follows from (4.72).

It remains to prove (4.65), (4.67), (4.68), (4.70) and (4.73).

We first prove that Ψ satisfies the property (4.65): $\|\Psi\|_{X(\mathbb{R})} = \infty$. Suppose the contrary that $\|\Psi\|_{X(\mathbb{R})} < \infty$. Then, a quite similar argument above works well, so that we obtain an absurd conclusion

$$\|\psi_n\|_{X(\mathbb{R})} < \infty \quad \text{for sufficiently large } n \in \mathbb{N}. \tag{4.194}$$

Hence, we have proved (4.65).

In order to prove (4.68) and (4.70), we shall show that there exists a subsequence of $\{f_n\}$ (still denoted by the same symbol) such that

$$\|f\|_{L^2} = \lim_{n\to\infty} \|f_n\|_{L^2}\,, \tag{4.195}$$

$$\lim_{n\to\infty} \mathcal{H}\left(e^{-\frac{i}{2}\tau_n\Delta} f\right) = \lim_{n\to\infty} \mathcal{H}\left(f_n\right). \tag{4.196}$$

Since $f_n^1 = e^{\frac{i}{2}\tau_n^1\Delta} e^{\eta_n^1\cdot\nabla} f_n(0)$ and $f = f^1$, the weak convergence result (4.152), with the help of extraction of some subsequence, leads to that

$$\|f\|_{L^2} \le \lim_{n\to\infty} \|f_n\|_{L^2}\,. \tag{4.197}$$

Extracting some subsequence further, we also have by (4.155) and (4.156) that

$$\lim_{n\to\infty} \mathcal{H}\left(e^{-\frac{i}{2}\tau_n\Delta} f\right) \le \lim_{n\to\infty} \mathcal{H}\left(f_n\right). \tag{4.198}$$

Here, we employ the mass conservation law (1.5) and (4.72) (or the formula (4.166) with $l = 1$) to obtain that

$$\|\Psi(0)\|_{L^2} = \lim_{n\to\infty} \|\Psi(-\tau_n)\|_{L^2} = \|f\|_{L^2}\,. \tag{4.199}$$

Moreover, the energy conservation law (1.6) and (4.166) with $l = 1$ give us that

$$\mathcal{H}(\Psi(0)) = \lim_{n\to\infty} \mathcal{H}(\Psi(-\tau_n)) = \lim_{n\to\infty} \mathcal{H}(e^{-\frac{i}{2}\tau_n\Delta} f). \tag{4.200}$$

Hence, supposing the contrary that (4.195) or (4.196) fails, we have by the minimizing property (4.62) that

$$\begin{aligned}
\widetilde{\mathcal{N}}_2(\Psi(0)) &= \|f\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \lim_{n\to\infty} \sqrt{\mathcal{H}\left(e^{-\frac{i}{2}\tau_n\Delta} f\right)}^{\frac{d}{2}(p-1)-2} \\
&< \lim_{n\to\infty} \|f_n\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \lim_{n\to\infty} \sqrt{\mathcal{H}\left(f_n\right)}^{\frac{d}{2}(p-1)-2} \\
&= \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(f_n) = \widetilde{N}_c.
\end{aligned} \tag{4.201}$$

This estimate, together with the definition of $\widetilde{N}_c$ (see (4.2)), leads to that $\|\Psi\|_{X(\mathbb{R})} < \infty$, which contradicts $\|\Psi\|_{X(\mathbb{R})} = \infty$. Thus, (4.195) and (4.196) holds. Then, an estimate similar to (4.201) gives (4.68). Moreover, (4.70) follows from (4.153) with $l = 1$, $s = 0$ and (4.195).

The properties (4.70) and (4.72) (or (4.74)), together with (4.60) and the mass conservation law (1.5), yield (4.67). Moreover, the formula (4.73) follows from the estimate (4.189) with $N = 1$, (4.157) and (4.166). $\square$

4.3 Proof of Proposition 4.1

We shall prove Proposition 4.1, showing that the candidate Ψ found in Lemma 4.3 is actually one-soliton-like solution as $t \to +\infty$.

Proof of Proposition 4.1. The properties (4.12) and (4.13) have been obtained in Lemma 4.3. Moreover, (2.26) in Proposition 2.5, together with (4.12) and (4.13), yields (4.14).

We shall prove (4.15): the momentum of Ψ is zero. Using the Galilei transformation, we set

$$\Psi_\xi(x,t) := e^{\frac{i}{2}(x\xi - t|\xi|^2)}\Psi(x-\xi t, t), \quad \xi \in \mathbb{R}^d. \tag{4.202}$$

It is easy to verify that

$$\|\Psi_\xi(t)\|_{L^q} = \|\Psi(t)\|_{L^q} \quad \text{for all } \xi \in \mathbb{R}^d,\ q \in [2, 2^*) \text{ and } t \in \mathbb{R}, \tag{4.203}$$

$$\|\Psi_\xi\|_{X(\mathbb{R})} = \|\Psi\|_{X(\mathbb{R})} = \infty \quad \text{for all } \xi \in \mathbb{R}^d. \tag{4.204}$$

Moreover, a simple calculation, together with the mass and momentum conservation laws (1.5) and (1.7), shows that

$$\begin{aligned}
\|\nabla\Psi_\xi(t)\|_{L^2}^2 &= \left\| i\xi\Psi_\xi(t) + e^{\frac{i}{2}(x\xi - t|\xi|^2)}(\nabla\Psi)(x-\xi t, t)\right\|_{L_x^2}^2 \\
&= \int_{\mathbb{R}^d} |\xi|^2|\Psi_\xi(x,t)|^2\,dx + \int_{\mathbb{R}^d} |\nabla\Psi|^2(x,t)\,dt \\
&\qquad + 2\Re\int_{\mathbb{R}^d} \overline{i\xi\Psi_\xi}(x,t)e^{\frac{i}{2}(x\xi - t|\xi|^2)}(\nabla\Psi)(x-\xi t,t)\,dx \\
&= |\xi|^2\|\Psi(t)\|_{L^2}^2 + \|\nabla\Psi(t)\|_{L^2}^2 + 2\xi\cdot\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,t)\nabla\Psi(x,t)\,dx \\
&= |\xi|^2\|\Psi(0)\|_{L^2}^2 + \|\nabla\Psi(t)\|_{L^2}^2 + 2\xi\cdot\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx \quad \text{for all } \xi\in\mathbb{R}^d.
\end{aligned} \tag{4.205}$$

This equality (4.205), together with the energy conservation law (1.6), yields that

$$\begin{aligned}
\mathcal{H}(\Psi_\xi(t)) &= \|\nabla\Psi_\xi(t)\|_{L^2}^2 - \frac{2}{p+1}\|\Psi_\xi(t)\|_{L^{p+1}}^{p+1} \\
&= \|\nabla\Psi_\xi(t)\|_{L^2}^2 - \frac{2}{p+1}\|\Psi(t)\|_{L^{p+1}}^{p+1} \\
&= \mathcal{H}(\Psi(0)) + |\xi|^2\|\Psi(0)\|_{L^2}^2 + 2\xi\cdot\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx \quad \text{for all } \xi\in\mathbb{R}^d.
\end{aligned} \tag{4.206}$$

Put

$$\xi_0 = -\frac{\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx}{\|\Psi(0)\|_{L^2}}. \tag{4.207}$$

Then, we have by (4.205) and (4.206) that

$$\|\nabla\Psi_{\xi_0}(t)\|_{L^2}^2 = \|\nabla\Psi(t)\|_{L^2}^2 - \left(\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx\right)^2 \quad \text{for all } t\in\mathbb{R}, \tag{4.208}$$

$$\mathcal{H}(\Psi_{\xi_0}(t)) = \mathcal{H}(\Psi(0)) - \left(\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx\right)^2 \quad \text{for all } t\in\mathbb{R}. \tag{4.209}$$

Now, we suppose that

$$\Im\int_{\mathbb{R}^d}\overline{\Psi}(x,t)\nabla\Psi(x,t)\,dx = \Im\int_{\mathbb{R}^d}\overline{\Psi}(x,0)\nabla\Psi(x,0)\,dx \neq 0. \tag{4.210}$$

Then, by (4.203) with $q = 2$, (4.208), (2.26) in Proposition 2.5 and (4.12), we have that

$$\begin{aligned}
&\mathcal{N}_2(\Psi_{\xi_0}(t)) < \mathcal{N}_2(\Psi(t)) \\
&\qquad \le \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \widetilde{\mathcal{N}}_2(\Psi(t)) = \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \widetilde{N}_c \qquad (4.211)\\
&\qquad < \sqrt{\frac{d(p-1)}{d(p-1)-4}}^{\frac{d}{2}(p-1)-2} \widetilde{N}_2 = N_2 \qquad \text{for all } t \in \mathbb{R}.
\end{aligned}$$

This inequality, together with the definition of N_2 (see 1.20), implies that

$$\mathcal{K}(\Psi_{\xi_0}(t)) > 0 \quad \text{for all } t \in \mathbb{R}. \tag{4.212}$$

Moreover, using (4.203) with $q = 2$, (4.209), and (4.12) again, we obtain that

$$\widetilde{\mathcal{N}}_2(\Psi_{\xi_0}(t)) < \widetilde{N}_c. \tag{4.213}$$

Hence, it follows from the definition of $\widetilde{N}_c$ that $\|\Psi_{\xi_0}\|_{X(\mathbb{R})} < \infty$, which contradicts (4.204). Thus, the momentum of Ψ must be zero.[2]

Next, we shall prove the tightness of either family $\{\Psi(t)\}_{t\in[0,\infty)}$ or $\{\Psi(t)\}_{t\in(-\infty,0]}$ in $H^1(\mathbb{R}^d)$. Since $\|\Psi\|_{X(\mathbb{R}))} = \infty$, it holds that $\|\Psi\|_{X([0,\infty))} = \infty$ or $\|\Psi\|_{X((-\infty,0]))} = \infty$. The time reversibility of our equation (1.1) allows us to assume that $\|\Psi\|_{X([0,\infty))} = \infty$; if not, we consider $\overline{\Psi(x,-t)}$ instead of $\Psi(x,t)$.

We introduce the following quantity to employ Proposition E.2:

$$A := \sup_{R>0} \liminf_{t\to\infty} \sup_{y\in\mathbb{R}^d} \int_{|x-y|\le R} |\Psi(x,t)|^2 \, dx. \tag{4.214}$$

By the definition of A, we can take a sequence $\{t_n\}$ in $[0,\infty)$ such that

$$\lim_{n\to\infty} t_n = +\infty, \tag{4.215}$$

$$\sup_{y\in\mathbb{R}^d} \int_{|x-y|\le R} |\Psi(x,t_n)|^2 \, dx \le \left(1 + \frac{1}{n}\right) A \quad \text{for all } R > 0 \text{ and } n \in \mathbb{N}. \tag{4.216}$$

Put

$$\Psi_n(x,t) := \Psi(x, t + t_n). \tag{4.217}$$

We investigate the behavior of this Ψ_n in order to show $A = 1$. Note here that since $\|\Psi(t)\|_{L^2} = 1$ for all $t \in \mathbb{R}$, we have $A \le 1$, so that $A \ge 1$ implies $A = 1$.

---

[2] The proof of zero momentum is quite similar to the one of Proposition 4.1 in [23]. We can find an analogous trick in Appendix D of [46].

The sequence $\{\Psi_n(0)\}$ satisfies the conditions of Lemma 4.3:

$$\Psi_n(t) \in PW_+ \quad \text{for all } t \in \mathbb{R}, \tag{4.218}$$

$$\|\Psi_n(t)\|_{L^2} = 1 \quad \text{for all } n \in \mathbb{N} \text{ and } t \in \mathbb{R}, \tag{4.219}$$

$$\sup_{n\in\mathbb{N}} \|\Psi_n(0)\|_{H^1} < \infty, \tag{4.220}$$

$$\widetilde{\mathcal{N}}_2(\Psi_n(t)) = \widetilde{N}_c \quad \text{for all } n \in \mathbb{N} \text{ and } t \in \mathbb{R}, \tag{4.221}$$

$$\|\Psi_n\|_{X(\mathbb{R})} = \infty \quad \text{for all } n \in \mathbb{N}, \tag{4.222}$$

$$\limsup_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\Psi_n(0) \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} > 0. \tag{4.223}$$

The assertion of (4.223) may need to be verified. If (4.223) fails, then we have

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\Psi(t_n) \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})} = 0, \tag{4.224}$$

which leads us to the contradictory conclusion $\|\Psi\|_{X(\mathbb{R})} < \infty$ by the small data theory (Proposition 3.7).

We apply Lemma 4.3 to $\{\Psi_n\}$, so that there exists a subsequence of $\{\Psi_n\}$ (still denoted by the same symbol) with the following properties: There exists
(i) a nontrivial global solution $\Psi_\infty \in C(\mathbb{R}; H^1(\mathbb{R}^d))$ to the equation (1.1) with

$$\|\Psi_\infty\|_{X(\mathbb{R})} = \infty, \tag{4.225}$$

$$\Psi_\infty(t) \in PW_+ \quad \text{for all } t \in \mathbb{R}, \tag{4.226}$$

$$\widetilde{\mathcal{N}}_2(\Psi_\infty(t)) = \widetilde{N}_c \quad \text{for all } t \in \mathbb{R}, \tag{4.227}$$

and
(ii) a nontrivial function $\Phi \in PW_+$, a sequence $\{\tau_n\}$ in $\mathbb{R}$ with $\lim\limits_{n\to\infty} \tau_n = \tau_\infty \in \mathbb{R} \cup \{\pm\infty\}$, and a sequence $\{\gamma_n\}$ in $\mathbb{R}^d$, such that, putting

$$\widetilde{\Psi}_n := e^{\frac{i}{2}\tau_n\Delta} e^{\gamma_n\cdot\nabla}\Psi_n, \quad \widetilde{\epsilon}_n := \Psi_n - e^{-\tau_n\frac{\partial}{\partial t} - \gamma_n\cdot\nabla}\Psi_\infty, \tag{4.228}$$

we have:

$$\lim_{n\to\infty} \widetilde{\Psi}_n(0) = \Phi \quad \text{weakly in } H^1(\mathbb{R}^d), \text{ and a.e. in } \mathbb{R}^d, \tag{4.229}$$

$$\lim_{n\to\infty} \left\| \widetilde{\Psi}_n(0) - \Phi \right\|_{L^2} = 0, \tag{4.230}$$

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\left(\widetilde{\Psi}_n(0) - \Phi\right) \right\|_{L^\infty(\mathbb{R};L^{\frac{d}{2}(p-1)})\cap X(\mathbb{R})} = 0, \tag{4.231}$$

$$\lim_{n\to\infty} \left\| \Psi_\infty(-\tau_n) - e^{-\frac{i}{2}\tau_n\Delta}\Phi \right\|_{H^1} = 0, \tag{4.232}$$

$$\lim_{n\to\infty} \left\| e^{\frac{i}{2}t\Delta}\widetilde{\epsilon}_n(0) \right\|_{X(\mathbb{R})} = 0. \tag{4.233}$$

Note here that the estimate (3.8) gives us that

$$\lim_{T\to\infty}\left\|e^{\frac{i}{2}t\Delta}\Phi\right\|_{X((-\infty,-T])}=\lim_{T\to\infty}\left\|e^{\frac{i}{2}t\Delta}\Phi\right\|_{X([T,+\infty))}=0. \tag{4.234}$$

We claim that $\tau_\infty\in\mathbb{R}$. We suppose the contrary that $\tau_\infty=+\infty$ or $\tau_\infty=-\infty$. If $\tau_\infty=+\infty$, then (4.231) and (4.234) show that

$$\begin{aligned}\left\|e^{\frac{i}{2}t\Delta}\Psi_n(0)\right\|_{X((-\infty,0])}&=\left\|e^{\frac{i}{2}(t+\tau_n)\Delta}\Psi_n(0)\right\|_{X((-\infty,-\tau_n])}\\&\le\left\|e^{\frac{i}{2}t\Delta}\Phi\right\|_{X((-\infty,-\tau_n])}+\left\|e^{\frac{i}{2}t\Delta}\left(\widetilde{\Psi}_n(0)-\Phi\right)\right\|_{X((-\infty,-\tau_n])}\\&\to 0\quad\text{as } n\to\infty.\end{aligned} \tag{4.235}$$

Then, the small data theory (Proposition 3.7) leads us to that

$$\|\Psi\|_{X((-\infty,t_n])}=\|\Psi_n\|_{X((-\infty,0])}\lesssim 1\quad\text{for all sufficiently large } n\in\mathbb{N}, \tag{4.236}$$

where the implicit constant is independent of n. Hence, taking $n\to\infty$ in (4.236), we obtain that

$$\|\Psi\|_{X(\mathbb{R})}<\infty, \tag{4.237}$$

which contradicts $\|\Psi\|_{X(\mathbb{R})}=\infty$. Thus, the case $\tau_\infty=+\infty$ never happens. On the other hand, if $\tau_\infty=-\infty$, then we have by (4.231) and (4.234) that

$$\begin{aligned}\left\|e^{\frac{i}{2}t\Delta}\Psi_n(0)\right\|_{X([0,+\infty))}&=\left\|e^{\frac{i}{2}(t+\tau_n)\Delta}\Psi_n(0)\right\|_{X([-\tau_n,+\infty))}\\&\le\left\|e^{\frac{i}{2}t\Delta}\Phi\right\|_{X([-\tau_n,+\infty))}+\left\|e^{\frac{i}{2}t\Delta}\left(\widetilde{\Psi}_n(0)-\Phi\right)\right\|_{X((-\tau_n,+\infty))}\\&\to 0\quad\text{as } n\to\infty.\end{aligned} \tag{4.238}$$

Hence, the small data theory (Proposition 3.7) shows that

$$\|\Psi\|_{X([t_n,+\infty))}=\|\Psi_n\|_{X([0,+\infty)}<\infty\quad\text{for all sufficiently large } n, \tag{4.239}$$

so that

$$\|\Psi\|_{X([0,+\infty))}<\infty. \tag{4.240}$$

However, (4.240) contradicts the our working hypothesis $\|\Psi\|_{X([0,+\infty))}=\infty$. Thus, we always have $\tau_\infty\in\mathbb{R}$.

We establish the strong convergence in $L^{p+1}(\mathbb{R}^d)$:

$$\lim_{n\to\infty}e^{\gamma_n\cdot\nabla}\Psi(t_n)=\lim_{n\to\infty}e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)=e^{-\frac{i}{2}\tau_\infty\Delta}\Phi\quad\text{strongly in } L^{p+1}(\mathbb{R}^d). \tag{4.241}$$

The Gagliardo-Nirenberg inequality shows that

$$\begin{aligned}&\left\|e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)-e^{-\frac{i}{2}\tau_\infty\Delta}\Phi\right\|_{L^{p+1}}^{p+1}\\&\lesssim\left\|e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)-e^{-\frac{i}{2}\tau_\infty\Delta}\Phi\right\|_{L^{\frac{d}{2}(p-1)}}^{p-1}\left\|\nabla\left(e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)-e^{-\frac{i}{2}\tau_\infty\Delta}\Phi\right)\right\|_{L^2}^2\\&\le\left\|e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)-e^{-\frac{i}{2}\tau_\infty\Delta}\Phi\right\|_{L^{\frac{d}{2}(p-1)}}^{p-1}\left(\|\nabla\Psi_n(0)\|_{L^2}^2+\|\nabla\Phi\|_{L^2}^2\right),\end{aligned} \tag{4.242}$$

where the implicit constant depends only on d and p. This estimate, together with (4.220) and (4.231), immediately yields (4.241).

Next, we shall show that

$$\lim_{n\to\infty} e^{\gamma_n\cdot\nabla}\Psi(t_n) = \lim_{n\to\infty} e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0) = e^{-\frac{i}{2}\tau_\infty\Delta}\Phi \quad \text{strongly in } H^1(\mathbb{R}^d). \tag{4.243}$$

Since (4.230) implies the strong convergence in $L^2(\mathbb{R}^d)$, if (4.243) fails, then we have by the weak convergence (4.229) that

$$\liminf_{n\to\infty} \|\nabla\Psi_n(0)\|_{L^2} = \liminf_{n\to\infty} \left\|\nabla\widetilde{\Psi}_n(0)\right\|_{L^2} > \|\nabla\Phi\|_{L^2}, \tag{4.244}$$

so that we have by (4.241) that

$$\lim_{n\to\infty} \mathcal{H}(\Psi_n(0)) = \lim_{n\to\infty} \mathcal{H}(e^{-\frac{i}{2}\tau_n\Delta}\widetilde{\Psi}_n(0)) > \mathcal{H}(e^{-\frac{i}{2}\tau_\infty\Delta}\Phi). \tag{4.245}$$

Moreover, it follows from (4.230), (4.245) and (4.221) that

$$\widetilde{\mathcal{N}}_2(e^{-\frac{i}{2}\tau_\infty\Delta}\Phi) < \lim_{n\to\infty} \widetilde{\mathcal{N}}_2(\Psi_n(0)) = \widetilde{N}_c. \tag{4.246}$$

Since Ψ_∞ is the solution to (1.1) with $\Psi_\infty(-\tau_\infty) = e^{-\frac{i}{2}\tau_\infty\Delta}\Phi$ (see (4.232)), the estimate (4.246), together with the definition of $\widetilde{N}_c$ (see (4.2)), leads to that

$$\|\Psi_\infty\|_{X(\mathbb{R})} < \infty, \tag{4.247}$$

which contradicts (4.225). Thus, (4.243) must hold.

Now, we are in a position to prove $A = 1$. We take an arbitrarily small number $\varepsilon > 0$, and fix it in the following argument; we assume that $\varepsilon < \frac{1}{4}$ at least.

Let $R_\varepsilon > 0$ be a number such that

$$\int_{|x|\le R_\varepsilon} |\Phi(x)|^2\,dx \ge \|\Phi\|_{L^2}^2 - \frac{\varepsilon}{2} = 1 - \frac{\varepsilon}{2}, \tag{4.248}$$

where we have used the fact that $\|\Phi\|_{L^2} = 1$ (see (4.219) and (4.230)). Then, it follows from (4.243) and (4.248) that

$$\int_{|x|\le R} |\Psi(x+\gamma_n, t_n)|^2\,dx > 1-\varepsilon \quad \text{for all sufficiently large } n\in\mathbb{N} \text{ and } R \ge R_\varepsilon. \tag{4.249}$$

Combining (4.216) and (4.249), we obtain that

$$1-\varepsilon \le \left(1+\frac{1}{n}\right)A \quad \text{for all sufficiently large } n\in\mathbb{N} \text{ and } R\ge R_\varepsilon. \tag{4.250}$$

Taking $\varepsilon \to 0$ and $n\to\infty$ in (4.250), we see that $1 \le A$. Hence, $A = 1$ as stated above.

We apply Proposition E.2 to $|\Psi|^2$ and find that: There exist $R_\varepsilon > 0$, $T_\varepsilon > 0$ and a continuous path $y_\varepsilon \in C([T_\varepsilon,\infty);\mathbb{R}^d)$ such that

$$\int_{|x-y_\varepsilon(t)|<R} |\Psi(x,t)|^2\,dx > 1-\varepsilon \quad \text{for all } t\in[T_\varepsilon,\infty) \text{ and } R > R_\varepsilon. \tag{4.251}$$

We claim that, if necessary, taking R_ε much larger , the following inequality holds for the same path y_ε just found above:

$$\int_{|x-y_\varepsilon(t)|<R} |\nabla\Psi(x,t)|^2\, dx > \|\nabla\Psi(t)\|_{L^2}^2 - \varepsilon \quad \text{for all } t\in[T_\varepsilon,\infty) \text{ and } R\ge R_\varepsilon. \tag{4.252}$$

We prove this by contradiction: Suppose the contrary that for any $k\in\mathbb{N}$, there exists $t_k^0\in[T_\varepsilon,\infty)$ such that

$$\int_{|x|<k} |\nabla\Psi(x+y_\varepsilon(t_k^0),t_k^0)|^2 \le \left\|\nabla\Psi(t_k^0)\right\|_{L^2}^2 - \varepsilon. \tag{4.253}$$

Put

$$\Psi_k^0(x,t) := \Psi(x+y_\varepsilon(t_k^0), t+t_k^0). \tag{4.254}$$

Then, in a similar argument for Ψ_n, we obtain a subsequence of $\{\Psi_k^0\}$ (still denoted by the same symbol), a sequence $\{\tau_k^0\}$ in $\mathbb{R}$ with $\lim\limits_{k\to\infty}\tau_k^0=\tau_\infty^0\in\mathbb{R}$, a sequence $\{y_k^0\}$ in $\mathbb{R}^d$, and a nontrivial function $\Phi^0\in H^1(\mathbb{R}^d)$ such that

$$\lim_{k\to\infty}\Psi_k^0(\cdot+y_k^0,0) = e^{-\frac{i}{2}\tau_\infty^0\Delta}\Phi^0 \quad \text{strongly in } H^1(\mathbb{R}^d). \tag{4.255}$$

Furthermore, (4.255) implies that there exists $\widetilde{R}_\varepsilon>0$ such that

$$\int_{|x-y_k^0|>\widetilde{R}_\varepsilon} \left|\Psi(x+y_\varepsilon(t_k^0),t_k^0)\right|^2 dx \le \varepsilon \quad \text{for all sufficiently large } k\in\mathbb{N}, \tag{4.256}$$

$$\int_{|x-y_k^0|\le\widetilde{R}_\varepsilon} \left|\nabla\Psi(x+y_\varepsilon(t_k^0),t_k^0)\right|^2 dx > \left\|\nabla\Psi(t_k^0)\right\|_{L^2}^2 - \varepsilon \quad \text{for all sufficiently large } k\in\mathbb{N}. \tag{4.257}$$

On the other hand, we have by (4.251) that

$$\int_{|x|\le R_\varepsilon} \left|\Psi(x+y_\varepsilon(t_k^0),t_k^0)\right|^2 dx > 1-\varepsilon. \tag{4.258}$$

If $\sup\limits_{k\in\mathbb{N}}|y_k^0|=\infty$, then we can take a subsequence of $\{y_k^0\}$ (still denoted by the same symbol) such that $\lim\limits_{k\to\infty}|y_k^0|=\infty$, so that

$$\left\{x\in\mathbb{R}^d \,\middle|\, |x-y_k^0|>\widetilde{R}_\varepsilon\right\} \supset \left\{x\in\mathbb{R}^d \,|\, |x|\le R_\varepsilon\right\} \quad \text{for all sufficiently large } k\in\mathbb{N}. \tag{4.259}$$

However, the result derived from (4.256) and (4.259) contradicts (4.258). Hence, it must hold that $\sup\limits_{k\in\mathbb{N}}|y_k^0|<\infty$. Put $R_0:=\sup\limits_{k\in\mathbb{N}}|y_k^0|$. Then, it follows from (4.257) that

$$\begin{aligned}\int_{|x|\le\widetilde{R}_\varepsilon+R_0} \left|\nabla\Psi(x+y_\varepsilon(t_k^0),t_k^0)\right|^2 dx &\ge \int_{|x-y_k^0|\le\widetilde{R}_\varepsilon} \left|\nabla\Psi(x+y_\varepsilon(t_k^0),t_k^0)\right|^2 dx \\ &> \left\|\nabla\Psi(t_k^0)\right\|_{L^2}^2 - \varepsilon,\end{aligned} \tag{4.260}$$

which contradicts (4.253) for $k>\widetilde{R}_\varepsilon+R_0$. Thus, (4.252) holds.

Finally, we obtain (4.16) and (4.17) from (4.251) and (4.252), respectively, by a space-time translation: regarding $\Psi(x+y_\varepsilon(T_\varepsilon),t+T_\varepsilon)$ as our $\Psi(x,t)$ with $\gamma_\varepsilon(t)=y_\varepsilon(t+T_\varepsilon)-y_\varepsilon(T_\varepsilon)$. ☐

4.4 Proofs of Theorem 1.1 and Corollary 1.5

We begin with the proof of Theorem 1.1.

Proof of Theorem 1.1. The claims (i) and (ii) are direct consequences of Proposition 2.5.

We shall prove (iii). Proposition 3.9 shows that the wave operators $W_\pm$ exist on Ω and continuous. It remains to prove the bijectivity of $W_\pm$ and the continuity of $W_\pm^{-1}$.

We first show that $W_\pm$ is surjective from Ω to PW_+. Let $\psi_0 \in PW_+$ and let ψ be the solution to (1.1) with $\psi(0) = \psi_0$. Since we have shown that $\widetilde{N}_c = \widetilde{N}_2$ in Section 4.1, $\|\psi\|_{X(\mathbb{R})} < \infty$. Therefore, it follows from Proposition 3.5 that ψ has asymptotic states $\phi_\pm$ in $H^1(\mathbb{R}^d)$. It remains to prove that $\phi_\pm \in \Omega$.

The energy conservation law (1.6) and Theorem 1.4 show that

$$0 < \mathcal{H}(\psi_0) = \lim_{t\to\pm\infty} \left(\|\nabla\psi(t)\|_{L^2}^2 - \frac{2}{p+1} \|\psi(t)\|_{L^{p+1}}^{p+1} \right) = \|\nabla\phi_\pm\|_{L^2}^2 . \tag{4.261}$$

Moreover, the mass conservation law (1.5) gives us that

$$\|\phi_\pm\|_{L^2}^2 = \lim_{t\to\pm\infty} \|\psi(t)\|_{L^2}^2 = \|\psi_0\|_{L^2}^2 . \tag{4.262}$$

Since $\psi_0 \in PW_+$, we obtain from (4.261), (4.262) and (1.44) that

$$\mathcal{N}_2(\phi_\pm) = \widetilde{\mathcal{N}}_2(\psi_0) < \widetilde{N}_2, \tag{4.263}$$

so that $\phi_\pm \in \Omega$.

Next, we shall show the injectivity of $W_\pm$. Take any $\phi_{+,1}, \phi_{+,2} \in \Omega$ and suppose that

$$W_+\phi_{+,1} = W_+\phi_{+,2}. \tag{4.264}$$

Put $\psi_0 = W_+\phi_{+,1} = W_+\phi_{+,2}$ and let ψ be the solution to the equation (1.1) with $\psi(0) = \psi_0$. Then, we have

$$\begin{aligned} \|\phi_{+,1} - \phi_{+,2}\|_{H^1} &\le \lim_{t\to+\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_{+,1} \right\|_{H^1} + \lim_{t\to+\infty} \left\| \psi(t) - e^{\frac{i}{2}t\Delta}\phi_{+,2} \right\|_{H^1} \\ &= 0, \end{aligned} \tag{4.265}$$

so that $\phi_{+,1} = \phi_{+,2}$. Similarly, we see that W_- is injective from Ω to PW_+.

Finally, we shall prove that W_+^{-1} is continuous. Take a function $\psi_0 \in PW_+$ and a sequence $\{\psi_{0,n}\}$ in PW_+ such that

$$\lim_{n\to\infty} \psi_{0,n} = \psi_0 \quad \text{strongly in } H^1(\mathbb{R}^d). \tag{4.266}$$

Let ψ and ψ_n be the solutions to (1.1) with $\psi(0) = \psi_0$ and $\psi_n(0) = \psi_{n,0}$. Since $\psi_0, \psi_{0,n} \in PW_+$ and $\widetilde{N}_c = \widetilde{N}_2$, we have by (4.2) that

$$\|\psi\|_{X(\mathbb{R})} < \infty, \quad \|\psi_n\|_{X(\mathbb{R})} < \infty. \tag{4.267}$$

Note that since $\|\psi\|_{X(\mathbb{R})} < \infty$, for any $\delta > 0$, there exists $T_\delta > 0$ such that

$$\|\psi\|_{X([T_\delta,\infty))} < \delta. \tag{4.268}$$

Now, we put

$$w_n = \psi_n - \psi. \tag{4.269}$$

Then, it follows from the inhomogeneous Strichartz estimate (3.11), an elementary inequality (D.2) and the Hölder inequality that

$$\begin{aligned} \|w_n\|_{X([t_0,t_1))} &\le \left\| e^{\frac{i}{2}(t-t_0)\Delta} w(t_0) \right\|_{X([t_0,t_1))} \\ &\quad + C\left(\|\psi\|_{X([t_0,t_1))}^{p-1} + \|w_n\|_{X([t_0,t_1))}^{p-1} \right) \|w_n\|_{X([t_0,t_1))} \\ &\qquad\qquad \text{for all } 0 \le t_0 < t_1 \le \infty, \end{aligned} \tag{4.270}$$

where C is some constant depending only on d, p and q_1. We claim that

$$\lim_{n\to\infty} \|w_n\|_{X([T_\delta,\infty))} = 0 \quad \text{for all } \delta \in \left(0,\ \left(\tfrac{1}{1+4C)}\right)^{\frac{1}{p-1}}\right), \tag{4.271}$$

where C is the constant found in (4.270) and T_δ is the time satisfying (4.268). Let ε be a number such that

$$0 < \varepsilon < \left(\frac{1}{(1+4C)^p} \right)^{\frac{1}{p-1}}. \tag{4.272}$$

Then, the estimate (3.8) and the continuous dependence of solutions on initial data show that there exists $N_\varepsilon \in \mathbb{N}$ such that

$$\left\| e^{\frac{i}{2}(t-T_\delta)} w_n(T_\delta) \right\|_{X(\mathbb{R})} < \varepsilon \quad \text{for all } n \ge N_\varepsilon. \tag{4.273}$$

We define a time $T_{\delta,n}$ by

$$T_{\delta,n} = \sup\left\{ t \ge T_\delta \mid \|w_n\|_{X([T_\delta,t))} \le (1+4C)\varepsilon \right\}. \tag{4.274}$$

For (4.271), it suffices to show that $T_{\delta,n} = \infty$ for all $n \ge N_\varepsilon$. We shall prove this. Note that (4.270) with $t_0 = T_\delta$, together with (4.273), implies that $T_{\delta,n} > T_\delta$ for all $n \ge N_\varepsilon$. Supposing the contrary that $T_{\delta,n} < \infty$ for some $n \ge N_\varepsilon$, we have from the continuity of w_n that

$$\|w_n\|_{X([T_\delta,T_{\delta,n}])} = (1+4C)\varepsilon. \tag{4.275}$$

However, (4.270), together with (4.268), (4.272), (4.273) and (4.275), shows that

$$\begin{aligned} \|w_n\|_{X([T_\delta,T_{\delta,n}])} &\le \varepsilon + C\left(\delta^{p-1} + (1+4C)^{p-1}\varepsilon^{p-1}\right)(1+4C)\varepsilon \\ &\le \varepsilon + 2C\varepsilon \qquad \text{for all } \delta \in \left(0,\ \left(\tfrac{1}{1+4C)}\right)^{\frac{1}{p-1}}\right) \text{ and } n \ge N_\varepsilon. \end{aligned} \tag{4.276}$$

This is a contradiction. Thus, we see that $T_{\delta,n} = \infty$ for all $n \ge N_\varepsilon$.

Besides (4.271), we see that there exists $\delta_0 > 0$ depending only on d, p and q_1 such that

$$\lim_{n\to\infty} \|w_n\|_{S([T_\delta,\infty))} = 0 \quad \text{for all } \delta \in (0,\delta_0). \tag{4.277}$$

Indeed, the Strichartz estimate, together with (D.2), yields that

$$\|w_n\|_{S([T_\delta,\infty))} \lesssim \|w_n(T_\delta)\|_{L^2} + \left(\|\psi\|_{X([T_\delta,\infty))}^{p-1} + \|w_n\|_{X([T_\delta,\infty))}^{p-1} \right) \|w_n\|_{S([T_\delta,\infty))}, \tag{4.278}$$

where the implicit constant depends only on d, p and q_1. This estimate, together with (4.271), gives (4.277).

We claim that

$$\lim_{n\to\infty} \|\psi_n - \psi\|_{L^\infty([T,\infty);H^1)} = \lim_{n\to\infty} \|w_n\|_{L^\infty([T,\infty);H^1)} = 0 \quad \text{for sufficiently large } T > 0. \tag{4.279}$$

Considering the integral equations of $\partial_j \psi$ and $\partial_j \psi_n$ for $1 \le j \le n$, we obtain that

$$\begin{aligned} \|\partial_j \psi_n - \partial_j \psi\|_{S([T_\delta,\infty))} &\lesssim \|\psi_n(T_\delta) - \psi(T_\delta)\|_{H^1} + \left\| |\psi_n|^{p-1} (\partial_j \psi_n - \partial_j \psi) \right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\ &\quad + \left\| \left(|\psi_n|^{p-1} - |\psi|^{p-1} \right) \partial_j \psi \right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\ &\quad + \left\| |\psi_n|^{p-3} \psi_n^2 \left(\partial_j \overline{\psi_n} - \partial_j \overline{\psi} \right) \right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\ &\quad + \left\| \left(|\psi_n|^{p-3} \psi_n^2 - |\psi|^{p-3} \psi^2 \right) \partial_j \overline{\psi} \right\|_{L^{r_0'}([T_\delta,\infty);L^{q_1'})} \\ &\lesssim \|\psi_n(T_\delta) - \psi(T_\delta)\|_{H^1} + \|\psi_n\|_{X([T_\delta,\infty)}^{p-1} \|\partial_j \psi_n - \partial_j \psi\|_{S([T_\delta,\infty))} \\ &\quad + \left\| |\psi_n|^{p-1} - |\psi|^{p-1} \right\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})} \|\partial_j \psi\|_{S([T_\delta,\infty))} \\ &\quad + \|\psi_n\|_{X([T_\delta,\infty))}^{p-1} \|\partial_j \psi_n - \partial_j \psi\|_{S([T_\delta,\infty))} \\ &\quad + \left\| |\psi_n|^{p-3} \psi_n^2 - |\psi|^{p-3} \psi^2 \right\|_{L^{\frac{r_2}{p-1}}([T_\delta,\infty);L^{\frac{q_2}{p-1}})} \|\partial_j \psi\|_{S([T_\delta,\infty))}, \end{aligned} \tag{4.280}$$

where the implicit constant depends only on d, p and q_1. This estimate, with the help of (4.271) and (4.277), leads to (4.279).

Now, put $\phi_{+,n} = W_+^{-1} \psi_{0,n}$. Then, we have by (4.279) that

$$\begin{aligned} \|\phi_{+,n} - \phi_+\|_{H^1} &\le \left\| \psi_n(t) - e^{\frac{i}{2}t\Delta} \phi_{+,n} \right\|_{H^1} + \left\| \psi(t) - e^{\frac{i}{2}t\Delta} \phi_+ \right\|_{H^1} \\ &\quad + \|\psi_n(t) - \psi(t)\|_{H^1} \\ &\to 0 \qquad \text{as } t \to \infty \text{ and } n \to \infty. \end{aligned} \tag{4.281}$$

Thus, we have proved that W_+ is a homeomorphism from Ω to PW_+. Similarly, we can prove that W_- is a homeomorphism from Ω to PW_+. ☐

We shall give the proof of Corollary 1.5.

Proof of Corollary 1.5. Let $f_1, f_2 \in PW_+ \cup \{0\}$. Then, Theorem 1.1 shows that the corresponding solutions ψ_1 and ψ_2 have asymptotic states at $+\infty$. Hence, it follows from Theorem 1.4 that

$$\lim_{t\to\infty} \|\psi_1(t)\|_{L^{p+1}} = \lim_{t\to\infty} \|\psi_2(t)\|_{L^{p+1}} = 0, \tag{4.282}$$

so that we can obtain

$$\lim_{t\to\infty} \|\psi_j(t)\|_{L^q} = 0 \quad \text{for all } q \in (2, 2^*) \text{ and } j = 1, 2. \tag{4.283}$$

Since the solutions are continuous in $H^1(\mathbb{R}^d)$, we find by (4.283) that f_1 and f_2 are connected by the path $\{\psi_1(t) \mid t \ge 0\} \cup \{0\} \cup \{\psi_2(t) \mid t \ge 0\}$ in $L^q(\mathbb{R}^d)$ with $q \in (2, 2^*)$. ☐

5 Analysis on PW_-

We shall give the proofs of Theorem 1.2, Theorem 1.6, Proposition 1.3 and Proposition 1.4.

We begin with the proof of Theorem 1.2. We employ the idea of Nawa [46]. Here, the generalized virial identity below (see (A.20)) plays an important role:

$$\begin{aligned}&(W_R, |\psi(t)|^2)\\&= (W_R, |\psi_0|^2) + 2t\Im(\vec{w}_R \cdot \nabla\psi_0, \psi_0) + 2\int_0^t\int_0^{t'} \mathcal{K}(\psi(t''))\,dt''dt'\\&\quad - 2\int_0^t\int_0^{t'} \mathcal{K}^R(\psi(t''))\,dt''dt' - \frac{1}{2}\int_0^t\int_0^{t'} (\Delta(\operatorname{div}\vec{w}_R), |\psi(t'')|^2)\,dt''dt'.\end{aligned}$$

Proof of Theorem 1.2. Since we have already proved (1.52) (see Proposition 2.6), proofs of (1.53) and (1.54) remain. For simplicity, we consider the forward time only. The problem for the backward time can be proved in a similar way.

Take any $\psi_0 \in PW_-$ and let ψ be the corresponding solution to our equation (1.1) with $\psi(0) = \psi_0$. When the maximal existence time $T_{\max}^+ < \infty$, we have (1.53) as mentioned in (1.4). Therefore, it suffices to prove (1.54).

We suppose the contrary that (1.54) fails when $T_{\max}^+ = \infty$, so that there exists $R_0 > 0$ such that

$$M_0 := \sup_{t\in[0,\infty)} \int_{|x|\geq R_0} |\nabla\psi(x,t)|^2\,dx < \infty. \tag{5.1}$$

Then, we shall derive a contradiction in three steps.
Step1. We claim that: there exists a constant $m_0 > 0$ such that

$$m_0 < \inf\left\{ \int_{|x|\geq R} |v(x)|^2\,dx \;\middle|\; \begin{array}{ll} v \in H^1(\mathbb{R}^d), & \mathcal{K}^R(v) \leq -\frac{1}{4}\varepsilon_0, \\ \|\nabla v\|_{L^2(|x|\geq R)}^2 \leq M_0, & \|v\|_{L^2} \leq \|\psi_0\|_{L^2} \end{array} \right\} \text{ for all } R > 0, \tag{5.2}$$

where $\varepsilon_0 = \mathcal{B}(\psi_0) - \mathcal{H}(\psi_0) > 0$. Let us prove this claim. Take any $v \in H^1(\mathbb{R}^d)$ with the following properties:

$$\mathcal{K}^R(v) \leq -\frac{1}{4}\varepsilon_0, \quad \|\nabla v\|_{L^2(|x|\geq R)}^2 \leq M_0, \quad \|v\|_{L^2} \leq \|\psi_0\|_{L^2}. \tag{5.3}$$

Note here that the second and third properties in (5.3) shows that

$$\|v\|_{H^1(|x|\geq R)}^2 \leq \|\psi_0\|_{L^2}^2 + M_0. \tag{5.4}$$

We also have by the first property in (5.3) and (A.28) that

$$\frac{1}{4}\varepsilon_0 \leq -\mathcal{K}^R(v) \leq \int_{|x|\geq R} \rho_3(x)|v(x)|^{p+1}\,dx. \tag{5.5}$$

Now, we define p_* by

$$p_* = \begin{cases} 2p-1 & \text{if} \quad d = 1, 2, \\ 2^*-1 & \text{if} \quad d \geq 3. \end{cases} \tag{5.6}$$

Then, the Hölder inequality and the Sobolev embedding show that

$$\begin{aligned}\int_{|x|\ge R} \rho_3(x)|v(x)|^{p+1}\,dx &\le \|\rho_3\|_{L^\infty}\|v\|_{L^2(|x|\ge R)}^{p+1-\frac{(p-1)(p_*+1)}{p_*-1}}\|v\|_{L^{p_*+1}(|x|\ge R)}^{\frac{(p-1)(p_*+1)}{p_*-1}} \\ &\lesssim \|\rho_3\|_{L^\infty}\|v\|_{L^2(|x|\ge R)}^{p+1-\frac{(p-1)(p_*+1)}{p_*-1}}\|v\|_{H^1(|x|\ge R)}^{\frac{(p-1)(p_*+1)}{p_*-1}},\end{aligned} \tag{5.7}$$

where the implicit constant depends only on d and p. This estimate, together with (5.4) and (5.5), yields that

$$\frac{\varepsilon_0}{\|\rho_3\|_{L^\infty}\sqrt{\|\psi_0\|_{L^2}^2+M_0}^{\frac{(p-1)(p_*+1)}{p_*-1}}} \lesssim \|v\|_{L^2(|x|\ge R)}^{p+1-\frac{(p-1)(p_*+1)}{p_*-1}}, \tag{5.8}$$

where the implicit constant depends only on d and p. Since $\|\rho_3\|_{L^\infty} \lesssim 1$ (see (A.31)), the estimate (5.8) gives us the desired result (5.2).

Step2. Let m_0 be a constant found in (5.2). Then, we prove that

$$\sup_{0\le t<\infty}\int_{|x|\ge R}|\psi(x,t)|^2\,dx \le m_0 \tag{5.9}$$

for all R satisfying the following properties:

$$R > R_0, \tag{5.10}$$

$$\frac{10d^2K}{R^2}\|\psi_0\|_{L^2}^2 < \varepsilon_0, \tag{5.11}$$

$$\int_{|x|\ge R}|\psi_0(x)|^2\,dx < m_0, \tag{5.12}$$

$$\frac{1}{R^2}\left(1+\frac{2}{\varepsilon_0}\|\nabla\psi_0\|_{L^2}^2\right)(W_R, |\psi_0|^2) < m_0, \tag{5.13}$$

where K is the constant given in (A.2). We remark that Lemma A.2 shows that we can take R satisfying (5.13).

Now, for $R > 0$ satisfying (5.10)–(5.13), we put

$$T_R = \sup\left\{ T>0 \;\middle|\; \sup_{0\le t<T}\int_{|x|\ge R}|\psi(x,t)|^2\,dx \le m_0 \right\}. \tag{5.14}$$

Note here that since $\psi \in C(\mathbb{R}; L^2(\mathbb{R}^d))$ and $\psi(0) = \psi_0$, we have by (5.12) that $T_R > 0$. It is clear that $T_R = \infty$ shows (5.9).

We suppose the contrary that $T_R < \infty$. Then, it follows from $\psi \in C(\mathbb{R}; L^2(\mathbb{R}^d))$ that

$$\int_{|x|\ge R}|\psi(x,T_R)|^2\,dx = m_0. \tag{5.15}$$

Hence, the definition of m_0 (see (5.2)), together with (5.1), (5.10) and the mass conservation law (1.5), leads us to that

$$-\frac{1}{4}\varepsilon_0 \le \mathcal{K}^R(\psi(T_R)). \tag{5.16}$$

Moreover, applying this inequality (5.16) and (1.52) (or (2.41)) to the generalized virial identity (A.20), we obtain that

$$\begin{aligned}(W_R, |\psi(T_R)|^2) < (W_R, |\psi_0|^2) + 2T_R \Im(\vec{w}_R \cdot \nabla\psi_0, \psi_0) - \varepsilon_0^2 T_R^2 \\ + \frac{1}{4}\varepsilon_0 T_R^2 - \frac{1}{2}\int_0^{T_R}\int_0^{t'} (\Delta(\operatorname{div}\vec{w}_R), |\psi(t'')|^2)\, dt''dt'.\end{aligned} \tag{5.17}$$

Here, the estimate (A.9), the mass conservation law (1.5) and (5.11) show that the last term on the right-hand side above is estimated as follows:

$$\begin{aligned}-\frac{1}{2}\int_0^{T_R}\int_0^{t'} (\Delta(\operatorname{div}\vec{w}_R), |\psi(t'')|^2)\, dt''dt' &\le \frac{1}{2}\int_0^{T_R}\int_0^{t'} \frac{10d^2K}{R^2}\left\|\psi(t'')\right\|_{L^2}^2\, dt''dt' \\ &\le \frac{1}{4}\varepsilon_0 T_R^2.\end{aligned} \tag{5.18}$$

Therefore, we have

$$\begin{aligned}(W_R, |\psi(T_R)|^2) &< (W_R, |\psi_0|^2) + 2T_R \Im(\vec{w}_R \cdot \nabla\psi_0, \psi_0) - \frac{1}{2}T_R^2\varepsilon_0. \\ &= (W_R, |\psi_0|^2) - \frac{1}{2}\varepsilon_0\left\{T_R - \frac{2}{\varepsilon_0}\Im(\vec{w}_R\cdot\nabla\psi_0, \psi_0)\right\}^2 + \frac{2}{\varepsilon_0}\left|(\vec{w}_R\cdot\nabla\psi_0, \psi_0)\right|^2 \\ &\le (W_R, |\psi_0|^2) + \frac{2}{\varepsilon_0}\left|(\vec{w}_R\cdot\nabla\psi_0, \psi_0)\right|^2 \\ &\le (W_R, |\psi_0|^2) + \frac{2}{\varepsilon_0}\left\|\nabla\psi_0\right\|_{L^2}^2 \int_{\mathbb{R}^d} W_R(x)|\psi_0(x)|^2\, dx,\end{aligned} \tag{5.19}$$

where we have used the Schwarz inequality and (A.5) to derive the last inequality. Combining this inequality (5.19) with (5.13), we see that

$$(W_R, |\psi(T_R)|^2) \le \left(1 + \frac{2}{\varepsilon_0}\left\|\nabla\psi_0\right\|_{L^2}^2\right)(W_R, |\psi_0|^2) < R^2 m_0. \tag{5.20}$$

On the other hand, since $W_R(x) \ge R^2$ for $|x| \ge R$ (see (A.4)), we have

$$\int_{|x|\ge R} |\psi(x, T_R)|^2\, dx = \frac{1}{R^2}\int_{|x|\ge R} R^2|\psi(x, T_R)|^2\, dx \le \frac{1}{R^2}(W_R, |\psi(T_R)|^2). \tag{5.21}$$

Thus, it follows form (5.20) and (5.21) that

$$\int_{|x|\ge R} |\psi(x, T_R)|^2\, dx < m_0, \tag{5.22}$$

which contradicts (5.15), so that $T_R = \infty$ and (5.9) hold.

Step3. We complete the proof of Theorem 1.2. The definition of m_0, together with the mass conservation law (1.5), (5.1) and (5.9), shows that

$$-\frac{1}{4}\varepsilon_0 \le \mathcal{K}^R(\psi(t)) \qquad \text{for all } R > 0 \text{ satisfying (5.10)–(5.13), and all } t \ge 0. \tag{5.23}$$

Applying this estimate (5.23) to the generalized virial identity (A.20), we obtain the following estimate as well as *Step2* (see (5.17) and (5.18)):

$$(W_R, |\psi(t)|^2) \le (W_R, |\psi_0|^2) + 2t\Im(\vec{w}_R \cdot \nabla\psi_0, \psi_0) - \frac{1}{2}t^2\varepsilon_0 \quad \text{for all } t \ge 0. \tag{5.24}$$

This inequality means that $(W_R, |\psi(t)|^2)$ becomes negative in a finite time, so that $T^+_{\max}$ must be finite. However, this contradicts $T^+_{\max} = \infty$. Hence, (5.1) derives an absurd conclusion: Thus, (1.54) holds. ☐

Next, we give the proof of Theorem 1.6.

Proof of Theorem 1.6. Let ψ_0 be a radially symmetric function in PW_-, and let ψ be the solution to (1.1) with $\psi(0) = \psi_0$.

To handle the term containing $\mathcal{K}^R$ in the generalized virial identity (A.20) for ψ, we consider the following variational problem, as well as Lemma 3.3 in [46]:

Lemma 5.1. *Assume that $d \ge 3$, $2 + \frac{4}{d} < p + 1 < 2^*$, and $p \le 5$ if $d = 2$. Then, there exists $R_* > 0$ and $m_* > 0$ such that*

$$m_* < \inf \left\{ \int_{|x| \ge R} |v(x)|^2\, dx \,\middle|\, \begin{array}{l} v \in H^1_{rad}(\mathbb{R}^d),\ \mathcal{K}^R(v) \le -\frac{1}{4}\varepsilon_0, \\ \|v\|_{L^2} \le \|\psi_0\|_{L^2} \end{array} \right\} \quad \text{for all } R \ge R_*,$$

where $H^1_{rad}(\mathbb{R}^d)$ is the set of radially symmetric functions in $H^1(\mathbb{R}^d)$, and $\varepsilon_0 = \mathcal{H}(\psi_0) - \mathcal{B}(\psi_0) > 0$.

Proof of Lemma 5.1. We take a function $v \in H^1_{rad}(\mathbb{R}^d)$ with the following properties:

$$\mathcal{K}^R(v) \le -\frac{1}{4}\varepsilon_0, \tag{5.25}$$

$$\|v\|_{L^2} \le \|\psi_0\|_{L^2}, \tag{5.26}$$

where R is a sufficiently large constant to be chosen later (see (5.32)). Since $v \in H^1_{rad}(\mathbb{R}^d)$, $\mathcal{K}^R(v)$ is written as follows (see Remark A.1):

$$\mathcal{K}^R(v) = \int_{\mathbb{R}^d} \rho_0(x)|\nabla v(x)|^2\, dx - \int_{\mathbb{R}^d} \rho_3(x)|v(x)|^{p+1}\, dx. \tag{5.27}$$

Hence, it follows from (5.25) that

$$\begin{aligned} \frac{1}{4}\varepsilon_0 + \int_{\mathbb{R}^d} \rho_0(x)|\nabla v(x)|^2\, dx &\le -\mathcal{K}^R(v) + \int_{\mathbb{R}^d} \rho_0(x)|\nabla v(x)|^2\, dx \\ &= \int_{\mathbb{R}^d} \rho_3(x)|v(x)|^{p+1}\, dx. \end{aligned} \tag{5.28}$$

To estimate the right-hand side of (5.28), we employ the following inequality (see Lemma 6.5.11 in [12]): Assume that $d \ge 1$. Let κ be a non-negative and radially symmetric function in $C^1(\mathbb{R}^d)$ with $|x|^{-(d-1)} \max\{-\frac{x\nabla\kappa}{|x|}, 0\} \in L^\infty(\mathbb{R}^d)$. Then, we have that

$$\begin{aligned} &\|\kappa^{\frac{1}{2}} f\|_{L^\infty} \\ &\le K_5 \|f\|^{\frac{1}{2}}_{L^2} \left\{ \left\| |x|^{-(d-1)} \kappa \frac{x\nabla f}{|x|} \right\|^{\frac{1}{2}}_{L^2} + \left\| |x|^{-(d-1)} \max\left\{ -\frac{x\nabla\kappa}{|x|},\ 0 \right\} \right\|^{\frac{1}{2}}_{L^\infty} \|f\|^{\frac{1}{2}}_{L^2} \right\} \\ &\qquad\qquad \text{for all } f \in H^1_{rad}(\mathbb{R}^d), \end{aligned} \tag{5.29}$$

where $K_5 > 0$ is some constant depending only on d.
This inequality (5.29), together with (A.31), (A.32) and $\operatorname{supp} \rho_3 = \{|x| \geq R\}$ (see (A.27)), yields the following estimate for the right-hand side of (5.28):

$$
\begin{aligned}
&\int_{\mathbb{R}^d} \rho_3(x)|v(x)|^{p+1}\,dx \\
&\leq \|\rho_3^{\frac{1}{4}} v\|_{L^\infty}^{p-1} \int_{\mathbb{R}^d} \rho_3^{\frac{5-p}{4}}(x)|v(x)|^2\,dx \\
&\leq \frac{K_5^{p-1}}{R^{\frac{p-1}{2}(d-1)}} \|v\|_{L^2(|x|\geq R)}^{\frac{p-1}{2}} \left\{ \left\| \sqrt{\rho_3}\frac{x\nabla v}{|x|} \right\|_{L^2}^{\frac{1}{2}} + \left\| \max\left\{ -\frac{x\nabla\sqrt{\rho_3}}{|x|},\ 0 \right\} \right\|_{L^\infty}^{\frac{1}{2}} \|v\|_{L^2}^{\frac{1}{2}} \right\}^{p-1} K_3^{\frac{5-p}{4}} \|v\|_{L^2}^2 \\
&\leq \frac{K_5^{p-1} K_3^{\frac{5-p}{4}}}{R^{\frac{p-1}{2}(d-1)}} \|v\|_{L^2(|x|\geq R)}^{\frac{p-1}{2}} \left\{ \|\sqrt{\rho_3}\nabla v\|_{L^2}^{\frac{1}{2}} + \frac{(K_3')^{\frac{1}{2}}}{R^{\frac{1}{2}}} \|v\|_{L^2}^{\frac{1}{2}} \right\}^{p-1} \|v\|_{L^2}^2 \\
&\leq \frac{K_5^{p-1} K_3^{\frac{5-p}{4}}}{R^{\frac{p-1}{2}(d-1)}} \|v\|_{L^2(|x|\geq R)}^{\frac{p-1}{2}} C_p \left\{ \|\sqrt{\rho_3}\nabla v\|_{L^2}^{\frac{p-1}{2}} + \frac{(K_3')^{\frac{p-1}{2}}}{R^{\frac{p-1}{2}}} \|v\|_{L^2}^{\frac{p-1}{2}} \right\} \|v\|_{L^2}^2
\end{aligned}
\tag{5.30}
$$

for some constant $C_p > 0$ depending only on p. Moreover, using $p \leq 5$ (so that $\frac{p-1}{2} \leq 2$), $\|v\|_{L^2} \leq \|\psi_0\|_{L^2}$, and the Young inequality ($ab \leq \frac{a^r}{r} + \frac{b^{r'}}{r'}$ with $\frac{1}{r} + \frac{1}{r'} = 1$), we obtain that

$$
\begin{aligned}
\text{R.H.S. of (5.30)} \leq \frac{p-1}{4} \|v\|_{L^2(|x|\geq R)}^2 \|\sqrt{\rho_3}\nabla v\|_{L^2}^2 + \frac{5-p}{4} \left(\frac{C_p \|\psi_0\|_{L^2}^2 K_5^{p-1} K_3^{\frac{5-p}{4}}}{R^{\frac{p-1}{2}(d-1)}} \right)^{\frac{4}{5-p}} \\
+ C_p \frac{\|\psi_0\|_{L^2}^{p+1} K_5^{p-1} K_3^{\frac{5-p}{4}} (K_3')^{\frac{p-1}{2}}}{R^{\frac{d(p-1)}{2}}}.
\end{aligned}
\tag{5.31}
$$

If R is so large that

$$
\frac{5-p}{4} \left(\frac{C_p \|\psi_0\|_{L^2}^2 K_5^{p-1} K_3^{\frac{5-p}{4}}}{R^{\frac{p-1}{2}(d-1)}} \right)^{\frac{4}{5-p}} + \frac{C_p \|\psi_0\|_{L^2}^{p+1} K_5^{p-1} K_3^{\frac{5-p}{4}} (K_3')^{\frac{p-1}{2}}}{R^{\frac{d(p-1)}{2}}} \leq \frac{1}{8}\varepsilon_0,
\tag{5.32}
$$

then (5.28), together with (5.30) and (5.31), shows that

$$
\int_{\mathbb{R}^d} \left\{ \rho_0(x) - \frac{p-1}{4} \|v\|_{L^2(|x|\geq R)}^2 \rho_3(x) \right\} |\nabla v(x)|^2 \leq -\frac{\varepsilon_0}{8}.
\tag{5.33}
$$

Hence, it must hold that

$$
\inf_{|x|\geq R} \frac{\rho_0(x)}{\rho_3(x)} < \frac{p-1}{4} \|v\|_{L^2(|x|\geq R)}^2,
\tag{5.34}
$$

which, together with (A.33), gives us the desired result. □

The following lemma tells us that the solution ψ is well localized:

Lemma 5.2 (cf. Lemma 3.4 in [46])**.** *Assume that* $d \geq 1$, $2 + \frac{4}{d} < p + 1 < 2^*$, *and* $p \leq 5$ *if* $d = 2$. *Let* R_* *and* m_* *be constants found in Lemma 5.1. Then, we have*

$$T_{\max}^{+} = \sup \left\{ T > 0 \;\middle|\; \sup_{0 \leq t < T} \int_{|x| \geq R} |\psi(x,t)|^2 \, dx < m_* \right\}$$

for all R *with the following properties:*

$$R > R_*, \tag{5.35}$$

$$\frac{10 d^2 K}{R^2} \|\psi_0\|_{L^2}^2 < \varepsilon_0, \tag{5.36}$$

$$\int_{|x| \geq R} |\psi_0(x)|^2 \, dx < m_*, \tag{5.37}$$

$$\frac{1}{R^2} \left(1 + \frac{2}{\varepsilon_0} \|\nabla \psi_0\|_{L^2}^2 \right) (W_R, |\psi_0|^2) < m_*. \tag{5.38}$$

Proof of Lemma 5.2. For $R > 0$ satisfying (5.35)–(5.38), we put

$$T_R := \sup \left\{ T > 0 \;\middle|\; \sup_{0 \leq t \leq T} \int_{|x| \geq R} |\psi(x,t)|^2 \, dx < m_* \right\}.$$

We suppose the contrary that $T_R < T_{\max}^{+}$, so that it follows from $\psi \in C([0, T_{\max}^{+}); L^2(\mathbb{R}^d))$ that

$$\int_{|x| \geq R} |\psi(x, T_R)|^2 \, dx = m_*. \tag{5.39}$$

Hence, we obtain by Lemma 5.1 that

$$-\frac{1}{4} \varepsilon_0 < \mathcal{K}^R(\psi(T_R)). \tag{5.40}$$

Then, the same computation as (5.17)–(5.22) yields that

$$\int_{|x| \geq R} |\psi(T_R, x)|^2 \, dx < m_*, \tag{5.41}$$

which contradicts (5.39): Thus, $T_R = T_{\max}^{+}$. □

Now, we are in a position to complete the proof of Theorem 1.6.

Proof of Theorem 1.6 (continued) We take $R > 0$ satisfying (5.35)–(5.38). Then, Lemma 5.2 gives us that

$$\int_{|x| \geq R} |\psi(t,x)|^2 \, dx < m_* \quad \text{for all } t \in [0, T_{\max}^{+}). \tag{5.42}$$

Moreover, Lemma 5.1, together with (5.42), shows that

$$-\frac{\varepsilon_0}{4} < \mathcal{K}^R(\psi(t)) \quad \text{for all } t \in [0, T_{\max}^{+}). \tag{5.43}$$

Applying (5.43) and (1.52) to the generalized virial identity (A.20), we obtain that

$$\begin{aligned}(W_R,|\psi(t)|^2) &< (W_R,|\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) - \varepsilon_0^2 t^2 \\ &\quad + \frac{1}{4}\varepsilon_0 t^2 - \frac{1}{2}\int_0^t\int_0^{t'}(\Delta(\operatorname{div}\vec{w}_R),|\psi(t'')|^2)\,dt''dt'.\end{aligned} \tag{5.44}$$

Here, it follows from (A.9), the mass conservation law (1.5) and (5.36) that the last term on the right-hand side above is estimated as follows :

$$\begin{aligned}-\frac{1}{2}\int_0^t\int_0^{t'}(\Delta(\operatorname{div}\vec{w}_R),|\psi(t'')|^2)\,dt''dt' &\le \frac{1}{2}\int_0^t\int_0^{t'}\frac{10d^2K}{R^2}\left\|\psi(t'')\right\|_{L^2}^2\,dt''dt' \\ &\le \frac{1}{4}\varepsilon_0 t^2.\end{aligned} \tag{5.45}$$

Combining (5.44) with (5.45), we obtain that

$$(W_R,|\psi(t)|^2) < (W_R,|\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) - \frac{1}{2}t^2\varepsilon_0. \tag{5.46}$$

This inequality means that $(W_R,|\psi(t)|^2)$ becomes negative in a finite time, so that $T_{\max}^+ < \infty$: This fact, together with (1.4), shows (1.59).

Now, we shall show the claims (i)–(iii) in Theorem 1.6.

We can find that Lemma 5.2 is still valid if we replace m_* with any $m \in (0,m_*)$. Thus, we see that (i) holds.

We next consider (ii). Since we have (A.29), in order to prove (1.61), it suffices to show that

$$\int_0^{T_{\max}^+}(T_{\max}^+-t)\left(\int_{\mathbb{R}^d}\rho_0(x)|\nabla\psi(x,t)|^2\,dx\right)dt < \infty \quad \text{for all sufficiently large } R>0. \tag{5.47}$$

Let us prove this. Integrating by parts, we find that

$$\begin{aligned}&\int_0^t(t-t')\left(\int_{\mathbb{R}^d}\rho_0(x)|\nabla\psi(x,t')|^2\,dx\right)dt' \\ &= \int_0^t\int_0^{t'}\left(\int_{\mathbb{R}^d}\rho_0(x)|\nabla\psi(x,t'')|^2\,dx\right)dt''\,dt' \quad \text{for all } t\ge 0.\end{aligned} \tag{5.48}$$

This formula (5.48) and the generalized virial identity (A.20) lead us to the following estimate:

$$\begin{aligned}&2\int_0^t(t-t')\left(\int_{\mathbb{R}^d}\rho_0(x)|\nabla\psi(x,t')|^2\,dx\right)dt' \\ &\le (W_R,|\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) + 2\int_0^t\int_0^{t'}\mathcal{K}(\psi(t''))\,dt''dt' \\ &\quad + 2\int_0^t(t-t')\left(\int_{\mathbb{R}^d}\rho_3(x)|\psi(x,t')|^{p+1}\,dx\right)dt' - \frac{1}{2}\int_0^t\int_0^{t'}(\Delta(\operatorname{div}\vec{w}_R),|\psi(t'')|^2)\,dt''dt'.\end{aligned} \tag{5.49}$$

Here, applying the estimate (5.45) to the right-hand side of (5.49) (abbreviated to R.H.S. of (5.49)), we have that

$$\begin{aligned}\text{R.H.S. of (5.49)} &\le (W_R, |\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) - \varepsilon_0 t^2 \\ &\qquad + 2\int_0^t (t-t')\left(\int_{\mathbb{R}^d}\rho_3(x)|\psi(x,t')|^{p+1}\,dx\right)dt' + \frac{\varepsilon_0}{4}t^2.\end{aligned} \tag{5.50}$$

Moreover, taking R so large that (5.32) holds, we find by the same computation as (5.30)–(5.31) that

$$\int_{\mathbb{R}^d}\rho_3(x)|\psi(x,t)|^{p+1}\,dx \le \frac{p-1}{4}\,\|\psi(t)\|^2_{L^2(|x|\ge R)}\int_{\mathbb{R}^d}\rho_3(x)|\nabla\psi(x,t)|^2\,dx + \frac{\varepsilon_0}{8}. \tag{5.51}$$

Combining (5.49), (5.50) and (5.51), we obtain that

$$\begin{aligned}&2\int_0^t (t-t')\left(\int_{\mathbb{R}^d}\left\{\rho_0(x) - \frac{p-1}{4}\,\|\psi(t)\|^2_{L^2(|x|\ge R)}\,\rho_3(x)\right\}|\nabla\psi(x,t)|^2\,dx\right)dt' \\ &\le (W_R,|\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) - \frac{\varepsilon_0}{2}t^2 \\ &\qquad\qquad\qquad \text{for all } R>0 \text{ satisfying (5.32) and (5.35)–(5.38).}\end{aligned} \tag{5.52}$$

Here, using the result of (i) (the formula (1.60)), we can take a constant $R_1 > 0$ such that

$$\frac{p-1}{4}\,\|\psi(t)\|^2_{L^2(|x|\ge R_1)}\,\frac{1}{K_4} \le \frac{1}{2} \quad \text{for all } t\in I_{\max}, \tag{5.53}$$

where K_4 is some positive constant found in (A.33). Therefore, (5.52), together with (5.53) and (A.33), shows that

$$\begin{aligned}&\int_0^t (t-t')\left(\int_{\mathbb{R}^d}\rho_0(x)|\nabla\psi(x,t)|^2\,dx\right)dt' \\ &\le (W_R,|\psi_0|^2) + 2t\Im(\vec{w}_R\cdot\nabla\psi_0,\psi_0) - \frac{\varepsilon_0}{2}t^2 \quad \text{for all sufficiently large } R \text{ and } t\in[0,T^+_{\max}).\end{aligned} \tag{5.54}$$

This, together with (A.28) and (A.29), gives (1.61).

Once (1.61) is proved, we can easily obtain (1.62). Indeed, (5.51), together with (1.61) and (A.30), immediately yields (1.62). Thus, we have proved (ii).

Finally, we prove (iii). We take $R > 0$ so large that (1.61) holds valid.

To prove (1.63), we employ the radial interpolation inequality by Strauss [51]:

$$\|f\|^2_{L^\infty} \lesssim \frac{1}{R^{d-1}}\,\|f\|_{L^2}\,\|\nabla f\|_{L^2} \quad \text{for all } f\in H^1_{rad}(\mathbb{R}^d), \tag{5.55}$$

where the implicit constant depends only on d.
This inequality (5.55), together with (1.61) and the mass conservation law (1.5), gives us (1.63).

The estimate (1.64) follows from (1.63) and the inequality

$$\|\psi(t)\|^{p+1}_{L^{p+1}(|x|>R)} \le \|\psi_0\|^2_{L^2}\|\psi(t)\|^{p-1}_{L^\infty(|x|>R)}.$$

We obtain (1.65) and (1.66) from (1.63) and (1.64), respectively. □

We also have variants of (1.63)–(1.66). For example: in their proofs, replacing R by $R(T_{\max}^{+}-t)^{-\frac{1}{2(d-1)}}$, we obtain

$$\int_0^{T_{\max}^{+}} \|\psi(t)\|^4_{L^\infty(|x|>R(T_{\max}^{+}-t)^{-\frac{1}{2(d-1)}})}\,dt<\infty$$

and

$$\int_0^{T_{\max}^{+}} \|\psi(t)\|^{\frac{4(p+1)}{p-1}}_{L^{p+1}(|x|>R(T_{\max}^{+}-t)^{-\frac{1}{2(d-1)}})}\,dt<\infty.$$

Moreover, replacing R by $R(T_{\max}^{+}-t)^{-\frac{1}{d-1}}$ in (1.63) and (1.64), we obtain

$$\liminf_{t\to T_{\max}^{+}} \|\psi(t)\|^2_{L^\infty\left(|x|>R(T_{\max}^{+}-t)^{-\frac{1}{d-1}}\right)}=0$$

and

$$\liminf_{t\to T_{\max}^{+}} \|\psi(t)\|^{p+1}_{L^{p+1}\left(|x|>R(T_{\max}^{+}-t)^{-\frac{1}{d-1}}\right)}=0,$$

respectively.

Now, we shall give the proof of Propositions 1.3.

Proof of Proposition 1.3. Suppose that ψ is a solution to the equation (1.1) satisfying that

$$\limsup_{t\to T_{\max}^{+}} \|\nabla\psi(t)\|_{L^2}=\limsup_{t\to T_{\max}^{+}} \|\psi(t)\|_{L^{p+1}}=\infty. \tag{5.56}$$

Then, there exists a sequence $\{t_n\}_{n\in\mathbb{N}}$ in $[0,T_{\max}^{+})$ such that

$$\lim_{n\to\infty} t_n=T_{\max}^{+}, \tag{5.57}$$

$$\|\psi(t_n)\|_{L^{p+1}}=\sup_{t\in[0,t_n)} \|\psi(t)\|_{L^{p+1}}. \tag{5.58}$$

Using such a sequence $\{t_n\}_{n\in\mathbb{N}}$, we define a number λ_n by

$$\lambda_n=\|\psi(t_n)\|_{L^{p+1}}^{-\frac{(p-1)(p+1)}{d+2-(d-2)p}},\quad n\in\mathbb{N}. \tag{5.59}$$

It is easy to see that

$$\lim_{n\to\infty}\lambda_n=0. \tag{5.60}$$

We consider the scaled functions ψ_n defined by

$$\psi_n(x,t):=\lambda_n^{\frac{2}{p-1}}\overline{\psi(\lambda_n x,t_n-\lambda_n^2 t)},\quad (x,t)\in\mathbb{R}^d\times\left(-\frac{T_{\max}^{+}-t_n}{\lambda_n^2},\frac{t_n}{\lambda_n^2}\right],\quad n\in\mathbb{N}. \tag{5.61}$$

We can easily verify that

$$2i\frac{\partial \psi_n}{\partial t} + \Delta \psi_n + |\psi_n|^{p-1}\psi_n = 0 \quad \text{in } \mathbb{R}^d \times \left(-\frac{T_{\max}^{+} - t_n}{\lambda_n^2},\, \frac{t_n}{\lambda_n^2}\right], \tag{5.62}$$

$$\|\psi_n(t)\|_{L^2} = \lambda_n^{-\frac{d(p-1)-4}{2(p-1)}} \|\psi_0\|_{L^2} \quad \text{for all } t \in \left(-\tfrac{T_{\max}^{+}-t_n}{\lambda_n^2},\, \tfrac{t_n}{\lambda_n^2}\right], \tag{5.63}$$

$$\mathcal{H}(\psi_n(t)) = \lambda_n^{\frac{2(p+1)}{p-1}-d}\mathcal{H}(\psi_0) \quad \text{for all } t \in \left(-\tfrac{T_{\max}^{+}-t_n}{\lambda_n^2},\, \tfrac{t_n}{\lambda_n^2}\right], \tag{5.64}$$

$$\sup_{t\in\left[0,\, \frac{t_n}{\lambda_n^2}\right]} \|\psi_n(t)\|_{L^{p+1}} = 1, \tag{5.65}$$

where we put $\psi_0 = \psi(0)$. Besides, we have that

$$\sup_{t\in\left[0, \frac{t_n}{\lambda_n^2}\right]} \|\nabla\psi_n(t)\|_{L^2}^2 \le 1 \quad \text{for all sufficiently large } n \in \mathbb{N}. \tag{5.66}$$

Indeed, (5.64) and (5.65) lead us to that

$$\begin{aligned} \|\nabla\psi_n(t)\|_{L^2}^2 &= \mathcal{H}(\psi_n(t)) + \frac{2}{p+1}\|\psi_n(t)\|_{L^{p+1}}^{p+1} \\ &\le \lambda_n^{\frac{2(p+1)}{p-1}-d}\mathcal{H}(\psi_0) + \frac{2}{p+1} \quad \text{for all } t \in \left[0,\, \tfrac{t_n}{\lambda_n^2}\right], \end{aligned} \tag{5.67}$$

which, together with (5.60), immediately yields (5.66).

Now, we suppose that

$$\|\psi\|_{L^\infty([0,T_{\max}^{+});L^{\frac{d}{2}(p-1)})} < \infty, \tag{5.68}$$

so that $\{\psi_n\}$ satisfies that

$$\sup_{n\in\mathbb{N}} \|\psi_n\|_{L^\infty([0,\frac{t_n}{\lambda_n^2}];L^{\frac{d}{2}(p-1)})} \le \|\psi\|_{L^\infty([0,T_{\max}^{+});L^{\frac{d}{2}(p-1)})} < \infty. \tag{5.69}$$

Take any $T > 0$. Then, extraction of some subsequence of $\{\psi_n\}$ allows us to assume that:

$$\sup_{t\in[0,T]} \|\psi_n(t)\|_{L^{\frac{d}{2}(p-1)}} \lesssim 1, \quad \sup_{t\in[0,T]} \|\psi_n(t)\|_{L^{p+1}} = 1, \quad \sup_{t\in[0,T]} \|\nabla\psi_n(t)\|_{L^2} \le 1 \quad \text{for all } n \in \mathbb{N}, \tag{5.70}$$

where the implicit constant is independent of n. The condition (5.70) enable us to apply Lemmata C.1 and C.2, so that: There exist a constant $\delta > 0$ and a sequence $\{y_n\}_{n\in\mathbb{N}}$ in $\mathbb{R}^d$ such that, putting $\widetilde{\psi}_n(x,t) = \psi_n(x + y_n, t)$, we have

$$\sup_{t\in[0,T]} \mathcal{L}^d\left(\left[\left|\widetilde{\psi}_n(\cdot + y_n, t)\right| \ge \delta\right] \cap B_1(0)\right) \gtrsim 1 \quad \text{for all } n \in \mathbb{N}, \tag{5.71}$$

where $B_1(0) = \{x \in \mathbb{R}^d | |x| < 1\}$ and the implicit constant is independent of n. Besides, we have by (5.62) and (5.70) that

$$2i\frac{\partial \widetilde{\psi}_n}{\partial t} + \Delta\widetilde{\psi}_n + |\widetilde{\psi}_n|^{p-1}\widetilde{\psi}_n = 0 \quad \text{in } \mathbb{R}^d \times [0,T], \tag{5.72}$$

$$\widetilde{\psi}_n \in C([0,T]; \dot{H}^1(\mathbb{R}^d) \cap L^{p+1}(\mathbb{R}^d)) \quad \text{for all } n \in \mathbb{N}, \tag{5.73}$$

$$\sup_{t\in[0,T]} \left\|\widetilde{\psi}_n(t)\right\|_{L^{p+1}} = 1, \quad \sup_{t\in[0,T]} \left\|\nabla\widetilde{\psi}_n(t)\right\|_{L^2} \le 1 \quad \text{for all } n \in \mathbb{N}. \tag{5.74}$$

We shall prove that $\{\widetilde{\psi}_n\}$ is an equicontinuous sequence in $C([0,T];L^2(\Omega))$ for any compact set $\Omega\subset\mathbb{R}^d$. Let χ be a function in $C_c^\infty(\mathbb{R}^d)$ such that $\chi\equiv 1$ on Ω. Then, we can verify that

$$\begin{aligned}
&\left\|\chi\widetilde{\psi}_n(t)-\chi\widetilde{\psi}_n(s)\right\|_{L^2}^2\\
&=\int_s^t \frac{d}{dt'}\left\|\chi\widetilde{\psi}_n(t')-\chi\widetilde{\psi}_n(s)\right\|_{L^2}^2 dt'\\
&=2\Re\int_s^t\int_{\mathbb{R}^d}\overline{\chi(x)(\partial_t\widetilde{\psi}_n)(x,t')}\left\{\chi(x)\widetilde{\psi}_n(x,t')-\chi(x)\widetilde{\psi}_n(x,s)\right\}dxdt' \quad \text{for all } s,t\in[0,T].
\end{aligned} \tag{5.75}$$

This identity (5.75), together with (5.72), yields that

$$\begin{aligned}
&\left\|\widetilde{\psi}_n(t)-\widetilde{\psi}_n(s)\right\|_{L^2(\Omega)}^2\\
&\le\left\|\chi\widetilde{\psi}_n(t)-\chi\widetilde{\psi}_n(s)\right\|_{L^2}^2\\
&\le 2|t-s|\left\|\chi\partial_t\widetilde{\psi}_n\right\|_{L^\infty([0,T];H^{-1})}\left\|\chi\widetilde{\psi}_n\right\|_{L^\infty([0,T];H^1)}\\
&\le 2|t-s|\left\{\left\|\chi\Delta\widetilde{\psi}_n\right\|_{L^\infty([0,T];H^{-1})}+\left\|\chi(|\widetilde{\psi}_n|^{p-1}\widetilde{\psi}_n)\right\|_{L^\infty([0,T];H^{-1})}\right\}\\
&\qquad\times\left\{\left\|\chi\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}+\left\|(\nabla\chi)\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}+\left\|\chi\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}\right\}\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\text{for all } s,t\in[0,T].
\end{aligned} \tag{5.76}$$

We consider the terms in the first parentheses on the right-hand side of (5.76). They are estimated by the Sobolev embedding and the Hölder inequality as follows:

$$\begin{aligned}
&\left\|\chi\Delta\widetilde{\psi}_n\right\|_{L^\infty([0,T];H^{-1})}+\left\|\chi(|\widetilde{\psi}_n|^{p-1}\widetilde{\psi}_n)\right\|_{L^\infty([0,T];H^{-1})}\\
&=\left\|\Delta(\chi\widetilde{\psi}_n)-(\Delta\chi)\widetilde{\psi}_n-2\nabla\chi\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];H^{-1})}+\left\|\chi(|\widetilde{\psi}_n|^{p-1}\widetilde{\psi}_n)\right\|_{L^\infty([0,T];H^{-1})}\\
&\lesssim\left\|\nabla(\chi\widetilde{\psi}_n)\right\|_{L^\infty([0,T];L^2)}+\left\|(\Delta\chi)\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^{\frac{p+1}{p}})}+\left\|\nabla\chi\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}\\
&\qquad\quad+\left\|\chi(|\psi_n|^{p-1}\widetilde{\psi}_n)\right\|_{L^\infty([0,T];L^{\frac{p+1}{p}})}\\
&\le\|\nabla\chi\|_{L^{\frac{2(p+1)}{p-1}}}\left\|\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^{p+1})}+\|\chi\|_{L^\infty}\left\|\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}\\
&\qquad+\|\Delta\chi\|_{L^{\frac{p+1}{p-1}}}\left\|\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^{p+1})}+\|\nabla\chi\|_{L^\infty}\left\|\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}\\
&\qquad\quad+\|\chi\|_{L^\infty}\left\|\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^{p+1})}^{p+1},
\end{aligned} \tag{5.77}$$

where the implicit constant depends only on d and p. On the other hand, the terms in the second parentheses on the right-hand side of (5.76) are estimated by the Hölder inequality as follows:

$$\begin{aligned}&\left\|\chi\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}+\left\|(\nabla\chi)\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}+\left\|\chi\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}\\&\le 2\left\|\chi\right\|_{W^{1,\frac{2(p+1)}{p-1}}}\left\|\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^{p+1})}+\left\|\chi\right\|_{L^\infty}\left\|\nabla\widetilde{\psi}_n\right\|_{L^\infty([0,T];L^2)}.\end{aligned}\tag{5.78}$$

Hence, combining these estimates (5.76)–(5.78) with (5.74), we obtain that

$$\left\|\widetilde{\psi}_n(t)-\widetilde{\psi}_n(s)\right\|_{L^2(\Omega)}^2\lesssim|t-s|,\tag{5.79}$$

where the implicit constant depends only on d, p and Ω (χ is determined by Ω). Thus, we see that $\{\widetilde{\psi}_n\}$ is equicontinuous in $C([0,T];L^2(\Omega))$ for all compact set $\Omega\subset\mathbb{R}^d$.

The estimate (5.79), with the help of the Gagliardo-Nirenberg inequality and (5.74), also shows that $\{\widetilde{\psi}_n\}$ is an equicontinuous sequence in $C([0,T];L^q(\Omega))$ for all compact set $\Omega\subset\mathbb{R}^d$ and $q\in[2,2^*)$. Hence, (5.74) and the Ascoli-Arzéra theorem, together with (5.71) and (5.72), yield that: There exist a subsequence of $\{\widetilde{\psi}_n\}$ (still denoted by the same symbol) and a nontrivial function $\psi_\infty\in L^\infty([0,\infty);\dot{H}^1(\mathbb{R}^d)\cap L^{p+1}(\mathbb{R}^d))$ such that

$$\lim_{n\to\infty}\widetilde{\psi}_n=\psi_\infty\quad\text{strongly in }L^\infty([0,T];L^q_{loc}(\mathbb{R}^d))\quad\text{for all }q\in[2,2^*),\tag{5.80}$$

$$\lim_{n\to\infty}\nabla\widetilde{\psi}_n=\nabla\psi_\infty\quad\text{weakly* in }L^\infty([0,T];L^2(\mathbb{R}^d)),\tag{5.81}$$

$$2i\frac{\partial\psi_\infty}{\partial t}+\Delta\psi_\infty+|\psi_\infty|^{p-1}\psi_\infty=0\quad\text{in }\mathcal{D}'([0,\infty);\dot{H}^{-1}(\mathbb{R}^d)+L^{\frac{p+1}{p}}(\mathbb{R}^d)).\tag{5.82}$$

Finally, we hall prove the formula (1.72). The Lebesgue dominated convergence theorem leads to that for all $\varepsilon>0$, there exists $R>0$ such that

$$\int_{|x|\le R}|\psi_\infty(0)|^{\frac{d(p-1)}{2}}\,dx\ge(1-\varepsilon)\int_{\mathbb{R}^d}|\psi_\infty(0)|^{\frac{d(p-1)}{2}}\,dx.\tag{5.83}$$

Combining this estimate with the strong convergence (5.80), we have that

$$\lim_{n\to\infty}\int_{|x|\le R}|\widetilde{\psi}_n(x,0)|^{\frac{d(p-1)}{2}}\,dx\ge(1-\varepsilon)\int_{\mathbb{R}^d}|\psi_\infty(0)|^{\frac{d(p-1)}{2}}\,dx.\tag{5.84}$$

Hence, it follows from (5.84) and the relation

$$\int_{|x|\le R}|\widetilde{\psi}_n(x,0)|^{\frac{d(p-1)}{2}}\,dx=\int_{|x-\lambda_n y_n|\le\lambda_n R}|\psi(x,t_n)|^{\frac{d(p-1)}{2}}\,dx\tag{5.85}$$

that the desired result holds. □

Finally, we shall give the proof of Proposition 1.4.

Proof of Proposition 1.4. We use the same assumptions, definitions and notation in Proposition 1.3, except the condition (1.71).

We find from the proof of Proposition 1.3 that: For all $T > 0$, there exists a subsequence of $\{\psi_n\}$ in $C([0,T]; H^1(\mathbb{R}^d))$ (still denoted by the same symbol) with the following properties:

$$\sup_{t\in[0,T]} \|\psi_n(t)\|_{L^{p+1}} = 1 \quad \text{for all } n \in \mathbb{N}, \tag{5.86}$$

$$\sup_{t\in[0,T]} \|\nabla\psi_n(t)\|_{L^2} \le 1 \quad \text{for all } n \in \mathbb{N}, \tag{5.87}$$

$$2i\frac{\partial\psi_n}{\partial t} + \Delta\psi_n + |\psi_n|^{p-1}\psi_n = 0 \quad \text{in } \mathbb{R}^d \times [0,T]. \tag{5.88}$$

For such a subsequence $\{\psi_n\}$, we define renormalized functions Φ_n^{RN} by

$$\Phi_n^{RN}(x,t) = \psi_n(x,t) - e^{\frac{i}{2}t\Delta}\psi_n(x,0), \quad n \in \mathbb{N}. \tag{5.89}$$

Here, it is worth while noting that

$$\Phi_n^{RN} \in C([0,T]; H^1(\mathbb{R}^d)) \quad \text{for all } n \in \mathbb{N}, \tag{5.90}$$

$$\Phi_n^{RN}(t) = \frac{i}{2}\int_0^t e^{\frac{i}{2}(t-t')\Delta}|\psi_n(t')|^{p-1}\psi_n(t')\, dt' \quad \text{for all } n \in \mathbb{N}. \tag{5.91}$$

We shall show that

$$\sup_{n\in\mathbb{N}} \|\Phi_n^{RN}\|_{L^\infty([0,T];H^1)} \le C_T \tag{5.92}$$

for some constant $C_T > 0$ depending only on d, p and T. Applying the Strichartz estimate to the formula (5.91), and using (5.86), we obtain the following two estimates:

$$\begin{aligned}
\|\Phi_n^{RN}\|_{L^\infty([0,T];L^2)} &\lesssim \||\psi_n|^{p-1}\psi_n\|_{L^{\frac{4(p+1)}{4(p+1)-d(p-1)}}([0,T];L^{\frac{p+1}{p}})} \\
&\le T^{1-\frac{d(p-1)}{4(p+1)}} \|\psi_n\|^p_{L^\infty([0,T];L^{p+1})} \\
&\le T^{1-\frac{d(p-1)}{4(p+1)}},
\end{aligned} \tag{5.93}$$

$$\begin{aligned}
\|\nabla\Phi_n^{RN}\|_{L^\infty([0,T];L^2)} &\lesssim \left\|\nabla\left(|\psi_n|^{p-1}\psi_n\right)\right\|_{L^{\frac{4(p+1)}{4(p+1)-d(p-1)}}([0,T];L^{\frac{p+1}{p}})} \\
&\le T^{1-\frac{d(p-1)}{2(p+1)}} \|\psi_n\|^{p-1}_{L^\infty([0,T];L^{p+1})} \|\nabla\psi_n\|_{L^{\frac{4(p+1)}{d(p-1)}}([0,T];L^{p+1})} \\
&\le T^{1-\frac{d(p-1)}{2(p+1)}} \|\nabla\psi_n\|_{L^{\frac{4(p+1)}{d(p-1)}}([0,T];L^{p+1})},
\end{aligned} \tag{5.94}$$

where the implicit constants depend only on d and p. Therefore, for the desired estimate (5.92), it suffices to show that

$$\sup_{n\in\mathbb{N}} \|\nabla\psi_n\|_{L^{\frac{4(p+1)}{d(p-1)}}([0,T];L^{p+1})} \le D_T \tag{5.95}$$

for some constant $D_T > 0$ depending only on d, p and T. Here, note that the pair $(p+1,\ \frac{4(p+1)}{d(p-1)})$ is admissible. In order to prove (5.95), we introduce an admissible pair

(q,r) with $q=p+2$ if $d=1,2$ and $q=\frac{1}{2}(p+1+2^{*})$ if $d\ge 3$, so that $p+1<q<2^{*}$. Then, it follows from the integral equation for ψ_n and the Strichartz estimate that

$$\begin{aligned}
&\|\nabla\psi_n\|_{L^r([0,T];L^q)}\\
&\lesssim \|\nabla\psi_n(0)\|_{L^2}+\left\|\nabla\left(|\psi_n|^{p-1}\psi_n\right)\right\|_{L^{\frac{4(p+1)}{4(p+1)-d(p-1)}}([0,T];L^{\frac{p+1}{p}})}\\
&\lesssim \|\nabla\psi_n(0)\|_{L^2}+\|\psi_n\|^{p-1}_{L^{\frac{2(p+1)(p-1)}{d+2-(d-2)p}}([0,T];L^{p+1})}\|\nabla\psi_n\|_{L^{\frac{4(p+1)}{d(p-1)}}([0,T];L^{p+1})}\\
&\le \|\nabla\psi_n(0)\|_{L^2}+T^{\frac{d+2-(d-2)p}{2(p+1)}}\|\psi_n\|^{p-1}_{L^\infty([0,T];L^{p+1})}\|\nabla\psi_n\|^{1-\frac{q(p-1)}{(q-2)(p+1)}}_{L^\infty([0,T];L^2)}\|\nabla\psi_n\|^{\frac{q(p-1)}{(q-2)(p+1)}}_{L^r([0,T];L^q)},
\end{aligned}\tag{5.96}$$

where the implicit constants depend only on d and p. Combining (5.96) with (5.86) and (5.87), we obtain that

$$\|\nabla\psi_n\|_{L^r([0,T];L^q)}\lesssim 1+T^{\frac{d+2-(d-2)p}{2(p+1)}}\|\nabla\psi_n\|^{\frac{q(p-1)}{(q-2)(p+1)}}_{L^r([0,T];L^q)}, \tag{5.97}$$

where the implicit constant depends only on d and p. Since $0<\frac{q(p-1)}{(q-2)(p+1)}<1$, this estimate, together with the Young inequality ($ab\le\frac{1}{\gamma}a^{\gamma}+\frac{1}{\gamma'}b^{\gamma'}$ for $\gamma,\gamma'>1$ with $\frac{1}{\gamma}+\frac{1}{\gamma'}=1$), yields that

$$\|\nabla\psi_n\|_{L^r([0,T];L^q)}\lesssim 1+T^{\frac{(q-2)\{d+2-(d-2)p\}}{4\{q-(p+1)\}}}, \tag{5.98}$$

where the implicit constant depends only on d and p. Hence, interpolating (5.87) and (5.98), we obtain (5.95), so that (5.92) holds.

Next, we shall show that $\{\Phi_n^{RN}\}$ is an equicontinuous sequence in $C([0,T];L^q(\mathbb{R}^d))$ for all $q\in[2,2^{*})$. Differentiating the both sides of (5.91), we obtain that

$$\begin{aligned}
\partial_t\Phi_n^{RN}(t)&=\frac{i}{2}|\psi_n(t)|^{p-1}\psi_n(t)-\frac{1}{4}\Delta\int_0^t e^{\frac{i}{2}(t-t')\Delta}|\psi_n(t')|^{p-1}\psi_n(t')\,dt'\\
&=\frac{i}{2}|\psi_n(t)|^{p-1}\psi_n(t)+\frac{i}{2}\Delta\Phi_n^{RN}(t).
\end{aligned}\tag{5.99}$$

This formula (5.99) and the Hölder inequality show that

$$\begin{aligned}
&\left\|\Phi_n^{RN}(t)-\Phi_n^{RN}(s)\right\|_{L^2}^2\\
&=\int_s^t\frac{d}{dt'}\left\|\Phi_n^{RN}(t')-\Phi_n^{RN}(s)\right\|_{L^2}^2dt'\\
&=2\Re\int_s^t\int_{\mathbb{R}^d}\overline{\partial_t\Phi_n^{RN}(x,t')}\left\{\Phi_n^{RN}(x,t')-\Phi_n^{RN}(x,s)\right\}dxdt'\\
&\lesssim |t-s|\,\|\psi_n\|^p_{L^\infty([0,T];L^{p+1})}\left\|\Phi_n^{RN}\right\|_{L^\infty([0,T];L^{p+1})}+|t-s|\left\|\nabla\Phi_n^{RN}\right\|^2_{L^\infty([0,T];L^2)},
\end{aligned}\tag{5.100}$$

where the implicit constant depends only on d and q. Combining this estimate with (5.86) and (5.92), we obtain

$$\left\|\Phi_n^{RN}(t)-\Phi_n^{RN}(s)\right\|_{L^2}^2\lesssim|t-s|\quad\text{for all } s,t\in[0,T], \tag{5.101}$$

where the implicit constant depends only on d, q and T. Moreover, the Gagliardo-Nirenberg inequality, together with (5.92) and (5.101), shows that $\{\Phi_n^{RN}\}$ is an equicontinuous sequence in $C([0,T]; L^q(\mathbb{R}^d))$ for all $q \in [2, 2^*)$.

Now, we are in a position to prove the main property of $\{\Phi_n^{RN}\}$, the alternatives (i) and (ii). We first suppose that

$$\lim_{n\to\infty} \sup_{t\in[0,T]} \left\|\Phi_n^{RN}(t)\right\|_{L^{\frac{d}{2}(p-1)}} = 0. \tag{5.102}$$

Then, a direct calculation immediately yields (1.77).

On the other hand, when

$$A := \limsup_{n\to\infty} \sup_{t\in[0,T]} \left\|\Phi_n^{RN}(t)\right\|_{L^{p+1}} > 0, \tag{5.103}$$

we can take a subsequence $\{\Phi_n^{RN}\}$ (still denoted by the same symbol) such that

$$\sup_{t\in[0,T]} \left\|\Phi_n^{RN}(t)\right\|_{L^{p+1}} \ge \frac{A}{2} \qquad \text{for all } n \in \mathbb{N}. \tag{5.104}$$

This inequality (5.104) and the uniform bound (5.92) enable us to apply Lemmata C.1 and C.2. Thus, we find that there exist a constant $\delta > 0$ and a sequence $\{y_n\}_{n\in\mathbb{N}}$ in $\mathbb{R}^d$ such that, putting $\widetilde{\Phi}_n^{RN}(x,t) = \Phi_n^{RN}(x + y_n, t)$, we have

$$\sup_{t\in[0,T]} \mathcal{L}^d\left(\left[\left|\widetilde{\Phi}_n^{RN}(t)\right| > \frac{\delta}{2}\right] \cap B_1(0)\right) > C \tag{5.105}$$

for some constant $C > 0$ being independent of n. Besides, we find by (5.92) and (5.101) that: $\{\widetilde{\Phi}_n^{RN}\}$ is

$$\text{a uniformly bounded sequence in } C([0,T]; H^1(\mathbb{R}^d)), \text{ and} \tag{5.106}$$

$$\text{an equicontinuous sequence in } C([0,T]; L^q(\mathbb{R}^d)) \text{ for all } q \in [2, 2^*). \tag{5.107}$$

Hence, the Ascoli-Arzelá theorem, together with (5.105), gives us that: There exist a subsequence of $\{\widetilde{\Phi}_n^{RN}\}$ (still denoted by the same symbol) and a nontrivial function $\Phi \in L^\infty([0,\infty); H^1(\mathbb{R}^d))$ such that

$$\lim_{n\to\infty} \widetilde{\Phi}_n^{RN} = \Phi \quad \text{weakly* in } L^\infty([0,T]; H^1(\mathbb{R}^d)), \tag{5.108}$$

$$\lim_{n\to\infty} \widetilde{\Phi}_n^{RN} = \Phi \quad \text{strongly in } C([0,T]; L^q_{loc}(\mathbb{R}^d)) \quad \text{for all } q \in [2, 2^*). \tag{5.109}$$

It remains to prove (1.80)–(1.82). We find by (5.86) that: There exists a function $F \in L^\infty([0,\infty); L^{\frac{p+1}{p}}(\mathbb{R}^d))$ such that

$$\lim_{n\to\infty} |\psi_n|^{p-1}\psi_n = F \quad \text{weakly* in } L^\infty([0,T]; L^{\frac{p+1}{p}}(\mathbb{R}^d)). \tag{5.110}$$

Then, it follows from the equation (5.99) and (5.108) that

$$2i\frac{\partial \Phi}{\partial t} + \Delta\Phi + F = 0. \tag{5.111}$$

Here, if F were trivial, then Φ is so since $\Phi(0) = \lim_{n\to\infty} \Phi_n^{RN}(0) = 0$. Therefore, F is nontrivial.

We can prove the formula (1.82) in a way similar to the proof of (1.72) (cf. the estimates (5.83)– (5.85)). ☐

A Generalized virial identity

The proofs of Theorems 1.1, 1.2 and 1.6 are based on a generalization of the virial identity. To state it, we first introduce a positive function w in $W^{3,\infty}([0,\infty))$, which is a variant of the function in [46, 48, 49]:

$$w(r)=\begin{cases} r & \text{if} \quad 0\le r<1, \\ r-(r-1)^{\frac{d}{2}(p-1)+1} & \text{if} \quad 1\le r<1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}, \\ \text{smooth and } w'\le 0 & \text{if} \quad 1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\le r<2, \\ 0 & \text{if} \quad 2\le r. \end{cases} \tag{A.1}$$

Since w is determined by d and p only, we may assume that

$$K:=\|w\|_{W^{3,\infty}}\lesssim 1, \tag{A.2}$$

where the implicit constant depends only on d and p. Using this w, we define

$$\vec{w}_R(x)=(\vec{w}_R^1(x),\dots,\vec{w}_R^d(x)):=\frac{x}{|x|}Rw\left(\frac{|x|}{R}\right), \quad R>0,\ x\in\mathbb{R}^d \tag{A.3}$$

and

$$W_R(x):=2R\int_0^{|x|}w\left(\frac{r}{R}\right)dr, \quad R>0,\ x\in\mathbb{R}^d. \tag{A.4}$$

These functions have the following properties:

Lemma A.1. *Assume that $d\ge 1$ and $2+\frac{4}{d}\le p+1<2^*$. Then, we have that*

$$|\vec{w}_R(x)|^2\le W_R(x) \quad \text{for all } R>0 \text{ and } x\in\mathbb{R}^d, \tag{A.5}$$

$$\|\vec{w}_R\|_{L^\infty}\le 2R \quad \text{for all } R>0, \tag{A.6}$$

$$\|W_R\|_{L^\infty}\le 8R^2 \quad \text{for all } R>0, \tag{A.7}$$

$$\left\|\nabla\vec{w}_R^j\right\|\le 2dK \quad \text{for all } R>0 \text{ and } j=1,\dots,d, \tag{A.8}$$

$$\|\Delta(\operatorname{div}\vec{w}_R)\|_{L^\infty}\le\frac{10d^2K}{R^2} \quad \text{for all } R>0, \tag{A.9}$$

where K is the constant given in (A.2).

Proof of Lemma A.1. Since $w(0)=0$ and $w'\le 1$, we have that

$$|\vec{w}_R(x)|^2=R^2w^2\left(\frac{|x|}{R}\right)\le 2R\int_0^{|x|}w\left(\frac{r}{R}\right)w'\left(\frac{r}{R}\right)dr\le W_R(x),$$

which shows (A.5).

Now, we can easily verify that that

$$\|w\|_{L^\infty}\le 2. \tag{A.10}$$

This estimate (A.10) immediately yields (A.6):

$$|\vec{w}_R(x)| = R\left|w\left(\frac{|x|}{R}\right)\right| \le 2R. \tag{A.11}$$

Moreover, (A.10), together with the fact $\operatorname{supp} w \subset [0,2]$, gives (A.7). Indeed, we have

$$|W_R(x)| \le 2R\int_0^{2R} w\left(\frac{r}{R}\right)\,dr \le 4R^2\left\|w\right\|_{L^\infty} \le 8R^2. \tag{A.12}$$

We shall prove (A.8). A simple calculation shows that

$$\begin{aligned}\left|\partial_k \vec{w}_R^j(x)\right| &= \left|\frac{\delta_{jk}|x|^2 - x_j x_k}{|x|^3}Rw\left(\frac{|x|}{R}\right) + \frac{x_j x_k}{|x|^2}w'\left(\frac{|x|}{R}\right)\right| \\ &\le \frac{R}{|x|}\left|w\left(\frac{|x|}{R}\right)\right| + \left|w'\left(\frac{|x|}{R}\right)\right|.\end{aligned} \tag{A.13}$$

Since we have

$$\frac{R}{|x|}\left|w\left(\frac{|x|}{R}\right)\right| \le \left\|w\right\|_{L^\infty} \quad \text{for all } x \in \mathbb{R}^d, \tag{A.14}$$

the estimate (A.13), together with (A.2), leads to (A.8).

The estimate (A.9) follows from (A.2) and the identity

$$\begin{aligned}\Delta(\operatorname{div}\vec{w}_R)(x) &= \frac{1}{R^2}w'''\left(\frac{|x|}{R}\right) + \frac{2(d-1)}{R|x|}w''\left(\frac{|x|}{R}\right) \\ &\quad + \frac{(d-1)(d-3)}{|x|^2}w'\left(\frac{|x|}{R}\right) - \frac{(d-1)(d-3)}{|x|^3}Rw\left(\frac{|x|}{R}\right).\end{aligned} \tag{A.15}$$

□

Lemma A.2. *Assume that $d \ge 1$ and $2 + \frac{4}{d} \le p+1 < 2^*$. Then, for any $m > 0$, $C > 0$ and $f \in L^2(\mathbb{R}^d)$, there exists $R_0 > 0$ such that*

$$\frac{C}{R^2}(W_R, |f|^2) < m \quad \textit{for all } R \ge R_0. \tag{A.16}$$

Proof of Lemma A.2. For any $m > 0$, $C > 0$ and $f \in L^2(\mathbb{R}^d)$, we can take $R_0' > 0$ such that

$$\int_{|x|\ge R_0'} |f(x)|^2\,dx < \frac{m}{16C}. \tag{A.17}$$

Hence, it follows from (A.7) that

$$\frac{C}{R^2}\int_{|x|\ge R_0'} W_R(x)|f(x)|^2\,dx \le 8C\int_{|x|\ge R_0'}|f(x)|^2\,dx < \frac{m}{2} \quad \text{for all } R > 0. \tag{A.18}$$

Moreover, we have by the definition of W_R (see (A.4)) that

$$\frac{C}{R^2}\int_{|x|<R_0'} W_R(x)|f(x)|^2\,dx \le C\frac{2R_0'K}{R}\left\|f\right\|_{L^2}^2 < \frac{m}{2} \quad \text{for all } R > \tfrac{4CKR_0'\|f\|_{L^2}^2}{m}. \tag{A.19}$$

Combining (A.18) and (A.19), we obtain the desired result. □

Now, we introduce our generalized virial identity:

Lemma A.3 (Generalized virial identity)**.** *Assume that* $d \geq 1$ *and* $2 + \frac{4}{d} \leq p + 1 < 2^*$. *Then, we have*

$$\begin{aligned}(W_R, |\psi(t)|^2) = (W_R, |\psi_0|^2) &+ 2t\Im(\vec{w}_R \cdot \nabla\psi_0, \psi_0) + 2\int_0^t \int_0^{t'} \mathcal{K}(\psi(t''))\, dt'' dt' \\ &- 2\int_0^t \int_0^{t'} \mathcal{K}^R(\psi(t''))\, dt'' dt' \qquad\qquad (A.20) \\ &- \frac{1}{2}\int_0^t \int_0^{t'} (\Delta(\operatorname{div}\vec{w}_R), |\psi(t'')|^2)\, dt'' dt' \qquad \textit{for all } R > 0.\end{aligned}$$

Here, $\mathcal{K}^R$ *is defined by*

$$\mathcal{K}^R(f) = \int_{\mathbb{R}^d} \rho_1(x)|\nabla f(x)|^2 + \rho_2(x)\left|\frac{x}{|x|} \cdot \nabla f(x)\right|^2 - \rho_3(x)|f(x)|^{p+1} dx, \quad f \in H^1(\mathbb{R}^d), \tag{A.21}$$

where

$$\rho_1(x) := 1 - \frac{R}{|x|} w\left(\frac{|x|}{R}\right), \tag{A.22}$$

$$\rho_2(x) := \frac{R}{|x|} w\left(\frac{|x|}{R}\right) - w'\left(\frac{|x|}{R}\right), \tag{A.23}$$

$$\rho_3(x) := \frac{p-1}{2(p+1)}\left\{d - w'\left(\frac{|x|}{R}\right) - \frac{d-1}{|x|} R w\left(\frac{|x|}{R}\right)\right\}. \tag{A.24}$$

Remark A.1. *If* $d = 1$ *or* ψ *is radially symmetric, then we have*

$$\mathcal{K}^R(\psi) = \int_{\mathbb{R}^d} \rho_0(x)|\nabla\psi(x)|^2 - \rho_3(x)|\psi(x)|^{p+1}\, dx, \tag{A.25}$$

where

$$\rho_0(x) := 1 - w'\left(\frac{|x|}{R}\right) = \rho_1(x) + \rho_2(x). \tag{A.26}$$

Proof of Lemma A.3. By the formula (B.17) in [46], we immediately obtain (A.20). ☐

In the next lemma, we give several properties of the weight functions ρ_1, ρ_2, ρ_3 and ρ_0:

Lemma A.4. *Assume that* $d \geq 1$ *and* $2 + \frac{4}{d} \leq p + 1 < 2^*$. *Then, for all* $R > 0$, *we have* *the followings:*

$$\operatorname{supp}\rho_j = \{x \in \mathbb{R}^d \mid |x| \geq R\} \quad \textit{for all } R > 0 \textit{ and } j = 0,1,2,3, \tag{A.27}$$

$$\inf_{x\in\mathbb{R}^d} \rho_j(x) \geq 0 \quad \textit{for all } R > 0 \textit{ and } j = 0,1,2,3, \tag{A.28}$$

$$\rho_0(x) = 1 \quad \textit{if } |x| \geq 2R, \tag{A.29}$$

$$\rho_3(x) = \frac{d(p-1)}{2(p+1)} \quad \textit{if } |x| \geq 2R, \tag{A.30}$$

$$\|\rho_j\|_{L^\infty} \leq K_j \quad \textit{for all } j = 0,1,2,3, \tag{A.31}$$

$$\sup_{x\in\mathbb{R}^d} \max\left\{-\frac{x}{|x|}\nabla\sqrt{\rho_3(x)},\ 0\right\} \leq \frac{K_3'}{R}, \tag{A.32}$$

$$\inf_{|x|\geq R} \frac{\rho_0(x)}{\rho_3(x)} \geq K_4, \tag{A.33}$$

where K_j $(j = 1,2,3)$, K_3' and K_4 are some constants independent of R.

Proof of Lemma A.4. We give proofs of (A.32) and (A.33) only.

We first prove (A.32). A direct calculation shows that

$$\frac{x}{|x|}\nabla\sqrt{\rho_3(x)} = \frac{-\frac{1}{R}w''\left(\frac{|x|}{R}\right) - \frac{d-1}{|x|}w'\left(\frac{|x|}{R}\right) + \frac{(d-1)R}{|x|^2}w\left(\frac{|x|}{R}\right)}{\sqrt{\frac{2(p+1)}{p-1}}\sqrt{d - w'\left(\frac{|x|}{R}\right) - \frac{(d-1)R}{|x|}w\left(\frac{|x|}{R}\right)}}. \tag{A.34}$$

Since $\operatorname{supp}\rho_3 = \{x \in \mathbb{R}^d \mid |x| \geq R\}$ (see (A.27)), it suffices to consider the case $|x| \geq R$. When $R \leq |x| < R\left\{1 + \left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\right\}$, we have from (A.34) that

$$\begin{aligned}
&\max\left\{-\frac{x}{|x|}\nabla\sqrt{\rho_3(x)},\ 0\right\} \\
&= \frac{\max\left\{\left(\frac{|x|}{R} - 1\right)^{\frac{d}{2}(p-1)-1}\left(\frac{d^2(p-1)^2+4d(p-1)+4(d-1)}{4R} + \frac{(d-1)R}{|x|^2} - \frac{d(d-1)(p-1)+2}{2|x|}\right),\ 0\right\}}{\sqrt{\frac{2(p+1)}{p-1}}\left(\frac{|x|}{R} - 1\right)^{\frac{d}{4}(p-1)}\sqrt{\frac{d}{2}(p-1) + d - \frac{(d-1)R}{|x|}}} \\
&\leq \left(\frac{|x|}{R} - 1\right)^{\frac{d}{4}(p-1)-1} \frac{\left\{\frac{d^2(p-1)^2+4d(p-1)+4(d-1)}{4R} + \frac{(d-1)R}{|x|^2} + \frac{d(d-1)(p-1)+2}{2|x|}\right\}}{\sqrt{\frac{2(p+1)}{p-1}}\sqrt{\frac{d}{2}(p-1) + 1}} \\
&\lesssim \frac{1}{R},
\end{aligned} \tag{A.35}$$

where the implicit constant depends only on d and p.

On the other hand, when $R\left\{1 + \left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\right\} \leq |x|$, we have

$$w'\left(\frac{|x|}{R}\right) \leq 0, \qquad \frac{R}{|x|}w\left(\frac{|x|}{R}\right) \leq 1 \tag{A.36}$$

and therefore we obtain from (A.34) that

$$\begin{aligned}\max\left\{-\frac{x}{|x|}\nabla\sqrt{\rho_3(x)},\ 0\right\} &\le \frac{\left|\frac{1}{R}w''\left(\frac{|x|}{R}\right)\right| + \left|\frac{d-1}{|x|}w'\left(\frac{|x|}{R}\right)\right| + \left|\frac{(d-1)R}{|x|^2}w\left(\frac{|x|}{R}\right)\right|}{\sqrt{\frac{2(p+1)}{p-1}}\sqrt{d-\frac{(d-1)R}{|x|}w\left(\frac{|x|}{R}\right)}} \\ &\lesssim \frac{1}{R}\|w\|_{W^{2,\infty}},\end{aligned} \tag{A.37}$$

where the implicit constant depends only on d and p. Thus, we see that (A.32) holds.

Next, we prove (A.33). The starting point of the proof is the identity:

$$\frac{p-1}{2(p+1)}\frac{\rho_0(x)}{\rho_3(x)} = \frac{\rho_0(x)}{\rho_0(x)+(d-1)\left\{1-\frac{R}{|x|}w\left(\frac{|x|}{R}\right)\right\}} \quad \text{for all } x\in\mathbb{R}^d \text{ with } |x|\ge R. \tag{A.38}$$

When $R\le|x|<R\left\{1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\right\}$, we have from (A.38) that

$$\begin{aligned}\frac{p-1}{2(p+1)}\frac{\rho_0(x)}{\rho_3(x)} &= \frac{\rho_0(x)}{\rho_0(x)+(d-1)\frac{R}{|x|}\left(\frac{|x|}{R}-1\right)^{\frac{d}{2}(p-1)+1}} \\ &= \frac{\left(\frac{|x|}{R}-1\right)^{\frac{d}{2}(p-1)}}{\left(\frac{|x|}{R}-1\right)^{\frac{d}{2}(p-1)}+\frac{2(d-1)}{d(p-1)+2}\frac{R}{|x|}\left(\frac{|x|}{R}-1\right)^{\frac{d}{2}(p-1)+1}} \\ &= \frac{1}{1+\frac{2(d-1)}{d(p-1)+2}\left(1-\frac{R}{|x|}\right)} \\ &\ge \frac{1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}}{1+\frac{d(p+1)}{d(p-1)+2}\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}}.\end{aligned} \tag{A.39}$$

On the other hand, when $R\left\{1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\right\}\le|x|$, we have

$$1\le\rho_0(x)\le 1+K, \qquad 0\le 1-\frac{R}{|x|}w\left(\frac{|x|}{R}\right)\le 1, \tag{A.40}$$

and therefore we obtain from (A.38) that

$$\frac{p-1}{2(p+1)}\frac{\rho_0(x)}{\rho_3(x)} \ge \frac{1}{K+d} \quad \text{for all } x\in\mathbb{R}^d \text{ with } R\left\{1+\left(\frac{2}{d(p-1)+2}\right)^{\frac{2}{d(p-1)}}\right\}\le|x|. \tag{A.41}$$

Thus, we see that (A.33) holds. ☐

B Compactness device I

We recall the following sequence of lemmata.

Lemma B.1 (Frölich, Lieb and Loss [18], Nawa [41]). *Let $1 < \alpha < \beta < \gamma < \infty$ and let f be a measurable function on $\mathbb{R}^d$ with*

$$\|f\|_{L^\alpha}^\alpha \le C_\alpha, \quad C_\beta \le \|f\|_{L^\beta}^\beta, \quad \|f\|_{L^\gamma}^\gamma \le C_\gamma$$

for some positive constants $C_\alpha, C_\beta, C_\gamma$. Then, we have

$$\mathcal{L}^d\left(\Big[|f| > \eta\Big]\right) > \frac{C_\beta}{2}\eta^\beta \quad \textit{for all } 0 < \eta < \min\left\{1,\ \left(\tfrac{C_\beta}{4C_\alpha}\right)^{\frac{1}{\beta-\alpha}},\ \left(\tfrac{C_\beta}{4C_\gamma}\right)^{\frac{1}{\gamma-\beta}}\right\}.$$

Proof of Lemma B.1. Let η be a constant satisfying

$$0 < \eta < \min\left\{1,\ \left(\frac{C_\beta}{4C_\alpha}\right)^{\frac{1}{\beta-\alpha}},\ \left(\frac{C_\beta}{4C_\gamma}\right)^{\frac{1}{\gamma-\beta}}\right\}. \tag{B.1}$$

Then, we can easily verify that

$$\begin{aligned} C_\beta &= \int_{[|f|\le\eta]} |f(x)|^\beta\,dx + \int_{\left[\eta<|f|\le\frac{1}{\eta}\right]} |f(x)|^\beta\,dx + \int_{\left[\frac{1}{\eta}<|f|\right]} |f(x)|^\beta\,dx \\ &\le \eta^{\beta-\alpha}\int_{[|f|\le\eta]} |f(x)|^\alpha\,dx + \left(\frac{1}{\eta}\right)^\beta \mathcal{L}\left(\Big[|f|\ge\eta\Big]\right) + \eta^{\gamma-\beta}\int_{[|f|\le\eta]} |f(x)|^\gamma\,dx \qquad \text{(B.2)} \\ &\le \frac{C_\beta}{2} + \left(\frac{1}{\eta}\right)^\beta \mathcal{L}\left(\Big[|f|\ge\eta\Big]\right). \end{aligned}$$

This estimate (B.2) immediately gives us the desired result. □

Lemma B.2 (Lieb [34], Nawa [41]). *Let $1 < q < \infty$, and let f be a function in $W^{1,q}(\mathbb{R}^d)$ with*

$$\|\nabla f\|_{L^q}^q \le D_1, \quad \mathcal{L}^d\left(\Big[|f| > \eta\Big]\right) \ge D_2 \tag{B.3}$$

for some positive constants D_1, D_2 and η. We put

$$q^\dagger = \begin{cases} \dfrac{qd}{d-q} & \textit{if} \quad q < d, \\ 2q & \textit{if} \quad q \ge d. \end{cases} \tag{B.4}$$

Then, there exists $y \in \mathbb{R}^d$ such that

$$\mathcal{L}^d\left(\left[|f(\cdot+y)| \ge \frac{\eta}{2}\right] \cap B_1(0)\right) \gtrsim \left(\frac{1+\eta^q D_2}{1+D_1}\right)^{\frac{q}{q^\dagger-q}}, \tag{B.5}$$

where the implicit constant depends only on d and q, and $B_1(0)$ is the ball in $\mathbb{R}^d$ with center 0 and radius 1.

Proof of Lemma B.2. Let f be a function in $W^{1,q}(\mathbb{R}^d)$ satisfying (B.3). Put

$$g(x) = \max\left\{|f(x)| - \frac{\eta}{2}, 0\right\}. \tag{B.6}$$

Then, we easily verify that

$$g \in W^{1,q}(\mathbb{R}^d), \qquad \|\nabla g\|_{L^q(\mathbb{R}^d)} \le D_1, \tag{B.7}$$

$$\operatorname{supp} g \subset \left[|f| \ge \frac{\eta}{2}\right]. \tag{B.8}$$

We first claim that

$$\int_{Q_y} |\nabla g(x)|^q \, dx < \left(1 + \frac{D_1}{\|g\|_{L^q}^q}\right) \int_{Q_y} |g(x)|^q \, dx \quad \text{for some } y \in \mathbb{R}^d, \tag{B.9}$$

where Q_y denotes the cube in $\mathbb{R}^d$ with the center y and side length $\frac{2}{\sqrt{d}}$. We note that Q_y is inscribed in $B_1(y)$ for all $y \in \mathbb{R}^d$.

Let $\{y_n\}_{n\in\mathbb{N}}$ be a sequence in $\mathbb{R}^d$ such that

$$\bigcup_{n\in\mathbb{N}} Q_{y_n} = \mathbb{R}^d, \qquad \overset{\circ}{Q}_{y_m} \cap \overset{\circ}{Q}_{y_n} = \emptyset \quad \text{for } m \neq n \tag{B.10}$$

Supposing the contrary that (B.9) fails, we have

$$\int_{Q_{y_n}} |\nabla g(x)|^q \, dx \ge \left(1 + \frac{D_1}{\|g\|_{L^q}^q}\right) \int_{Q_{y_n}} |g(x)|^q \, dx \quad \text{for all } n \in \mathbb{N}. \tag{B.11}$$

Then, summing (B.11) over all $n \in \mathbb{N}$ yields that

$$D_1 \ge \int_{\mathbb{R}^d} |\nabla g(x)|^q \, dx \ge \left(1 + \frac{D_1}{\|g\|_{L^q}^q}\right) \int_{\mathbb{R}^d} |g(x)|^q \, dx > D_1, \tag{B.12}$$

which is a contradiction. Hence, (B.9) holds.

Now, it follows from (B.9) that: There exists $y_0 \in \mathbb{N}$ such that

$$\int_{Q_{y_0}} |\nabla g(x)|^q \, dx + |g(x)|^q \, dx < \left(2 + \frac{D_1}{1 + \|g\|_{L^q}^q}\right) \int_{Q_{y_0}} |g(x)|^q \, dx. \tag{B.13}$$

On the other hand, the Sobolev embedding leads us to that

$$\left(\int_{Q_{y_0}} |g(x)|^{q^\dagger} \, dx,\right)^{\frac{q}{q^\dagger}} \lesssim \int_{Q_{y_0}} |\nabla g(x)|^q \, dx + |g(x)|^q \, dx, \tag{B.14}$$

where $q^\dagger$ is the exponent defined in (B.4), and the implicit constant depends only on d and q. Combining these estimates (B.13) and (B.14), we obtain that

$$\begin{aligned} \left(\int_{Q_{y_0}} |g(x)|^{q^\dagger} \, dx\right)^{\frac{q}{q^\dagger}} &\lesssim \left(2 + \frac{D_1}{1 + \|g\|_{L^q}^q}\right) \int_{Q_{y_0}} |g(x)|^q \, dx \\ &\lesssim \left(2 + \frac{D_1}{1 + \|g\|_{L^q}^q}\right) \mathcal{L}\left(Q_{y_0} \cap \operatorname{supp} g\right)^{1-\frac{q}{q^\dagger}} \left(\int_{Q_{y_0}} |g(x)|^{q^\dagger} \, dx\right)^{\frac{q}{q^\dagger}}, \end{aligned} \tag{B.15}$$

where the implicit constant depends only on d and q. Hence, we see that

$$\left(\frac{1+\|g\|_{L^q}^q}{D_1+2+2\|g\|_{L^q}^q}\right)^{\frac{q^\dagger}{q^\dagger-q}} \lesssim \mathcal{L}\left(Q_{y_0}\cap \operatorname{supp} g\right), \tag{B.16}$$

where the implicit constant depends only on d and q. Here, it follows from the definition of g (see (B.6)) and the assumption (B.3) that

$$\|g\|_{L^q}^q \geq \int_{[|f|\geq \eta]} \left(|f(x)|-\frac{\eta}{2}\right)^q \geq \left(\frac{\eta}{2}\right)^q D_2. \tag{B.17}$$

Since $\operatorname{supp} g \subset \left[|f| \geq \frac{\eta}{2}\right]$ (see (B.8)), the estimate (B.16), together with (B.17), gives us the desired result. ☐

Lemma B.3 (Lieb [34], Nawa [41])**.** *Let* $1<q<\infty$, *and let* $\{f_n\}_{n\in\mathbb{N}}$ *be a uniformly bounded sequence in* $W^{1,q}(\mathbb{R}^d)$ *with*

$$\inf_{n\in\mathbb{N}} \mathcal{L}^d\left(\left[|f_n|>\delta\right]\right) \geq C$$

for some constants $\delta>0$ *and* $C>0$. *Then, there exists a sequence* $\{y_n\}_{n\in\mathbb{R}^d}$ *in* $\mathbb{R}^d$ *and a nontrivial function* $f\in W^{1,q}(\mathbb{R}^d)$ *such that*

$$f_n(\cdot+y_n)\to f \quad \text{weakly in } W^{1,q}(\mathbb{R}^d). \tag{B.18}$$

Lemma B.4 (Brezis and Lieb [8], Nawa [41])**.** *Let* $0<q<\infty$ *and let* $\{f_n\}_{n\in\mathbb{N}}$ *be a uniformly bounded sequence in* $L^q(\mathbb{R}^d)$ *with* $f_n\to f$ *a.e. in* $\mathbb{R}^d$ *for some* $f\in L^q(\mathbb{R}^d)$. *Then, we have*

$$\lim_{n\to\infty}\int_{\mathbb{R}^d}\left||f_n(x)|^q-|f_n(x)-f(x)|^q-|f(x)|^q\right|dx=0. \tag{B.19}$$

C Compactness device II

We recall the following sequence of lemmas: All their proofs are found in Appendix C of [46].

Lemma C.1 (Nawa [46])**.** *Let* $1<\alpha<\beta<\gamma<\infty$, $I\subset\mathbb{R}$ *and let* $u\in C(I;L^\alpha(\mathbb{R}^d)\cap L^\gamma(\mathbb{R}^d))$ *with*

$$\sup_{t\in I}\|u(t)\|_{L^\alpha}^\alpha \leq C_\alpha, \quad C_\beta \leq \sup_{t\in I}\|u(t)\|_{L^\beta}^\beta, \quad \sup_{t\in I}\|u(t)\|_{L^\gamma}^\gamma \leq C_\gamma$$

for some positive constants C_α, C_β *and* C_γ. *Then, we have*

$$\sup_{t\in I}\mathcal{L}^d\left(\left[|u(t)|>\delta\right]\right) > C$$

for some constants $C>0$ *and* $\delta>0$ *depending only on* α, β, γ, C_α, C_β *and* C_γ.

Lemma C.2 (Nawa [46])**.** *Let* $1 \le q < \infty$, $I \Subset \mathbb{R}$ *and let* u *be a function in* $C(I; W^{1,q}(\mathbb{R}^d))$ *such that*

$$\sup_{t\in I} \|\nabla u(t)\|_{L^q} \le C_q,$$

$$\sup_{t\in I} \mathcal{L}^d \left(\Big[|u(t)| > \delta \Big] \right) \ge C$$

for some positive constants C_q, δ *and* C*. Then, there exists* $y \in \mathbb{R}^d$ *such that*

$$\sup_{t\in I} \mathcal{L}^d \left(\left[|u(\cdot + y, t)| > \frac{\delta}{2} \right] \cap B_1(0) \right) \ge C'$$

for some constant $C' > 0$ *depending only on* C_q, C, δ *and* d*, where* $B_1(0)$ *is the ball in* $\mathbb{R}^d$ *with center* 0 *and radius* 1*.*

Lemma C.3 (Nawa [46])**.** *Let* $0 < q < \infty$, $I \Subset \mathbb{R}$ *and let* $\{u_n\}_{n\in\mathbb{N}}$ *be a equicontinuous and uniformly bounded sequence in* $C(I; L^q(\mathbb{R}^d))$ *with* $\lim_{n\to\infty} u_n = u$ *a.e. in* $\mathbb{R}^d \times I$ *for some* $u \in C(I; L^q(\mathbb{R}^d))$*. Then, we have that:*

$$\lim_{n\to\infty} \sup_{t\in I} \int_{\mathbb{R}^d} \Big| |u_n(x,t)|^q - |u_n(x,t) - u(x,t)|^q - |u(x,t)|^q \Big| \, dx = 0, \tag{C.1}$$

$$\lim_{n\to\infty} \left\{ |u_n|^{q-1}u_n - |u_n - u|^{q-1}(u_n - u) - |u|^{q-1}u \right\} = 0 \quad \textit{strongly in } L^\infty(I; L^{q'}(\mathbb{R}^d)). \tag{C.2}$$

D Elementary inequalities

In this section, we summarize elementary inequalities often used in this paper. We begin with the following obvious inequalities.

Lemma D.1. *Let* $1 < q < \infty$*. Then, we have that*

$$\big||a|^q - |b|^q\big| \lesssim \left(|a|^{q-1} + |b|^{q-1} \right) |a - b| \qquad \textit{for all } a, b \in \mathbb{C}, \tag{D.1}$$

$$\Big| |a|^{q-1}\, a - |b|^{q-1}\, b \Big| \lesssim \left(|a|^{q-1} + |b|^{q-1} \right) |a - b| \qquad \textit{for all } a, b \in \mathbb{C}, \tag{D.2}$$

where the implicit constants depend only on q*.*

Besides the above, we also have the following inequality.

Lemma D.2. *Let* $0 < q < \infty$, $L \in \mathbb{N}$ *and* $a_1, \dots, a_L \in \mathbb{C}$*. Then, we have*

$$\left| \left| \sum_{k=1}^{L} a_k \right|^q \sum_{l=1}^{L} a_l - \sum_{l=1}^{L} |a_l|^q \, a_l \right| \le C \sum_{l=1}^{L} \sum_{\substack{1\le k\le L \\ k\ne l}} |a_l| \, |a_k|^q$$

for some constant $C > 0$ *depending only on* q *and* L*.*

Proof of Lemma D.2. Since

$$\left|\left|\sum_{k=1}^{L} a_k\right|^q \sum_{l=1}^{L} a_l - \sum_{l=1}^{L} |a_l|^q a_l\right| \leq \sum_{l=1}^{L} |a_l| \left|\left|\sum_{k=1}^{L} a_k\right|^q - |a_l|^q\right|,$$

it suffices to prove that

$$\sum_{l=1}^{L} |a_l| \left|\left|\sum_{k=1}^{L} a_k\right|^q - |a_l|^q\right| \leq C \sum_{l=1}^{L} \sum_{\substack{1\leq k\leq L \\ k\neq l}} |a_l|\,|a_k|^q \tag{D.3}$$

for some constant $C > 0$ depending only on q and L. Rearranging the sequence, we may assume that

$$|a_1| \leq |a_2| \leq \cdots \leq |a_{L-1}| \leq |a_L|.$$

Then, we have

$$\begin{aligned}\sum_{l=1}^{L-1} |a_l| \left|\left|\sum_{k=1}^{L} a_k\right|^q - |a_l|^q\right| &\leq \sum_{l=1}^{L-1} |a_l| \left\{\left(\sum_{k=1}^{L} |a_k|\right)^q + |a_l|^q\right\} \\ &\leq \sum_{l=1}^{L-1} |a_l| \left\{(L|a_L|)^q + |a_L|^q\right\} \\ &= (L^q+1) \sum_{l=1}^{L-1} |a_l|\,|a_L|^q \\ &\leq (L^q+1) \sum_{l=1}^{L} \sum_{\substack{1\leq k\leq L \\ k\neq l}} |a_l|\,|a_k|^q.\end{aligned}$$

It remains an estimate for the term

$$|a_L| \left|\left|\sum_{k=1}^{L} a_k\right|^q - |a_L|^q\right|.$$

When $q > 1$, we have by (D.1) that

$$\begin{aligned}|a_L| \left|\left|\sum_{k=1}^{L} a_k\right|^q - |a_L|^q\right| &\lesssim |a_L| \left(\left|\sum_{k=1}^{L} a_k\right|^{q-1} + |a_L|^{q-1}\right) \left|\sum_{l=1}^{L-1} a_l\right| \\ &\leq \left(L^{q-1}+1\right) |a_L|^q \sum_{l=1}^{L-1} |a_l| \\ &\leq \left(L^{q-1}+1\right) \sum_{l=1}^{L} \sum_{\substack{1\leq k\leq L \\ k\neq l}} |a_l|\,|a_k|^q,\end{aligned}$$

where the implicit constant depends only on q. Hence, in the case $q > 1$, we have obtained the result. We consider the case $0 < q \leq 1$. When $|a_{L-1}| \geq |a_L|/L$, we have

$$\begin{aligned}
|a_L| \left| \left| \sum_{l=1}^{L} a_l \right|^q - |a_L|^q \right| &\leq |a_L| \left(L^q + 1 \right) |a_L|^q \\
&\leq L \, |a_{L-1}| \left(L^q + 1 \right) |a_L|^q \\
&\leq L \left(L^q + 1 \right) \sum_{l=1}^{L} \sum_{\substack{1 \leq k \leq L \\ k \neq l}} |a_l| \, |a_k|^q .
\end{aligned}$$

On the other hand, when $|a_{L-1}| \leq |a_L|/L$, one can see that

$$\frac{|a_L|}{L} \leq |a_L| - \left| \sum_{l=1}^{L-1} a_l \right| \leq \left| \sum_{l=1}^{L} a_l \right| \leq |a_L| + \left| \sum_{l=1}^{L-1} a_l \right| .$$

Moreover, we have by the convexity of the function $f(t) = t^q$ $(0 < q \leq 1)$ that

$$\left(|a_L| + \left| \sum_{l=1}^{L-1} a_l \right| \right)^q - |a_L|^q \leq |a_L|^q - \left(|a_L| - \left| \sum_{l=1}^{L-1} a_l \right| \right)^q .$$

Hence, it follows from these estimates that

$$\begin{aligned}
\left| \left| \sum_{l=1}^{L} a_l \right|^q - |a_L|^q \right| &\leq |a_L|^q - \left(|a_L| - \left| \sum_{l=1}^{L-1} a_l \right| \right)^q \\
&\leq q \left| \sum_{l=1}^{L-1} a_l \right| \left(|a_L| - \left| \sum_{l=1}^{L-1} a_l \right| \right)^{q-1} \\
&\leq q \left| \sum_{l=1}^{L-1} a_l \right| |a_L|^{q-1}
\end{aligned}$$

and therefore

$$|a_L| \left| \left| \sum_{l=1}^{L} a_l \right|^q - |a_L|^q \right| \leq q \left| \sum_{l=1}^{L-1} a_l \right| |a_L|^q \leq q \sum_{l=1}^{L} \sum_{\substack{1 \leq k \leq L \\ k \neq l}} |a_l| \, |a_k|^q .$$

Thus, we have completed the proof. ☐

E Concentration function

In this section, $B_R(a)$ denotes an open ball in $\mathbb{R}^d$ with the center $a \in \mathbb{R}^d$ and the radius R:

$$B_R(a) := \left\{ x \in \mathbb{R}^d \;\middle|\; |x - a| < R \right\}.$$

In order to trace the "fake" soliton, we prepare Proposition E.2 below. We begin with:

Lemma E.1 (Nawa [42]). *Let $0 < T_m \le \infty$, $1 < q < \infty$ and let ρ be a non-negative function in $C([0,T_m); L^1(\mathbb{R}^d) \cap L^q(\mathbb{R}^d))$. Suppose that*

$$\int_{|x-y_0|<R_0} \rho(x,t_0)\,dx > C_0 \tag{E.1}$$

for some $C_0 > 0$, $R_0 > 0$ and $(y_0,t_0) \in \mathbb{R}^d \times [0,T_m)$. Then, there exist $\theta > 0$ and $\Gamma > 0$, both depending only on d, q, ρ, C_0, R_0 and t_0, such that

$$\int_{|x-y|<R_0} \rho(x,t)\,dx > C_0, \quad \forall (y,t) \in B_\Gamma(y_0) \times (t_0-\theta, t_0+\theta).$$

Remark E.1. *When $\rho\colon [0,T_m) \to L^1(\mathbb{R}^d)$ is uniformly continuous, we can take θ independent of t_0. Unfortunately, we are not in a case to assume the uniform continuity.*

Proof of Lemma E.1. Put

$$\varepsilon_0 = \int_{|x-y_0|<R_0} \rho(x,t_0)\,dx - C_0.$$

Then, we have by (E.1) that $\varepsilon_0 > 0$. Since we have by the Hölder inequality that

$$\int_{B_{R_0}(y_0)\setminus B_{R_0}(y)} \rho(x,t_0)\,dx \le \mathcal{L}^d\left(B_{R_0}(y_0)\setminus B_{R_0}(y)\right)^{1-\frac{1}{q}} \|\rho(t_0)\|_{L^q(\mathbb{R}^d)}, \quad \forall y \in \mathbb{R}^d, \tag{E.2}$$

we can take $\Gamma > 0$, depending only on d, q, ρ, C_0, R_0 and t_0, such that

$$\int_{B_{R_0}(y_0)\setminus B_{R_0}(y)} \rho(x,t_0)\,dx < \frac{\varepsilon_0}{3}, \quad \forall y \in B_\Gamma(y_0). \tag{E.3}$$

Moreover, it follows from (E.3) that

$$\begin{aligned}
&\int_{B_{R_0}(y_0)} \rho(x,t_0)\,dx \\
&= \int_{B_{R_0}(y)} \rho(x,t_0)\,dx + \int_{B_{R_0}(y_0))\setminus B_{R_0}(y)} \rho(x,t_0)\,dx - \int_{B_{R_0}(y)\setminus B_{R_0}(y_0)} \rho(x,t_0)\,dx \\
&< \int_{B_{R_0}(y)} \rho(x,t_0)\,dx + \frac{\varepsilon_0}{3}, \qquad \forall y \in B_\Gamma(y_0),
\end{aligned} \tag{E.4}$$

so that we have

$$\int_{B_{R_0}(y)} \rho(x,t_0)\,dx > \int_{B_{R_0}(y_0)} \rho(x,t_0)\,dx - \frac{\varepsilon_0}{3}, \quad \forall y \in B_\Gamma(y_0). \tag{E.5}$$

Now, since ρ is non-negative, we have that, for any measurable set $\Omega \subset \mathbb{R}^d$,

$$\int_\Omega \rho(x,t_0)\,dx - \int_\Omega \rho(x,t)\,dx \le \|\rho(t)-\rho(t_0)\|_{L^1(\mathbb{R}^d)}, \quad \forall t \in [0,T_m). \tag{E.6}$$

This estimate (E.6), together with the continuity of $\rho\colon [0,T_m) \to L^1(\mathbb{R}^d)$, gives us that there exists $\theta > 0$, depending only on d, ρ, C_0, R_0 and t_0, such that

$$\begin{aligned}
\int_{B_{R_0}(y)\cap B_{R_0}(y_0)} \rho(x,t_0)\,dx - \int_{B_{R_0}(y)\cap B_{R_0}(y_0)} \rho(x,t)\,dx < \frac{\varepsilon_0}{3},& \\
\forall (y,t) \in \mathbb{R}^d \times (t_0-\theta, t_0+\theta),&
\end{aligned} \tag{E.7}$$

and

$$\int_{B_{R_0}(y)\setminus B_{R_0}(y_0)} \rho(x,t_0)\,dx - \int_{B_{R_0}(y)\setminus B_{R_0}(y_0)} \rho(x,t)\,dx < \frac{\varepsilon_0}{3}, \qquad \forall (y,t) \in \mathbb{R}^d \times (t_0-\theta, t_0+\theta), \tag{E.8}$$

Combining these estimates, we see that there exists $\theta > 0$, depending only on d, ρ, C_0, R_0 and t_0, such that

$$\int_{B_{R_0}(y)} \rho(x,t_0)\,dx - \int_{B_{R_0}(y)} \rho(x,t)\,dx < \frac{2\varepsilon_0}{3}, \qquad \forall (y,t) \in \mathbb{R}^d \times (t_0-\theta, t_0+\theta). \tag{E.9}$$

Moreover, it follows from the estimates (E.5) and (E.9) that there exist $\theta > 0$ and $\Gamma > 0$, both depending only on d, q, ρ, C_0, R_0 and t_0, such that

$$\begin{aligned}\int_{|x-y|<R_0} \rho(x,t)\,dx &= \int_{B_{R_0}(y)} \rho(x,t_0)\,dx - \left(\int_{B_{R_0}(y)} \rho(x,t_0)\,dx - \int_{B_{R_0}(y)} \rho(x,t)\,dx \right) \\ &> \int_{B_{R_0}(y)} \rho(x,t_0)\,dx - \frac{2\varepsilon_0}{3} \\ &\geq \int_{B_{R_0}(y_0)} \rho(x,t_0)\,dx - \varepsilon_0 = C_0\end{aligned}$$

for all $y \in \mathbb{R}^d$ with $|y - y_0| < \Gamma$, and all $t \in [0, T_m)$ with $|t - t_0| < \theta$. Thus, we have proved the lemma. $\square$

Proposition E.2 (Nawa [42, 43]). *Assume that* $d \geq 1$. *Let* $0 < T_m \leq \infty$, $1 < q < \infty$, *and let* ρ *be a non-negative function in* $C([0,T_m); L^1(\mathbb{R}^d) \cap L^q(\mathbb{R}^d))$ *with*

$$\|\rho(t)\|_{L^1(\mathbb{R}^d)} = 1 \quad \text{for all } t \in [0, T_m). \tag{E.10}$$

We put

$$A_\rho = \sup_{R>0} \liminf_{t \to T_m} \sup_{y \in \mathbb{R}^d} \int_{|x-y| \leq R} \rho(x,t)\,dx.$$

If $A_\rho > \frac{1}{2}$, *then for all* $\varepsilon \in (0,1)$, *there exist* $R_\varepsilon > 0$, $T_\varepsilon > 0$ *and a continuous path* $\gamma_\varepsilon \in C([T_\varepsilon, T_m); \mathbb{R}^d)$ *such that*

$$\int_{|x-\gamma_\varepsilon(t)| \leq R} \rho(x,t)\,dx \geq (1-\varepsilon) A_\rho \quad \text{for all } R \geq 3R_\varepsilon \text{ and } t \in [T_\varepsilon, T_m).$$

Proof of Proposition E.2. We put $\eta := 2A_\rho - 1$, so that $A_\rho = \frac{1}{2} + \frac{\eta}{2}$. Then, the assumptions (E.10) and $A_\rho > \frac{1}{2}$ imply that $0 < \eta \leq 1$. We choose arbitrarily $\varepsilon \in (0, \frac{\eta}{1+\eta})$ and fix it.

It follows from the definition of A_ρ that for our $\varepsilon > 0$, there exist $R_\varepsilon > 0$ and $T_\varepsilon > 0$ with the following property: for all $t \in [T_\varepsilon, T_m)$, there exists a point $y_\varepsilon(t) \in \mathbb{R}^d$ such that

$$\int_{|x-y_\varepsilon(t)| \leq R_\varepsilon} \rho(x,t)\,dx > (1-\varepsilon)\, A_\rho. \tag{E.11}$$

For these R_ε and T_ε, we shall construct a continuous path $\gamma_\varepsilon\colon [T_\varepsilon, T_m) \to \mathbb{R}^d$ satisfying that

$$\int_{|x-\gamma_\varepsilon(t)|\le 3R_\varepsilon} \rho(x,t)\,dx \ge (1-\varepsilon)\,A_\rho \quad \text{for all } t\in[T_\varepsilon,T_m), \tag{E.12}$$

which will lead us to the desired conclusion. To this end, we define:

$$T_\varepsilon^* = \sup\left\{ t\in[T_\varepsilon,T_m) \;\middle|\; \int_{|x-y_\varepsilon(T_\varepsilon)|\le R_\varepsilon} \rho(x,t)\,dx > (1-\varepsilon)\,A_\rho \right\}, \tag{E.13}$$

where $y(T_\varepsilon)$ is a point found in (E.11). By Lemma E.1, it follows from (E.11) that $T_\varepsilon^* > T_\varepsilon$. If $T_\varepsilon^* = T_m$, then there is nothing to prove: $\gamma_\varepsilon(t)\equiv y_\varepsilon(T_\varepsilon)$ $(t\in[T_\varepsilon,T_m))$ is the desired continuous path.

Hence, we consider the case of $T_\varepsilon^* < T_m$. In this case, by Lemma E.1 again, we find that: there exists a constant $\theta_\varepsilon(T_\varepsilon^*)>0$ such that

$$\int_{|x-y_\varepsilon(T_\varepsilon^*)|\le R_\varepsilon} \rho(x,t)\,dx > (1-\varepsilon)\,A_\rho \qquad \text{for all } t\in(T_\varepsilon^*-\theta_\varepsilon(T_\varepsilon^*), T_\varepsilon^*+\theta_\varepsilon(T_\varepsilon^*))\;. \tag{E.14}$$

Here, we claim that:

$$(B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\times\{t\})\cap(B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))\times\{t\})\ne\emptyset \quad \text{for} \quad t\in(T_\varepsilon^*-\theta_\varepsilon(T_\varepsilon^*), T_\varepsilon^*]. \tag{E.15}$$

For: we suppose the contrary that $t_1\in(T_\varepsilon^*-\theta_\varepsilon(T_\varepsilon^*), T_\varepsilon^*]$ such that

$$(B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\times\{t_1\})\cap(B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))\times\{t_1\})=\emptyset. \tag{E.16}$$

Then, it follows from the assumption (E.10) and (E.16) that

$$\begin{aligned} 1 &\ge \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cup B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx \\ &= \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))} \rho(x,t_1)\,dx + \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx. \end{aligned} \tag{E.17}$$

Moreover, (E.14) and the definition of T_ε^* yeild that the right-hand side of (E.17) is greater than $2(1-\varepsilon)A_\rho$. Hence, we obtain

$$1 > 2(1-\varepsilon)A_\rho = 2(1-\varepsilon)\left(\frac{1}{2}+\frac{\eta}{2}\right) = 1+\eta-\varepsilon(1+\eta),$$

so that we have

$$\varepsilon > \frac{\eta}{1+\eta}, \tag{E.18}$$

which (E.18) contradicts our choice of $\varepsilon < \dfrac{\eta}{1+\eta}$. Therefore (E.15) holds valid.

Now, for our $\varepsilon\in(0,\frac{\eta}{1+\eta})$, we define a path $\gamma_\varepsilon^*\colon[T_\varepsilon,T_\varepsilon^*]\to\mathbb{R}^d$ by

$$\gamma_\varepsilon^*(t) = \begin{cases} y_\varepsilon(T_\varepsilon) & \text{if} \quad T_\varepsilon \le t < T_\varepsilon^* - \theta_\varepsilon(T_\varepsilon^*), \\ y_\varepsilon(T_\varepsilon^*) + \dfrac{T_\varepsilon^*-t}{\theta_\varepsilon(T_\varepsilon^*)}\,(y_\varepsilon(T_\varepsilon)-y_\varepsilon(T_\varepsilon^*)) & \text{if} \quad T_\varepsilon^*-\theta_\varepsilon(T_\varepsilon^*) \le t \le T_\varepsilon^*, \\ y_\varepsilon(T_\varepsilon^*) & \text{if} \quad T_\varepsilon^* \le t < T_\varepsilon^*+\theta_\varepsilon(T_\varepsilon^*). \end{cases} \tag{E.19}$$

Clearly we have $\gamma_\varepsilon^* \in C([T_\varepsilon, T_\varepsilon^* + \theta_\varepsilon(T_\varepsilon^*)]; \mathbb{R}^d)$. Moreover, it follows from (E.15) that

$$B_{3R_\varepsilon}(\gamma_\varepsilon^*(t)) \supset B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*)) \cup B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon)) \quad \text{for all } t \in [T_\varepsilon, T_\varepsilon^* + \theta_\varepsilon(T_\varepsilon^*)] \tag{E.20}$$

which, together with the definition of T_ε^*, yields that

$$\begin{aligned} \int_{|x-\gamma_\varepsilon^*(t)|\le 3R_\varepsilon} \rho(x,t)\,dx &\ge \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cup B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t)\,dx \\ &\ge (1-\varepsilon)\,A_\rho \qquad \text{for all } t \in [T_\varepsilon, T_\varepsilon^* + \theta_\varepsilon(T_\varepsilon^*)]. \end{aligned} \tag{E.21}$$

Here, we have enlarged the radius of the ball centered at the $\gamma_\varepsilon^*(t)$ for $t \in [T_\varepsilon, T_\varepsilon^* + \theta_\varepsilon(T_\varepsilon^*)]$.

We shall extend the path γ_ε^* to the whole interval $[T_\varepsilon, T_m)$. We recall (E.11), so that we have from Lemma E.1 that, for all $\tau \in (T_\varepsilon^*, T_m)$, there exists $y_\varepsilon(\tau)$, and exists a constant $\theta_\varepsilon(\tau) > 0$ depending only on d, q, ρ, ε and τ such that

$$\int_{|x-y_\varepsilon(\tau)|\le R_\varepsilon} \rho(x,t)\,dx > (1-\varepsilon)\,A_\rho \quad \text{for all } t \in (\tau - \theta_\varepsilon(\tau), \tau + \theta_\varepsilon(\tau)) \subset (T_\varepsilon^*, T_m) \tag{E.22}$$

Now we put $I_\tau := (\tau - \theta_\varepsilon(\tau), \tau + \theta_\varepsilon(\tau))$. Then, we have

$$\bigcup_{\tau \in (T_\varepsilon^*, T_m)} I_\tau = (T_\varepsilon^*, T_m),$$

so that $\{I_\tau\}_{\tau \in (T_\varepsilon^*, T_m)}$ is an open covering of (T_ε^*, T_m). Since (T_ε^*, T_m) is a Linderöf space, we can take a countable subcovering $\{I_{\tau_k}\}_{k\in\mathbb{N}}$, where $\{\tau_k\}_{k\in\mathbb{N}}$ is some increasing sequence in (T_ε^*, T_m) such that $\tau_k < \tau_{k+1}$, arranging that $I_{\tau_k} \cap I_{\tau_{k+2}} = \emptyset$ for $k = 1, 2, \cdots$. We note that one can take $y_\varepsilon(\tau_1) = y_\varepsilon(T_\varepsilon^*)$.

We define $\gamma_\varepsilon := \gamma_\varepsilon^*$ on $[T_\varepsilon, T_\varepsilon^*]$. If necessary, we make an analogous procedure as in constructing γ_ε^* in (E.19) to define γ_ε on $[T_\varepsilon^*, T_m)$: by writing $I_{\tau_k} \cap I_{\tau_{k+1}} = (a_k, b_k)$ and $y_k := y_\varepsilon(\tau_k)$,

$$\gamma_\varepsilon(t) = \begin{cases} y_k & \text{if} \quad t \in I_{\tau_k} \setminus (a_k, b_k), \\ y_{k+1} + \dfrac{b_k - t}{b_k - a_k}(y_k - y_{k+1}) & \text{if} \quad t \in (a_k, b_k), \\ y_{k+1} & \text{if} \quad t \in I_{\tau_{k+1}} \setminus (a_k, b_k). \end{cases}$$

Then, $\gamma_\varepsilon \colon [T_\varepsilon, T_m) \to \mathbb{R}^d$ is continuous and satisfies that

$$\int_{|x-\gamma_\varepsilon(t)|\le 3R_\varepsilon} \rho(x,t)\,dx \ge (1-\varepsilon)A_\rho \quad \text{for all } t \in [T_\varepsilon, T_m),$$

since we have

$$(B_{R_\varepsilon}(y_k) \times \{t\}) \cap (B_{R_\varepsilon}(y_{k+1}) \times \{t\}) \ne \emptyset \quad \text{for all } t \in (a_k, b_k).$$

□

Remark E.2. *Here, we remark that: if $A_\rho = 1$, the proof becomes easier. Then the following estimate holds: for all $\varepsilon > 0$,*

$$\int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cap B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx > 1 - 2\varepsilon \quad \textit{for all } t \in (T_\varepsilon^* - \theta_\varepsilon(T_\varepsilon^*), T_\varepsilon^*]\ . \tag{E.23}$$

Indeed, we have from (E.14) and the definition of T_ε^* *that*

$$\begin{aligned}
1 &\geq \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cup B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx \\
&= \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))} \rho(x,t_1)\,dx + \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx \\
&\qquad\qquad - \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cap B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx \\
&> 2\,(1-\varepsilon) - \int_{B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon^*))\cap B_{R_\varepsilon}(y_\varepsilon(T_\varepsilon))} \rho(x,t_1)\,dx
\end{aligned}$$

for all $t \in (T_\varepsilon^* - \theta_\varepsilon(T_\varepsilon), T_\varepsilon^*]$ *, which gives (E.23). Thus we do not need to enlarge the radius of the ball centered at the* $\gamma(t)$ *for* $t \in [T_\varepsilon, T_m)$*.*

F Variational problems

We give the proofs of the variational problems, Proposition 1.1 and Proposition 1.2.

We begin with the proof of Proposition 1.1.

Proof of Proposition 1.1. We first prove the relation (1.27):

$$N_1^{\frac{p-1}{2}} = \left(\frac{2}{d}\right)^{\frac{p-1}{2}} \left\{\frac{d(p-1)}{(d+2)-(d-2)p}\right\}^{\frac{1}{4}\{(d+2)-(d-2)p\}} N_2.$$

Take any $f \in H^1(\mathbb{R}^d)\setminus\{0\}$ with $\mathcal{K}(f) \leq 0$ and put $f_\lambda(x) = \lambda^{\frac{2}{p-1}} f(\lambda x)$ for $\lambda > 0$. Then, one can easily verify that:

$$\|f_\lambda\|_{\widetilde{H}^1}^2 = \lambda^{\frac{4}{p-1}+2-d}\left\{\frac{p-(1+\frac{4}{d})}{p-1}\right\}\|\nabla f\|_{L^2}^2 + \lambda^{\frac{4}{p-1}-d}\|f\|_{L^2}^2, \tag{F.1}$$

$$\mathcal{K}(f_\lambda) = \lambda^{\frac{4}{p-1}+2-d}\mathcal{K}(f) \leq 0, \tag{F.2}$$

$$\mathcal{N}_2(f_\lambda) = \mathcal{N}_2(f). \tag{F.3}$$

Moreover, an elementary calculus shows that $\|f_\lambda\|_{\widetilde{H}^1}^2$ takes the minimum at

$$\lambda = \left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{2}} \frac{\|f\|_{L^2}}{\|\nabla f\|_{L^2}}, \tag{F.4}$$

so that

$$\min_{\lambda>0}\|f_\lambda\|_{\widetilde{H}^1}^{p-1} = \left(\frac{d}{2}\right)^{\frac{p-1}{2}} \left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{4}\{d+2-(d-2)p+4\}} \mathcal{N}_2(f). \tag{F.5}$$

Now, we consider a minimizing sequence $\{f_n\}_{n\in\mathbb{N}}$ of the variational problem for N_2 (see (1.20)). Then, it follows from (F.5) and the definition of the variational value N_1 (see (1.19)) that

$$N_1^{\frac{p-1}{2}} \leq \left(\frac{d}{2}\right)^{\frac{p-1}{2}} \left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{4}\{d+2-(d-2)p+4\}} N_2. \tag{F.6}$$

On the other hand, considering a minimizing sequence $\{f_n\}_{n\in\mathbb{N}}$ of the variational problem for N_1, we obtain

$$N_1^{\frac{p-1}{2}} \geq \left(\frac{d}{2}\right)^{\frac{p-1}{2}} \left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{4}\{d+2-(d-2)p+4\}} N_2. \tag{F.7}$$

This inequality (F.7) together with (F.6) gives (1.27).

Next, we prove (1.28):

$$N_3 = \frac{d(p-1)}{2(p+1)} N_2.$$

Let $\{f_n\}_{n\in\mathbb{N}}$ be a minimizing sequence of the variational problem for N_3 (see (1.21)). For each f_n, there is a constant $s_n > 0$ such that $\mathcal{K}(s_n f_n) = 0$. We put $g_n = s_n f_n$. Then, one can easily verify that

$$\mathcal{I}(f_n) = \mathcal{I}(g_n),$$

so that

$$\lim_{n\to\infty} \mathcal{I}(g_n) = N_3. \tag{F.8}$$

Moreover, using $\mathcal{K}(g_n) = 0$, we find that

$$\mathcal{I}(g_n) = \frac{d(p-1)}{2(p+1)} \mathcal{N}_2(g_n) \geq \frac{d(p-1)}{2(p+1)} N_2 \quad \text{for all } n \in \mathbb{N}. \tag{F.9}$$

Hence, (F.8) and (F.9) give us that

$$\frac{d(p-1)}{2(p+1)} N_2 \leq N_3. \tag{F.10}$$

Now, we consider a minimizing sequence $\{f_n\}_{n\in\mathbb{N}}$ of the variational problem for N_2 (see (1.20)). Then, it follows from $\mathcal{K}(f_n) \leq 0$ and the definition of N_3 that

$$N_3 \frac{2(p+1)}{d(p-1)} \|\nabla f_n\|_{L^2}^2 \leq N_3 \|f_n\|_{L^{p+1}}^{p+1} \leq \|f_n\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \|\nabla f_n\|_{L^2}^{\frac{d}{2}(p-1)}. \tag{F.11}$$

Dividing the both sides of (F.11) by $\frac{2(p+1)}{d(p-1)}\|\nabla f_n\|_{L^2}^2$, we have

$$N_3 \leq \frac{d(p-1)}{2(p+1)} \|f_n\|_{L^2}^{p+1-\frac{d}{2}(p-1)} \|\nabla f_n\|_{L^2}^{\frac{d}{2}(p-1)-2} = \frac{d(p-1)}{2(p+1)} \mathcal{N}_2(f_n). \tag{F.12}$$

Since $\{f_n\}_{n\in\mathbb{N}}$ is a minimizing sequence of the variational problem for N_2, taking $n \to \infty$ in (F.12), we obtain

$$N_3 \leq \frac{d(p-1)}{2(p+1)} N_2,$$

which together with (F.10) gives (1.28). □

Next, we give the proof of Proposition 1.2.

Proof of Proposition 1.2. We first solve the variational problem for

$$N_1 = \inf\left\{ \|f\|_{\widetilde{H}^1}^2 \mid f \in H^1(\mathbb{R}^d) \setminus \{0\},\ \mathcal{K}(f) \leq 0 \right\}.$$

Considering a concrete function, we verify that

$$N_1 \lesssim 1, \tag{F.13}$$

where the implicit constant depends only on d and p. Moreover, the functional $\|\cdot\|_{\widetilde{H}^1}$ satisfies that

$$\|f\|_{H^1}^2 \le \frac{d(p-1)}{d(p-1)-4}\|f\|_{\widetilde{H}^1}^2 \quad \text{for all } f \in H^1(\mathbb{R}^d). \tag{F.14}$$

We take a minimizing sequence $\{f_n\}_{n\in\mathbb{N}}$ for this problem, so that

$$\lim_{n\to\infty}\|f_n\|_{\widetilde{H}^1}^2 = N_1, \tag{F.15}$$

$$\mathcal{K}(f_n) \le 0 \quad \text{for all } n \in \mathbb{N}. \tag{F.16}$$

Here, the property (F.16) is equivalent to the inequality

$$\frac{2(p+1)}{d(p-1)}\|\nabla f_n\|_{L^2}^2 \le \|f_n\|_{L^{p+1}}^{p+1} \quad \text{for all } n \in \mathbb{N}. \tag{F.17}$$

By (F.13), (F.14) and (F.15), we obtain the uniform bound of $\{f_n\}_{n\in\mathbb{N}}$ in $H^1(\mathbb{R}^d)$: there exists $C_1 > 0$ depending only on d and p such that

$$\sup_{n\in\mathbb{N}}\|f_n\|_{H^1} \le C_1. \tag{F.18}$$

Moreover, by the Gagliardo-Nirenberg inequality (we can obtain this inequality without solving the variational problem for N_3) and (F.18), we have

$$\|f_n\|_{L^{p+1}}^{p+1} \lesssim C_1^{p+1-\frac{d}{2}(p-1)}\|\nabla f_n\|_{L^2}^{\frac{d}{2}(p-1)}, \tag{F.19}$$

where the implicit constant depends only on d and p. This inequality (F.19), together with (F.17), yields that

$$\frac{2(p+1)}{d(p-1)} \lesssim C_1^{p+1-\frac{d}{2}(p-1)}\|\nabla f_n\|_{L^2}^{\frac{d}{2}(p-1)-2}. \tag{F.20}$$

Hence, using (F.17) again, we have that: there exists $C_2 > 0$ depending only on d and p such that

$$\inf_{n\in\mathbb{N}}\|f_n\|_{L^{p+1}} \ge C_2. \tag{F.21}$$

The properties (F.18) and (F.21) enable us to apply Lemma B.1, so that we obtain

$$\mathcal{L}^d\left[|f_n| \ge \delta\right] > C$$

for some constants C and $\delta > 0$ independent of n. Moreover, applying Lemma B.2, we find that: there exists $y_n \in \mathbb{R}^d$ such that, putting $\widetilde{f_n}(x) = f_n(x+y_n)$, we have

$$\mathcal{L}^d\left(\left[\left|\widetilde{f_n}\right| \ge \frac{\delta}{2}\right] \cap B_1(0)\right) > C' \tag{F.22}$$

for some constant $C' > 0$ independent of n. Here, we can easily verify that this sequence $\{\widetilde{f}_n\}_{n\in\mathbb{N}}$ has same properties as the original one $\{f_n\}_{n\in\mathbb{N}}$:

$$\lim_{n\to\infty} \|\widetilde{f}_n\|_{\widetilde{H}^1}^2 = N_1, \tag{F.23}$$

$$\mathcal{K}(\widetilde{f}_n) \le 0 \quad \text{for all } n \in \mathbb{N}, \tag{F.24}$$

$$\sup_{n\in\mathbb{N}} \|\widetilde{f}_n\|_{H^1} \le C_1. \tag{F.25}$$

We apply Lemma B.3 to $\{\widetilde{f}_n\}_{n\in\mathbb{N}}$ and obtain a subsequence $\{\widetilde{f}_n\}_{n\in\mathbb{N}}$ (still denoted by the same symbol) and a nontrivial function $Q \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n\to\infty} \widetilde{f}_n = Q \quad \text{weakly in } H^1(\mathbb{R}^d). \tag{F.26}$$

The property (F.26) also gives us that

$$\begin{aligned} \|\nabla\widetilde{f}_n\|_{L^2}^2 - \|\nabla\widetilde{f}_n - \nabla Q\|_{L^2}^2 - \|\nabla Q\|_{L^2}^2 &= 2\Re\int_{\mathbb{R}^d} (\nabla\widetilde{f}_n - \nabla Q)\overline{\nabla Q}\, dx \\ &\to 0 \qquad \text{as } n \to \infty, \end{aligned} \tag{F.27}$$

and

$$\|Q\|_{\widetilde{H}^1}^2 \le N_1. \tag{F.28}$$

Here, (F.28) follows from the lower continuity in the weak topology and (F.23). Moreover, Lemma B.4, together with (F.26), gives us that: for all $2 \le q < 2^*$,

$$\|\widetilde{f}_n\|_{L^q}^q - \|\widetilde{f}_n - Q\|_{L^q}^q - \|Q\|_{L^q}^q \to 0 \quad \text{as } n \to \infty. \tag{F.29}$$

It follows from (F.27) and (F.29) that

$$\mathcal{K}(\widetilde{f}_n) - \mathcal{K}(\widetilde{f}_n - Q) - \mathcal{K}(Q) \to 0 \quad \text{as } n \to \infty \tag{F.30}$$

and

$$\|\widetilde{f}_n\|_{\widetilde{H}^1}^2 - \|\widetilde{f}_n - Q\|_{\widetilde{H}^1}^2 - \|Q\|_{\widetilde{H}^1}^2 \to 0 \quad \text{as } n \to \infty. \tag{F.31}$$

We show that the function Q is a minimizer of the variational problem for N_1. To this end, it suffices to prove that

$$\mathcal{K}(Q) \le 0. \tag{F.32}$$

Indeed, by the definition of N_1, (F.32) implies that $N_1 \le \|Q\|_{\widetilde{H}^1}$, which, together with (F.28), yields that

$$\|Q\|_{\widetilde{H}^1}^2 = N_1. \tag{F.33}$$

We prove (F.32) by contradiction: suppose that $\mathcal{K}(Q) > 0$. Then, (F.24) and (F.30) imply that

$$\mathcal{K}(\widetilde{f}_n - Q) \le 0 \qquad \text{for all sufficiently large } n \in \mathbb{N}.$$

Therefore, by the definition of N_1, we have

$$\|\widetilde{f}_n - Q\|_{\widetilde{H}^1}^2 \ge N_1 \qquad \text{for all sufficiently large } n \in \mathbb{N}. \tag{F.34}$$

Combining (F.31), (F.23) and (F.34), we obtain that $\|Q\|_{\widetilde{H}^1} = 0$, which is a contradiction. Hence, we have proved (F.32).

Now, it follows from (F.23), (F.26) and (F.33) that

$$\begin{aligned}\|\widetilde{f}_n - Q\|_{\widetilde{H}_1}^2 &= \|\widetilde{f}_n\|_{\widetilde{H}^1}^2 - \frac{2\left\{p-\left(1+\frac{4}{d}\right)\right\}}{p-1}\Re\int_{\mathbb{R}^d} \nabla\widetilde{f}_n(x)\overline{\nabla Q(x)}\,dx \\ &\qquad - 2\Re\int_{\mathbb{R}^d}\widetilde{f}_n(x)\overline{Q(x)}\,dx + \|Q\|_{\widetilde{H}^1}^2 \\ &\to 0 \qquad \text{as } n\to\infty.\end{aligned}$$

This, together with (F.14), immediately yields the strong convergence:

$$\lim_{n\to\infty}\widetilde{f}_n = Q \ \text{ strongly in } H^1. \tag{F.35}$$

We shall prove

$$\mathcal{K}(Q) = 0. \tag{F.36}$$

Putting $Q_s = sQ$ for $s\in\mathbb{R}$, we have that

$$\mathcal{K}(Q_s) > 0 \quad \text{for all } s \in \Bigg(0,\ \left\{\tfrac{2(p+1)}{d(p-1)}\tfrac{\|\nabla Q\|_{L^2}^2}{\|Q\|_{L^{p+1}}^{p+1}}\right\}^{\frac{1}{p-1}}\Bigg).$$

Here, (F.32) implies that

$$\left\{\frac{2(p+1)}{d(p-1)}\frac{\|\nabla Q\|_{L^2}^2}{\|Q\|_{L^{p+1}}^{p+1}}\right\}^{\frac{1}{p-1}} \le 1.$$

Supposing the undesired situation $\mathcal{K}(Q) < 0$ ($\mathcal{K}(Q)\le 0$ has been proved already), we have by the intermediate value theorem that there exists $s_0\in(0,1)$ such that $\mathcal{K}(Q_{s_0}) = 0$, so that $\|Q_{s_0}\|_{\widetilde{H}^1}^2 \ge N_1$ by the definition of N_1. However, it follows from (F.33) that

$$\|Q_{s_0}\|_{\widetilde{H}^1}^2 = s_0^2\|Q\|_{\widetilde{H}^1}^2 < \|Q\|_{\widetilde{H}^1}^2 = N_1,$$

which is a contradiction. Hence, (F.36) holds valid.

Next, we consider the variational problem for N_2. We show that the function Q obtained above is a minimizer of this problem. Put $Q_\lambda(x) = \lambda^{\frac{2}{p-1}}Q(\lambda x)$ for $\lambda > 0$. We easily verify that

$$\mathcal{K}(Q_\lambda) = 0 \quad \text{for all } \lambda > 0,$$

which implies that

$$\|Q_\lambda\|_{\widetilde{H}^1}^2 \ge N_1 \quad \text{for all } \lambda > 0.$$

Then, the same argument as (F.4), together with (F.33), shows that $\|Q_\lambda\|_{\widetilde{H}^1}\colon (0,\infty)\to[0,\infty)$ takes the minimum at

$$\lambda = \left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{2}}\frac{\|Q\|_{L^2}}{\|\nabla Q\|_{L^2}} = 1. \tag{F.37}$$

Therefore, as well as (F.5), we have that

$$\begin{aligned}N_1^{\frac{p-1}{2}} &= \|Q\|_{\widetilde{H}^1}^{p-1} = \|Q_\lambda\|_{\widetilde{H}^1}^{p-1}|_{\lambda=1} \\ &= \left(\frac{d}{2}\right)^{\frac{p-1}{2}}\left(\frac{d(p-1)}{d+2-(d-2)p}\right)^{\frac{1}{4}\{d+2-(d-2)p+4\}}\mathcal{N}_2(Q).\end{aligned}$$

This, together with (1.27) in Proposition 1.1, leads us to the conclusion that

$$\mathcal{N}_2(Q) = N_2. \tag{F.38}$$

Here, we remark that (F.36) and (F.37) yield the relation (1.33):

$$\|Q\|_{L^2}^2 = \frac{d+2-(d-2)p}{d(p-1)}\|\nabla Q\|_{L^2}^2 = \frac{d+2-(d-2)p}{2(p+1)}\|Q\|_{L^{p+1}}^{p+1}.$$

We shall show that Q is also a minimizer of the variational problem for N_3. Indeed, (1.28) in Proposition 1.1, together with (F.36) and (F.38), yields that

$$\begin{aligned} \mathcal{I}(Q) &= \frac{\|Q\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)}}{\|Q\|_{L^{p+1}}^{p+1}} \\ &= \frac{d(p-1)}{2(p+1)}\mathcal{N}_2(Q) = \frac{d(p-1)}{2(p+1)}N_2 = \mathcal{N}_3. \end{aligned} \tag{F.39}$$

Finally, we prove that Q satisfies the equation (1.16) with $\omega = 1$:

$$\Delta Q - Q + |Q|^{p-1}Q = 0. \tag{F.40}$$

Since $\mathcal{I}(Q)$ is the critical value of $\mathcal{I}$ (see (F.39)), we have

$$\begin{aligned} 0 &= \frac{d}{d\varepsilon}\mathcal{I}(Q+\varepsilon\phi)\bigg|_{\varepsilon=0} \\ &= \frac{(p+1-\frac{d}{2}(p-1))\|Q\|_{L^2}^{p-1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)}\Re\int Q\overline{\phi}dx}{\|Q\|_{L^{p+1}}^{p+1}} \\ &\quad + \frac{\frac{d}{2}(p-1)\|Q\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)-2}\Re\int \nabla Q\overline{\nabla\phi}dx}{\|Q\|_{L^{p+1}}^{p+1}} \\ &\quad - \frac{(p+1)\|Q\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)}\Re\int |Q|^{p-1}Q\overline{\phi}dx}{\|Q\|_{L^{p+1}}^{2(p+1)}} \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}^d). \end{aligned} \tag{F.41}$$

Combining (F.41) with (F.36) ($\|Q\|_{L^{p+1}}^{p+1} = \frac{2(p+1)}{d(p-1)}\|\nabla Q\|_{L^2}^2$), we obtain that

$$\begin{aligned} 0 &= \frac{d^2(p-1)^2}{4(p+1)}\left(\frac{2(p+1)}{d(p-1)} - 1\right)\|Q\|_{L^2}^{p-1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)-2}\Re\int_{\mathbb{R}^d} Q(x)\overline{\phi(x)}\,dx \\ &\quad + \frac{d^2(p-1)^2}{4(p+1)}\frac{\|Q\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)-2}}{\|\nabla Q\|_{L^2}^2}\Re\int_{\mathbb{R}^d}\nabla Q(x)\overline{\nabla\phi(x)}\,dx \\ &\quad - \frac{d^2(p-1)^2}{4(p+1)}\frac{\|Q\|_{L^2}^{p+1-\frac{d}{2}(p-1)}\|\nabla Q\|_{L^2}^{\frac{d}{2}(p-1)-2}}{\|\nabla Q\|_{L^2}^2}\Re\int_{\mathbb{R}^d}|Q(x)|^{p-1}Q(x)\overline{\phi(x)}\,dx. \end{aligned} \tag{F.42}$$

This, together with (F.37), shows that Q satisfies the equation (F.40) in a weak sense. Moreover, it turns out that Q has the following properties: Q is positive (see [35]), radially symmetric (see [19]), unique up to the translations and phase shift (see [32]), and satisfies the decay estimate (see [4, 35]):

$$|\partial^\alpha Q(x)| \le Ce^{-\delta|x|} \quad \text{for all multi-index } \alpha \text{ with } |\alpha| \le 2, \tag{F.43}$$

where C and δ are some positive constants.

Finally, we shall show that Q belongs to the Schwartz space $\mathcal{S}(\mathbb{R}^d)$. Since Q is a smooth and radially symmetric solution to (F.40), we have that

$$\frac{d^2}{dr^2}\partial^\alpha Q = \partial^\alpha Q + \partial^\alpha Q^p - \partial^\alpha \frac{d-1}{r}\frac{d}{dr}Q \quad \text{for all multi-index } \alpha \text{ and } r = |x|. \tag{F.44}$$

Then, an induction argument, together with (F.43), leads to that $Q \in \mathcal{S}(\mathbb{R}^d)$. □

Acknowledgments

The authors would like to express their deep gratitude to Professor Kenji Nakanishi for pointing out a mistake in the earliest version of this paper. The authors also thank Doctor Takeshi Yamada for his valuable comments.

H. Nawa is partially supported by the Grant-in-Aid for Scientific Research (Challenging Exploratory Research # 19654026) of JSPS.

References

[1] Bahouri, H. and Gérard, P., High frequency approximation of solutions to critical nonlinear wave equations. Amer. J. Math. **121** (1999), 131–175.

[2] Bégout, P., Necessary conditions and sufficient conditions for global existence in the nonlinear Schrödinger equation. Adv. Math. Sci. Appl. **12** (2002), 817–827.

[3] Berestycki, H. and Cazenave, T., Instability des etats stationnaires dans les equations de Schrodinger et de Klein Gordon non lineaires. [Instability of stationary states in nonlinear Schrodinger and Klein Gordon equations] C. R. Acad. Sci. Paris Ser. I Math. **293** (1981), 489–492.

[4] Berestycki, H. and Lions, P.L., Nonlinear scalar field equations I, II. Archi. Rat. Mech. Anal. **82** (1983), 313–376.

[5] Bourgain, J., Scattering in the energy space and below for 3D NLS, J. Anal. Math. **75** (1998), 267-297.

[6] Bergh, J., Interpolation Spaces, Springer-Verlag, Berlin Heidelberg New York 1976.

[7] Brézis, H. and Coron, J.M., Convergence of $H-$system or how to blow bubbles. Archi. Rat. Mech. Anal. **89** (1985) 21–56.

[8] Brézis, H. and Lieb, E.H., A relation between pointwise convergence of functions and convergence of functionals. Proc. Amer. Math. Soc. **88** (1983) 485–489. MR0699419 (84e:28003)

[9] Brézis, H. and Lieb, E.H., Minimum action solutions of some vector field equations. Commun. Math. Phys. **96** (1984) 97–113. MR0765961 (86d:35045)

[10] Carles, R., Critical nonlinear Schrödinger equations with and without harmonic potential. Math. Models Methods Appl. Sci. **12** (2002), 1513–1523.

[11] Carles, R. and Nakamura,Y., Nonlinear Schrödinger equations with Stark potential. Hokkaido Math. J. **33** (2004), 719–729.

[12] Cazenave, T., Semilinear Schrödinger equations. Courant Lecture Notes in Mathematics, **10**. New York university, Courant Institute of Mathematical Sciences, New York (2003) .

[13] Cazenave, T. and Weissler, F.B., The Cauchy problem for the critical nonlinear Schödinger equation in H^s, Nonlinear Anal. **14** (1990), 807–836.

[14] Colliander, J., Keel, M., Staffilani, G., Takaoka, H. and Tao, T., Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in $\mathbf{R}^3$,

[15] Cycon, H.L., Froese, R.G., Kirsch, W. and Simon, B., Schrödinger operators and application to quantum mechanics and global geometry, Texts and Monographs in Physics, Springer-Verlag, Berlin (1987).

[16] Duyckaerts, T., Holmer, J. and Roudenko, S., Scattering for the non-radial 3D cubic nonlinear Schrödinger equation. Math. Res. Lett. **15** (2008), 1233–1250.

[17] Foschi, D., Inhomogeneous Strichartz estimates, J. Hyperbolic Differ. Equ. **2** (2005), 1–24.

[18] Fröhlich, J., Lieb, E.H. and Loss, M., Stability of coulomb systems with magnetic fields I, The one- electron atom. Commun. Math. Phys. **78** (1989) 160–190.

[19] Gidas, B., Ni, W.M. and Nirenberg, L., Symmetry of positive solutions of nonlinear elliptic equations in $\mathbf{R}^n$. Mathematical analysis and applications, Part A, Adv. in Math. Suppl. Stud., 7a, Academic Press, New York-London, (1981) 369–402.

[20] Glassey, R.T., On the blowing up solution to the Cauchy problem for nonlinear Schrödinger equations. J. Math. Phys. **18** (1977) 1794–1797.

[21] Ginibre, J. and Velo, G., On a class of Schrödinger equations. I, II: The Cauchy problem, a general case, J. Funct. Anal. **32** (1979) 1–71.

[22] Ginibre, J. and Velo, G., Scattering theory in the energy space for a class of nonlinear Schödinger equations. J. Math. Pures Appl. (9) **64** (1985), 363–401.

[23] Holmer, J. and Roudenko, S., A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation, Comm. Math. Phys. **282** (2008), 435–467.

[24] Holmer, J. and Roudenko, S., On blow-up solutions to the 3D cubic nonlinear Schrödinger equation. Appl. Math. Res. Express. AMRX 2007, no. 1, Art. ID abm004, 31 pp.

[25] Ibrahim, S., Masmoudi, N. and Nakanishi, K., Scattering threshold for the focusing nonlinear Klein-Gordon equation, preprint, arXiv:1001.1474.

[26] Kato, T., On nonlinear Schrödinger equations, Ann. Inst. Henri Poincaré, Physique Theorique **46** (1987), 113–129.

[27] Kato, T., Nonlinear Schrödinger equations. Schrödinger operators (Sonderborg, 1988), 218–263, Lecture Notes in Phys., **345**, Springer, Berlin, 1989.

[28] Kato, T., On nonlinear Schrödinger equations. II. H^s-solutions and unconditional well-posedness, J. Anal. Math. **67** (1995), 281–306. MR1383498 (98a:35124a)

[29] Keel, M. and Tao, T., Endpoint Strichartz estimates. (English summary) Amer. J. Math. **120** (1998), 955–980.

[30] Kenig, C.E. and Merle, F., Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. Invent. Math. **166** (2006), 645–675.

[31] Keraani, S., On the defect of compactness for the Strichartz estimates of the Schrödinger equations. J. Differential Equations **175** (2001), 353–392.

[32] Kwong, M.K., Uniqueness of positive solutions of $\triangle u - u + u^p = 0$ in $\mathbf{R}^N$. Archi. Rat. Mech. Anal. **105** (1989) 243–266.

[33] Leggett, A.J., Bose-Einstein condensation in the alkali gases. some fundamental concepts, Rev. Mod. Phys. **73** (2001) 307–356.

[34] Lieb, E.H., On the lowest eigenvalue of Laplacian for the intersection of two domains. Invent. Math. **74** (1983) 441–448.

[35] Lieb, E.H. and Loss, M., Analysis. Graduate Studies in Mathematics, 14. American Mathematical Society (1997).

[36] Lin, J.E. and Strauss, W.A., Decay and scattering of solutions of a nonlinear Schrödinger equation. J. Funct. Anal. **30** (1978), 245–263.

[37] Lions, P.L., The concentration compactness principle in the calculus of variations. The locally compact case. I, II. Inst. Henri Poincaré, Analyse Nonlinéaire. **1** (1984) 109–145, 223–283.

[38] Lions, P.L., The concentration compactness principle in the calculus of variations. The limit case. I, II. Riv. Math. Iberoamericana **1** (1985) 45–121, 145–201.

[39] Merle, F. and Raphaël, P., Blow up of the critical norm for some radial L^2 super critical nonlinear Schrödinger equations, Amer. J. Math. **130** (2008), 945–978.

[40] Nakanishi, K., Energy scattering for nonlinear Klein-Gordon and Schrödinger equations in spatial dimensions 1 and 2. J. Funct. Anal. **169** (1999), 201–225.

[41] Nawa, H., "Mass concentration" phenomenon for the nonlinear Schrödinger equation with the critical power nonlinearity II. Kodai Math. J. **13** (1990) 333–348.

[42] Nawa. H., Formation of singularities in solutions of the nonlinear Schrödinger equation, Proc. Japan Acad. Ser. A Math. Sci. **67** (1991), 29–34.

[43] Nawa, H., Asymptotic profiles of blow-up solutions of the nonlinear Schrödinger equation. Singularities in fluids, plasmas and optics (Heraklion, 1992), 221–253, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 404, Kluwer Acad. Publ., Dordrecht, 1993.

[44] Nawa, H., Limiting profiles of blow-up solutions of the nonlinear Schrödinger equation with critical power nonlinearity. Proc. Japan Acad. Ser. A Math. Sci. **73** (1997), 171–175.

[45] Nawa, H., Asymptotic profiles of blow-up solutions of the nonlinear Schrödinger equation with critical power nonlinearity. J. Math. Soc. Japan **46** (1994) 557–586.

[46] Nawa, H., Asymptotic and Limiting Profiles of Blowup Solutions of the Nonlinear Schrödinger Equations with Critical Power, Comm. Pure Appl. Math. **52** (1999), 193-270.

[47] Niederer, U., The maximal kinematical invariance groups of Schrödinger equations with arbitrary potentials. Helv. Phys. Acta, **47** (1974) 167–172.

[48] Ogawa, T. and Tsutsumi, Y., Blow-up of H^1-solution for the nonlinear Schrödinger equation. J. Differential Equations **92** (1991) 317–330.

[49] Ogawa, T. and Tsutsumi, Y., Blow-up of H^1-solution for the one dimensional nonlinear Schrödinger equation with critical power nonlinearity. Proc. Amer. Math. Soc. **111** (1991) 487–496.

[50] Sattinger, D. H., On global solution of nonlinear hyperbolic equations. Arch. Rational Mech. Anal. **30** (1968) 148–172.

[51] Strauss, W.A., Existence of solitary waves in higher dimensions. Commun. Math. Phys. **55** (1977) 149–162.

[52] Stubbe, J., Global solutions and stable ground states of nonlinear Schrödinger equations, Physica D **48** (1991) 259–272.

[53] Sulem, C. and Sulem, P-L., The Nonlinear Schrödinger Equation, Self-focusing and wave collapse, Applied Mathematical Sciences, **139**, Springer-Verlag, New York (1999).

[54] Tao, T., Nonlinear dispersive equations. Local and global analysis. CBMS Regional Conference Series in Mathematics, 106 (2006).

[55] Weinstein, M.I., Nonlinear Schrödinger equations and sharp interpolation estimates. Commun. Math. Phys. **87** (1983) 511–517.

[56] Zhang, J., Sharp conditions of global existence for nonlinear Schrödinger and Klein-Gordon equations. Nonlinear Anal. **48** (2002), Ser. A: Theory Methods, 191–207.

Takafumi Akahori,
Graduate School of Science and Engineering
Ehime University,

2-5 Bunkyo-cho, Matsuyama, 790-8577, Japan
E-mail: akahori@math.sci.ehime-u.ac.jp

Hayato Nawa,
Division of Mathematical Science, Department of System Inovation
Graduate School of Engineering Science
Osaka University,
Toyonaka 560-8531 , JAPAN
E-mail: nawa@sigmath.es.osaka-u.ac.jp